



DUAL HIV-1/M+O INFECTIONS AND HIV-1/MO RECOMBINANTS IN FRANCE FROM 2004 TO 2014

Pierre Cappy, Fabienne de Oliveira, Veronique Lemée, Jean-Louis Gaillard, Laurence Bocquet, Jean-Dominique Poveda, Magali Bouvier, Anne Maillard, Thomas Mourez, Jean-Christophe Plantier

► To cite this version:

Pierre Cappy, Fabienne de Oliveira, Veronique Lemée, Jean-Louis Gaillard, Laurence Bocquet, et al.. DUAL HIV-1/M+O INFECTIONS AND HIV-1/MO RECOMBINANTS IN FRANCE FROM 2004 TO 2014. CROI, Feb 2015, Seattle, United States. hal-02115306

HAL Id: hal-02115306

<https://normandie-univ.hal.science/hal-02115306>

Submitted on 30 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pierre Cappy^{*1}, Fabienne De Oliveira¹, Véronique Lemée¹, Jean-Louis Gaillard², Laurence Bocquet³, Jean-Dominique Poveda⁴, Magali Bouvier⁵, Anne Maillard⁶, **Thomas Mourez¹**, Jean-Christophe Plantier^{1*}.

1. Virology Unit, CHU Rouen, France ; 2. Microbiology Unit, Amproise Paré Hospital, AP-HP, France; 3. Virology unit, CHU Lille, France; 4. Laboratoire CERBA, France; 5. Virology Unit, Henri Mondor Hospital, APHP, France ; 6. Virology Unit, CHU Rennes, France ;

*corresponding authors : pierre.cappy@chu-rouen.fr; jean-christophe.plantier@univ-rouen.fr

BACKGROUND

HIV-1 exhibits a high genetic diversity. It is currently divided into 4 groups: M, N, O and P as shown on figure 1^[1]. Recombination is a crucial mechanism of HIV evolution, and HIV-M recombinants now account for more than 20% of HIV-1 infections^[2]. HIV-M and HIV-O co-circulation in Cameroon led in few patients to dual infections and to subsequent generation of MO recombinants (MOrecs). The first dual M+O infections were described in Benin in 1995^[3] and 3 MOrecs were identified in Cameroon in 1999^[4,5] and 2004^[6]. Due to historical links with Western Central Africa, sporadic cases of superinfection/dual infections^[7,8] and MO recombinant infection^[9] were also described in France.

The aim of this study was to determine the circulation level of dual M+O Infections and associated MO recombinant forms in the last decade in France.




Figure 1. Genetic diversity of HIV-1¹

RESULTS

The presence of **both M and O genomes** were detected in **10/141** (7%) HIV-O infected patients.

The analysis of these cases allowed us to identify different situations:

- **2 M+O dual infections** in 2 patients (**RBF140** in 2004 and **RBF216** in 2009),

- **3 MO recombinants associated** :

- **with the HIV-O parental strain** in 2 epidemiologically linked patients (**BCF174**, 2005 and **BCF212**, 2006)

- **with both the M and O parental strains** in 1 patient (**RBF235**, 2007)

- **6 MO recombinants without parental strains** :

- **RBF208** (2008), **RBF221** (2 recombinants in the same patient, 2010), **RBF222**, (2010), **RBF237** (2013), **BCF204** (2013)




Figure 2. Genomic patterns of HIV MO recombinants detected in France






Figure 3. Breakpoints found *in vivo* in MOrecs in France (blue arrows) and in previous publications (orange arrows)

METHODS

1. The presence of both M and O genomes were looked for HIV-M+O / HIV-MO screening in **141** HIV-O patient samples collected from 2004 to 2014 by the RESO/French HIV-National Reference Centre



2. Regions amplified by group specific PCR



Conclusions

Although all described in Cameroonian native patients, we show 11 HIV-M+O dual infections and/or HIV-MO recombinant infections in the past decade in France, highlighting these phenomena are not so rare. Moreover, the presence of MO recombinants circulating without the presence of parental O or M strains, and in epidemiologically linked patients demonstrate they are fit enough to be transmitted and spread. Finally, as HIV-O are naturally resistant to NNRTIs, the transmission of dual HIV-M+O infections or MO recombinants carrying HIV-O genomic fragments underline the importance of searching for these forms, evaluating their spread dynamics and following the possible emergence of a CRF_MO.

References: 1. Mourez T, et al., CMR, 2013 ; 2. Hemelaar J, et al., AIDS, 2011 ; 3. Heyndrickx L, et al., Lancet, 1995 ; 4. Takehisa J, et al., J Virol, 1999 ; 5. Peeters M et al., J Virol, 1999 ; 6. Yamaguchi, J., et al., AIDS Res Hum Retroviruses, 2004 ; 7. Plantier JC, et al., AIDS, 2004 ; 8. Brand, D., et al., AIDS, 2004 ; 9. re A, et al., AIDS, 2010

The 2 viruses found in the same patient RBF221 exhibited multiple breakpoints in the accessory genes (at least 2 and 5 respectively).