



HAL
open science

Evolution of HIV-1 groups M and O: genetic comparative analysis of 23 HIV-1/MO inter-group recombinant forms

Alice Moisan, Fabienne de Oliveira, Pierre Cappy, Paul-Alain Ngoupo, Richard Njouom, Jean-Christophe Plantier

► **To cite this version:**

Alice Moisan, Fabienne de Oliveira, Pierre Cappy, Paul-Alain Ngoupo, Richard Njouom, et al.. Evolution of HIV-1 groups M and O: genetic comparative analysis of 23 HIV-1/MO inter-group recombinant forms. IAS, Jul 2017, Paris, France. <hal-02115882>

HAL Id: hal-02115882

<https://normandie-univ.hal.science/hal-02115882v1>

Submitted on 30 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Evolution of HIV-1 groups M and O: genetic comparative analysis of 23 HIV-1/MO inter-group recombinant forms

A. Moisan^{1,2}, F. De Oliveira¹, P. Cappy^{1,2,3}, P.A. Ngoupo^{2,4}, R. Njouom⁴, J.C. Plantier^{1,2}

¹Charles Nicolle University Hospital, Virology, Rouen, France - ²Normandy University, Groupe de Recherche sur l'Adaptation Microbienne (GRAM2.0), Rouen, France - ³Institut de biologie moléculaire et cellulaire (IBMC), CNRS:FR1589, Strasbourg, France - ⁴Centre Pasteur du Cameroun, Virology, Yaounde, Cameroon

Background

HIVs are characterized by a **high genetic diversity**, due to their simian origins and their replication mode, enhanced by recombination events. Despite the great genetic divergence between pandemic **HIV-1/M** and **HIV-1/O** (endemic in Cameroon), **dual infections** between the two groups can **generate HIV-1/MO inter-group recombinants**. Only three cases were described until 2004 [1-3]. Emergence of such mosaic forms may have a strong impact on diagnosis and treatment due to the presence of highly divergent group O fragments in their genome. Between 2006 and 2017, the implementation of a **sero-molecular algorithm in Cameroon** and the **exploration of atypical HIV-1 infection reactivities** at HIV diagnosis or during follow-up in **France** allowed the description of **new HIV-1/MO infections** [4-6].

Aim

Analyze and compare the **recombination patterns** and **genetic peculiarities** of all HIV-1/MO recombinants currently characterized.

Materials and methods

- **Collection of available data** on infected patients and on all recombinant forms characterized between 1999 and 2017: geographic origin of patients, partial/whole genome sequencing, possible co-infection with parental strains.
- **Comparison of frequencies of their characteristics**: phylogenetic analysis, breakpoints numbers in each mosaic genome and location.

Results

- In addition to the first three cases described, **20 new recombinants or putative forms** were identified in Cameroon and France between 2006 and 2017. Thereby, **23 distinct, partially or completely sequenced, recombinant forms**, found in **22 patients**, were analyzed and compared, revealing a complexity of patterns [Fig.1].
- This work also highlighted a **variable degree of complexity** in profiles, with on average **two to three breakpoints** per recombinant genome [Fig.1].
- For the 23 MO recombinants, the **presence of HIV-1/M and/or HIV-1/O parental strains** was searched and results showed:
 - 12 recombinants with **no parental strains**
 - 10 recombinants **associated to one or two parental strains**
 - one case of **superinfection**, which led to the emergence of a recombinant form.
- Phylogenetic analysis showed that the 23 recombinant forms corresponded to **19 distinct unique recombinant forms (URFs)** [Fig.2]:
 - two URFs in two pairs of patients without any epidemiological link (BCF174/BCF212 and RBF222/RBF243 [6]),
 - one URF in two married patients (REC003/REC024 [4]),
 - one URF described by two different research teams, likely in the same patient (97CA.MP645MO [1]/YBF298),
 - two URFs in the same patient (RBF221-1 and RBF221-2),
 - no known link for the 13 other URFs.




Fig. 1 Genomic profiles of the 23 described recombinants
Fragments belonging to HIV-1/O and /M are shown in blue and red respectively. Codes of recombinants described in 1999 and 2004 [1-4] are written in green and those of recombinants described between 2006 and 2017 in France and in Cameroon [4-6] are written in gray.

- These 23 recombinants were all found in **patients of Cameroonian origin** and genetic diversity of the mosaic fragments **matched the molecular epidemiology in Cameroon**. Indeed, a clear predominance of **HIV-1/M CRF02_AG** (53%) and of **HIV-1/O sub-group H** (87%) was observed [Fig.3].

- Analysis of the genomic profiles and of the breakpoints frequency in the 23 described recombinant forms revealed [Fig.4]:

- **hotspots in the "central" accessory genes** (*vif*, *vpr*, *vpu*), **reverse transcriptase**, **gp41** and **LTRs**
- **no recombination event in protease**, **gp120** and **nef**.




Fig. 2 Phylogenetic relationships of the HIV-1/MO fully sequenced Neighbour-joining trees of the PR-RT [899bp] (a) and gp41 [443bp] (b) regions were constructed with HIV-1/O and /M reference subset.




Fig. 3a Distribution of subtypes of HIV-1/M of the 23 described recombinants




Fig. 3b Distribution of sub-groups of HIV-1/O of the 23 described recombinants [7]

Conclusions

- Despite the complexity of the recombination patterns, these recombinants could be detected with a **suitable sero-molecular strategy** or **following discordances** between detection and/or follow-up of results.
- The phenomenon of MO recombination is rare but **not anecdotal**, and **numerous situations** have been observed (single recombinants, or associated with one or two parental strains, or even a case of superinfection)
- **Inter-groups MO recombination hot/coldspot gradients** and the **location of breakpoints** were identified and remain to be compared with those in intra-group M recombinants.
- The analysis showed that recombinants were often single, with no parental strains, suggesting an **advantageous phenotype** of recombinant forms and a **possible better fitness** relative to parental strains, as observed by Peeters *et al.* [1].

In conclusion, this work allowed us to describe and analyze a unique series of HIV-1/MO recombinants, **highlighting a greater diversity and complexity than originally supposed**, and offering new research perspectives on the **conditions of emergence and virological properties of these recombinant forms**. Moreover, these multiple URFs-MO, single and transmitted in pairs of patients, could lead to **the diffusion and the emergence of a CRF_MO**.

References

- Peeters M, Liegeois F, Torimiro N, Bourgeois A, Mpoudi E, Vergne L, et al. Characterization of a highly replicative intergroup M/O human immunodeficiency virus type 1 recombinant isolated from a Cameroonian patient. *J Virol* 1999;73:7368-7375.
- Takehisa J, Zekeng L, Ido E, Yamaguchi-Kabata Y, Mboudjeka I, Harada Y, et al. Human immunodeficiency virus type 1 intergroup (M/O) recombination in Cameroon. *J Virol* 1999;73:6810-6820.
- Yamaguchi J, Bodelle P, Vallari AS, Coffey R, McArthur CP, Schochetman G, et al. HIV infections in northwestern Cameroon: identification of HIV type 1 group O and dual HIV type 1 group M and group O infections. *AIDS Res Hum Retroviruses* 2004;20:944-957.
- Ngoupo PA, Sadeuh-Mba SA, De Oliveira F, Ngono V, Ngono L, Tchendjou P, et al. First evidence of transmission of an HIV-1 M/O intergroup recombinant virus. *AIDS* 2016;30:1-8.
- Vessiere A, Leoz M, Brodard V, Stracy C, Lemeze V, Depatureaux A, et al. First evidence of a HIV-1 M/O recombinant form circulating outside Cameroon. *AIDS* 2010;24:1079-1082.
- NRCI data
- Leoz M, Feyertag F, Kfutwah A, Mauclere P, Lachenal G, Damond F, et al. The Two-Phase Emergence of Non Pandemic HIV-1 Group O in Cameroon. *PLoS Pathog* 2015;11:e1005029.