

First Description of Azole-Resistant Aspergillus fumigatus Due to TR 46 /Y121F/T289A Mutation in France

Rose-Anne Lavergne, Florent Morio, Loïc Favennec, Stéphane Dominique, Jacques Meis, Gilles Gargala, Paul E Verweij, Patrice Le Pape

▶ To cite this version:

Rose-Anne Lavergne, Florent Morio, Loïc Favennec, Stéphane Dominique, Jacques Meis, et al.. First Description of Azole-Resistant Aspergillus fumigatus Due to TR 46 /Y121F/T289A Mutation in France. Antimicrobial Agents and Chemotherapy, 2015, 59 (7), pp.4331-4335. 10.1128/AAC.00127-15 . hal-02317942

HAL Id: hal-02317942 https://normandie-univ.hal.science/hal-02317942v1

Submitted on 16 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. AAC Accepted Manuscript Posted Online 27 April 2015 Antimicrob. Agents Chemother. doi:10.1128/AAC.00127-15 Copyright © 2015, American Society for Microbiology. All Rights Reserved.

- 1 First description of azole-resistant Aspergillus fumigatus due to TR₄₆/Y121F/T289A mutation
- 2 in France
- 3
- 4 Rose-Anne Lavergne,^{a,b} Florent Morio, ^{a,b} Loïc Favennec,^c Stéphane Dominique,^d Jacques F.
- 5 Meis,^{e,f} Gilles Gargala,^c Paul E. Verweij,^f Patrice Le Pape^{a,b}#
- 6
- Département de Parasitologie et Mycologie Médicale, Université de Nantes, Nantes 7 Atlantique Universités, EA1155-IICiMed, Faculté de Pharmacie, Nantes, France^a; 8 Laboratoire de Parasitologie-Mycologie, Institut de Biologie, CHU de Nantes, France^b; 9 EA3800, Laboratoire de Parasitologie-Mycologie, Université de Rouen et CHU de Rouen, 10 France^c; Department of Pneumology, Cystic Fibrosis Center, Rouen University Hospital, 11 France^d; Department of Medical Microbiology and Infectious Diseases, Canisius Wilhelmina 12 Hospital, Nijmegen, The Netherlands^e; Department of Medical Microbiology, Radboud 13 University Medical Center, Nijmegen, The Netherlands^f 14 15
- 16 Running Head: TR46 resistant Aspergillus fumigatus in France
- 17
- 18 #Address correspondence to Pr Patrice Le Pape, patrice.le-pape@univ-nantes.fr
- 19
- 20 R-A.L and F.M. contributed equally to this work.

21	Abstract: Azole resistance in Aspergillus fumigatus is an emerging public health concern.
22	Recently, a novel fungicide-driven mutation in the cyp51A gene and its promoter,
23	$TR_{46}/Y121F/T289A$, leading to high-level resistance to voriconazole has been identified in
24	the Netherlands, Belgium, Germany, Denmark, Tanzania and India in both clinical and
25	environmental samples. Here we report the first description of A. fumigatus carrying this
26	mutation in France, in a cystic fibrosis patient, underlining the need for extensive monitoring
27	of Aspergillus resistance.

2

untimicrobial Agents and Chemotherapy

Azole-resistant Aspergillus fumigatus are increasingly reported in Europe since the late 2000s. 28 This emerging public health concern occurs through two distinct routes of acquisition: in vivo 29 selection of resistance as a consequence of long-term azole treatment or de novo acquisition 30 of a resistant isolate directly from the environment, linked to the widespread use of azole 31 32 fungicides in agriculture. Besides the TR₃₄/L98H mutation in the cyp51A gene first described in the Netherlands, a novel fungicide-driven mutation, $TR_{46}/Y121F/T289A$ has been recently 33 identified. Up to now, the $TR_{46}/Y121F/T289A$ mutation has been reported in both 34 environmental and/or clinical samples in four countries across Europe (1-7), in Asia (8) and 35 more recently in Africa (9), suggesting a large geographical spread. Here we provide the first 36 37 description of A. fumigatus carrying TR₄₆/Y121F/T289A mutation in France, in a cystic fibrosis patient. 38

39

A 23-year-old male cystic fibrosis patient with follow-up at the Pneumology Department at 40 Rouen University Hospital, France, was seen in consultation in March 2014. This patient had 41 42 high levels of total IgE and Aspergillus-specific IgE with positive Aspergillus-specific IgG antibodies, being suggestive of diagnosis of allergic bronchopulmonary aspergillosis. He had 43 a history of A. fumigatus airway colonization and mold-active azoles exposure since 2002 44 (itraconazole and voriconazole). At the time of the consultation, he was treated with 45 46 voriconazole. Mycological cultures of the sputum collected during the consultation grew A. fumigatus (strain 1). Species identification was obtained by both macroscopic and 47 microscopic characteristics on Sabouraud's agar medium together with sequencing of the 48 beta-tubulin gene (10). According to a local research protocol aiming at the surveillance of 49 azole resistance, this isolate was tested for antifungal susceptibility by the Etest[®] method 50 51 (bioMérieux, Marcy-l'Etoile, France). Unexpectedly, this strain exhibited a high-level 52 resistance to voriconazole (MIC>32µg/mL) in contrast with itraconazole (MIC=8µg/mL) and

posaconazole (MIC=1µg/mL). Antifungal susceptibility was confirmed by the EUCAST broth 53 microdilution reference method (Table 1) (11, 12). Nucleotide sequencing of the cyp51A gene 54 and its promoter, using previously described primers (13, 14) and home-designed primers 55 (CYP51AF-F1: 5'-ATTTCCCTCATCACTGCAA, CYP51AF-R1: 5'-56 5'-AGAAGCGAGATGCTGCTCAT, 57 CATCATGTGCGCAATCTCTT, CYP51AF-F2: CYP51AF-R2: 5'-CCTTTGAAGTCCTCGATGGT), showed the TR₄₆/Y121F/T289A 58 mutation. Antifungal therapy was therefore switched to posaconazole in April 2014 then to 59 caspofungin (50 mg per day) in July 2014 because of pulmonary exacerbation. 60

Given these findings, we retrospectively analyzed all A. fumigatus strains previously isolated 61 62 from this patient since 2007 (n=13), for itraconazole and voriconazole susceptibility, cyp51A sequencing and microsatellite genotyping. As shown in Table 1, our patient had already been 63 colonized by a TR₄₆/Y121F/T289A isolate one year before, in February 2013 (strain 3). All 64 remaining isolates collected before February 2013 were azole-susceptible being either wild-65 type for the *cyp51A* gene or carrying mutations previously found in both azole-resistant and 66 67 azole-susceptible isolates (15). As a single colony was subjected to in vitro susceptibility testing other azole-resistant isolates could have been missed. Microsatellite genotyping was 68 performed using a panel of nine short tandem repeats as described previously (16). As 69 illustrated in Table 1, both TR₄₆/Y121F/T289A isolates from our patient had the same 70 71 genotype as a strain previously isolated in Germany (7) (Table 1). To gain further insights into the route of acquisition of this azole-resistant isolate in our patient, we conducted an 72 environmental study by soil samplings next to the patient's home as described previously 73 (17), as well as surface samplings (contact agar plates) in his office. Neither A. fumigatus 74 carrying TR₄₆/Y121F/T289A nor TR₃₄/L98H was identified. 75

76

Aspergillus fumigatus isolates carrying the TR₄₆/Y121F/T289A mutation have been first 77 described in December 2009 in the Netherlands (2). Since then, such isolates have been 78 evidenced in three others European countries namely Belgium (1, 5), Germany (3, 7) and 79 Denmark (4) and recently in India (8) and Tanzania (9) (Table 2, Figure 1). Taken together, 80 81 these findings suggest, as discussed previously for $TR_{34}/L98H$ isolates, a large geographical 82 spread of this resistance mechanism. Several lines of evidence indicate that, as for TR₃₄/L98H, TR₄₆/Y121F/T289A has emerged through a fungicide-driven route (18), such 83 isolates being found in both azole-naïve (1, 2, 6, 7) and azole-exposed patients (2, 3, 5, 7) as 84 well as from the environment (2, 8, 9). Here we report the first description of A. fumigatus 85 86 carrying the TR₄₆/Y121F/T289A mutation isolated from a French patient.

Interestingly, our patient organized trips to the Netherlands as a tour operator. For these 87 working purposes, he traveled to Amsterdam in November 2012, three months before the first 88 isolation of the TR₄₆/Y121F/T289A strain from his sputum (February 2013). Moreover he 89 regularly received advertising postal packages from Dutch flower producers which were 90 91 opened in his office. Three hypotheses can explain the route of acquisition of this $TR_{46}/Y121F/T289A$ strain in our patient: (i) inhalation of spores carrying $TR_{46}/Y121F/T289A$ 92 during his trip to the Netherlands (2); (ii) colonization after inhaling spores carrying 93 TR46/Y121F/T289A from his environment in France. Our environmental study conducted 94 95 next to the patient's home, less than 100 km from Belgium (where $TR_{46}/Y121F/T289A$ strains have been recently identified (1)), failed to detect $TR_{46}/Y121F/T289A$ environmental isolates. 96 Nevertheless, environmental isolates carrying this mutation have been recently identified by 97 our team in the same region in France supporting this hypothesis (unpublished data). (iii) The 98 hypothesis is colonization after inhaling A. fumigatus spores carrying 99 last 100 TR₄₆/Y121F/T289A, escaping while opening the packages received from the Netherlands. 101 Though the French strains are genetically indistinguishable from the German and genetically different from Dutch isolates, the route of acquisition in our patient is unclear, as spores probably follow an airborne migration as hypothesized previously for $TR_{34}/L98H$ (19, 20).

The present report now provides evidence that A. fumigatus voriconazole-resistant isolates 104 carrying the TR₄₆/Y121F/T289A mutation can be now isolated from clinical samples in 105 106 France. As observed with $TR_{34}/L98H$, a geographical spread of this resistance mechanism is ongoing across Europe and possibly worldwide. These findings together with the high-level 107 voriconazole resistance of the TR₄₆/Y121F/T289A strains both in vitro and in vivo (1, 2, 4-6) 108 underline the need for intensive investigations to determine its prevalence in both clinical and 109 environmental samples. In line with this, as recommended by the ECDC Technical report 110 111 (18), antifungal susceptibility testing of triazoles should be performed on all clinical A. *fumigatus* isolates before starting antifungal therapy. 112

113

Acknowledgments: We are grateful to all the technicians of the Parasitology and Mycology
Laboratories at Nantes University Hospital and Rouen University hospital for technical
assistance.

117

118 Conflict of interest: PLP received grants from Astellas and Pfizer and speaker's fees from 119 Merck and Gilead. FM received speaker's fees from Gilead and MSD and travel grants from 120 Gilead, MSD, Pfizer and Astellas. JFM received grants from Astellas, Basilea and Merck, has 121 been a consultant to Astellas, Basilea and Merck and received speaker's fees from Merck and 122 Gilead.

123 References

Vermeulen E, Maertens J, Schoemans H, Lagrou K. 2012. Azole-resistant
 Aspergillus fumigatus due to TR46/Y121F/T289A mutation emerging in Belgium, July 2012.
 Euro Surveill 17. (48). pii: 20326.

van der Linden JW, Camps SM, Kampinga GA, Arends JP, Debets-Ossenkopp
 YJ, Haas PJ, Rijnders BJ, Kuijper EJ, van Tiel FH, Varga J, Karawajczyk A, Zoll J,
 Melchers WJ, Verweij PE. 2013. Aspergillosis due to voriconazole highly resistant
 Aspergillus fumigatus and recovery of genetically related resistant isolates from domiciles.
 Clin Infect Dis 57:513-520.

Fischer J, van Koningsbruggen-Rietschel S, Rietschel E, Vehreschild MJ,
 Wisplinghoff H, Kronke M, Hamprecht A. 2014. Prevalence and molecular characterization
 of azole resistance in *Aspergillus spp.* isolates from German cystic fibrosis patients. J
 Antimicrob Chemother 69:1533-1536.

Astvad KM, Jensen RH, Hassan TM, Mathiasen EG, Thomsen GM, Pedersen UG,
 Christensen M, Hilberg O, Arendrup MC. 2014. First detection of TR46/Y121F/T289A
 and TR34/L98H alterations in *Aspergillus fumigatus* isolates from azole-naive patients in
 Denmark despite negative findings in the environment. Antimicrob Agents Chemother
 58:5096-5101.

Montesinos I, Dodemont M, Lagrou K, Jacobs F, Etienne I, Denis O. 2014. New
case of azole-resistant *Aspergillus fumigatus* due to TR46/Y121F/T289A mutation in
Belgium. J Antimicrob Chemother 69:3439-3440.

Kuipers S, Bruggemann RJ, de Sevaux RG, Heesakkers JP, Melchers WJ,
 Mouton JW, Verweij PE. 2011. Failure of posaconazole therapy in a renal transplant patient
 with invasive aspergillosis due to *Aspergillus fumigatus* with attenuated susceptibility to
 posaconazole. Antimicrob Agents Chemother 55:3564-3566.

7. Steinmann J, Hamprecht A, Vehreschild MJ, Cornely OA, Buchheidt D, Spiess B,
Koldehoff M, Buer J, Meis JF, Rath PM. 2015. Emergence of azole-resistant invasive
aspergillosis in HSCT recipients in Germany. J Antimicrob Chemother 70:
doi:10.1093/jac/dku566.

Chowdhary A, Sharma C, Kathuria S, Hagen F, Meis JF. 2014. Azole-resistant
 Aspergillus fumigatus with the environmental TR46/Y121F/T289A mutation in India. J
 Antimicrob Chemother 69:555-557.

Chowdhary A, Sharma C, van den Boom M, Yntema JB, Hagen F, Verweij PE,
 Meis JF. 2014. Multi-azole-resistant *Aspergillus fumigatus* in the environment in Tanzania. J
 Antimicrob Chemother 69:2979-2983.

Balajee SA, Gribskov JL, Hanley E, Nickle D, Marr KA. 2005. *Aspergillus lentulus*sp. nov., a new sibling species of A. fumigatus. Eukaryot Cell 4:625-632.

160 11. Rodriguez-Tudela JL, Alcazar-Fuoli L, Mellado E, Alastruey-Izquierdo A,
161 Monzon A, Cuenca-Estrella M. 2008. Epidemiological cutoffs and cross-resistance to azole
162 drugs in *Aspergillus fumigatus*. Antimicrob Agents Chemother 52:2468-2472.

163 12. European Committee on Antimicrobial Susceptibility Testing. 2014. Antifungal
164 Agents - breakpoint tables for interpretation of MICs version 7.1. Available from:
165 http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/AFST/Antifungal_breakpoi
166 nts_v_7.0.pdf

Morio F, Aubin GG, Danner-Boucher I, Haloun A, Sacchetto E, Garcia-Hermoso
D, Bretagne S, Miegeville M, Le Pape P. 2012. High prevalence of triazole resistance in *Aspergillus fumigatus*, especially mediated by TR/L98H, in a French cohort of patients with
cystic fibrosis. J Antimicrob Chemother 67:1870-1873.

171 14. Alanio A, Sitterle E, Liance M, Farrugia C, Foulet F, Botterel F, Hicheri Y,
172 Cordonnier C, Costa JM, Bretagne S. 2011. Low prevalence of resistance to azoles in

- 173 Aspergillus fumigatus in a French cohort of patients treated for haematological malignancies.
- 174 J Antimicrob Chemother **66**:371-374.
- 175 15. Snelders E, Karawajczyk A, Schaftenaar G, Verweij PE, Melchers WJ. 2010.
- 176 Azole resistance profile of amino acid changes in Aspergillus fumigatus CYP51A based on
- 177 protein homology modeling. Antimicrob Agents Chemother 54:2425-2430.
- 178 16. de Valk HA, Meis JF, Curfs IM, Muehlethaler K, Mouton JW, Klaassen CH.
- 179 2005. Use of a novel panel of nine short tandem repeats for exact and high-resolution
- 180 fingerprinting of *Aspergillus fumigatus* isolates. J Clin Microbiol **43:**4112-4120.
- 2009. Possible environmental origin of resistance of *Aspergillus fumigatus* to medical
 triazoles. Appl Environ Microbiol **75**:4053-4057.

Snelders E, Huis In 't Veld RA, Rijs AJ, Kema GH, Melchers WJ, Verweij PE.

- 184 18. European Center for Disease Prevention and Control. 2013. Technical Report. Risk
 185 assessment on the impact of environmental usage of triazoles on the development and spread
 186 of resistance to medical triazoles in *Aspergillus* species. doi 10.2900/76274
- 187 19. Vermeulen E, Lagrou K, Verweij PE. 2013. Azole resistance in *Aspergillus*188 *fumigatus*: a growing public health concern. Curr Opin Infect Dis 26:493-500.
- 189 20. Chowdhary A, Kathuria S, Xu J, Meis JF. 2013. Emergence of azole-resistant
- 190 Aspergillus fumigatus strains due to agricultural azole use creates an increasing threat to
- human health. PLoS Pathog 9:e1003633. doi:10.1371/journal.ppat.1003633
- 192

17.

181

9

- 193 Table 1- Overview of the characteristics of all Aspergillus fumigatus strains isolated from
- 194 sputum samples of the patient
- 195 Table 2- Literature review of all studies reporting TR₄₆/Y121F/T289A A. fumigatus isolates
- 196 Figure 1- Geographical spread of the TR₄₆/Y121F/T289A resistance mechanism (for each
- 197 strain, the exact location and origin (clinical or environmental) is indicated)
- 198 Figure 2- STRAf dendrogram highlighting the genetic relatedness between Aspergillus
- 199 *fumigatus* isolated from our patient and previously reported TR₄₆/Y121F/T289A isolates


			MIC	s mg/L				
Strain	Reference in the dendrogram	Month/Year of isolation -	(EUCAST)		Cyp51A mutations			
number			ITC	VRC	-			
1	14-105-2468	Mar 2014	8	>8	TR ₄₆ /Y121F/T289A			
2	14-148-2457	Nov 2013	0.5	0.5	Wild-type			
3	14-148-2460	Feb 2013	>8	>8	TR ₄₆ /Y121F/T289A			
4	141428-459	Jan 2013	0.5	1	Wild-type			
5	14-148-2458	Jan 2013	0.25	0.25	Wild-type			
6	14-148-2456	Dec 2010	Not performed		F46Y, G89G, M172V, N248T, D255E, L358L, E427K, C45-			
7	14-148-2455	Sept 2010	Not performed		F46Y, G89G, M172V, N248T, D255E, L358L, E427K, C45-			
8	14-148-2454	Jul 2010	Not performed		F46Y, G89G, M172V, N248T, D255E, L358L, E427K, C45-			
9	Not performed	Jul 2009	Not performed		F46Y, G89G, M172V, N248T, D255E, L358L, E427K, C454			
10	14-148-2450	Jul 2009	0.25	1	Wild-type			
11	14-148-2448	Mar 2009	0.25	1	Wild-type			
12	14-148-2447	Dec 2007	0.25	1	Wild-type			
13	14-148-2445	May 2007	0.5	2	Wild-type			
14	Not performed	Feb 2007	0.5	1	F46Y, G89G, M172V, N248T, D255E, L358L, E427K, C45			

Table 1- Overview of the characteristics of all *Aspergillus fumigatus* strains isolated from sputum samples of the patient

Table 2- Literature review of	of all studies reporting	TR46/Y121F/T289A A.	fumigatus isolates
-------------------------------	--------------------------	---------------------	--------------------

Reference	Date	Type of sample	Underlying condition	Infection	Antifungal susceptibility CLSI (EUCAST*)			Outcome	Country	
					VRC	ITC	POS	_		
(1)	July 2012	BAL fluid	HSCT	Probable IA	>16	4	1	Death	Belgium	
	Dec 2009	Sputum	HSCT	Probable IA	>16	4	0.25	Persistent infection		
	Jan 2010	Ear	Chronic otitis externals/sinusitis	IA	>16	>16	2	Persistent infection		
	Jan 2010	Abdominal abscess	SOT	Proven IA	>16	2	0.5	Death		
	Feb 2010	Sputum	Cystic fibrosis	No IA	>16	4	0.5	Alive		
	Feb 2010	Sputum	Lung carcinoma	No IA	>16	>16	2	Alive		
	March 2010	Sputum	HSCT	Probable IA	>16	1	0.25	Death		
	March 2010	Sputum	Cystic fibrosis, SOT	Proven IA	>16	>16	0.5	Alive		
(2)	May 2010	Biopsy	Chronic otitis, surgery	Proven IA	>16	4	1	Alive	The Netherlands	
	May 2010	Sputum	Lung fibrosis	None	>16	>16	1	Alive		
	June 2010	Sputum	Traumatism	None	>16	1	0.25	Death		
	July 2010	Brain biopsy	Beta thalassemia, diabetes mellitus	Proven IA	>16	4	1	Death		
	Sept 2010	Sputum	Cystic fibrosis	ABPA	>16	2	0.5	Alive		
	Oct 2010	Sputum	COPD, SOT	None	>16	>16	2	Alive		
	Nov 2010	Sputum	COPD	No IA	>16	>16	2	Alive		
	Jan 2011	Sputum	HSCT	Probable IA	>16	>16	1	Death		
	Dec 2009 to Jan 2011	Air sampling			ND	ND	ND			
(8)	2012-2013	Soil sampling			>16	1-2	0.25-0.5		India	
(3)	Sept 2012	Sputum	Cystic fibrosis	Colonization	>8*	>8*	2*	Alive	Germany	
(4)	Jan 2014	Sputum	Bruton's agammaglobulinemia, SOT	Probable IA	>4*	0.25- 0.5*	0.125- 0.25*	Death	Denmark	
(9)		Soil sampling			16->16	1-2	0.25-0.5		Tanzania	
(5)	Nov 2013	BAL fluid	HSCT	Probable IA	>8	>16	1	Death	Belgium	
	Sept 2012	pt 2012 BAL fluid HSCT		Probable IA	16*	>16*	0.5*	Death		
(7)	July 2012	BAL fluid	HSCT	Proven IA	1*	>16*	0.5*	Death	Germany	
Present report	Feb 2013 March 2014	Sputum	Cystic fibrosis	Colonization	>8* >8*	>8* 8*	ND	Alive	France	

POS: posaconazole BAL: bronchoalveolar lavage; HSCT: hematopoietic stem cell transplantation; IA: invasive aspergillosis; SOT: solid organ transplantation; ABPA: allergic bronchopulmonary aspergillosis; COPD: chronic obstructive pulmonary disease; ND: Not determined



AAC

Antimicrobial Agents and Chemotherapy STRA

STRAf

-100 2 38 30 æ 38 g 44 ₽ 4 ş 50 Ģ -70 율 Ģ 10.0 CF/NL1659 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL1859 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL1866 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL2244 TR46/Y121F/T289A 26.0 14.0 The Netherlands 21.0 9.0 33.0 11.0 22.0 8.0 10.0 CF/NL2340 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL2604 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL2992 The Netherlands TR46/Y121F/T289A 26.0 21.0 33.0 11.0 22.0 9.0 8.0 14.0 10.0 CF/NL0682 TR46/Y121F/T289A The Netherlands 25.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL0723 The Netherlands TR46/Y121F/T289A 25.0 21.0 9.0 33.0 11.0 22.0 8.0 14.0 10.0 CF/NL0645 TR46/Y121F/T289A The Netherlands 26.0 26.0 9.0 33.0 11.0 22.0 8.0 14.0 5.0 141428-459 France Wildtype 18.0 12.0 16.0 25.0 11.0 22.0 8.0 8.0 9.0 CF/NL0032 The Netherlands TR46/Y121F/T289A 26.0 21.0 12.0 26.0 9.0 20.0 13.3 9.0 9.0 CF/NL2753 TR46/Y121F/T289A 26.0 21.0 12.0 26.0 9.0 20.0 13.3 9.0 The Netherlands 9.0 3F The Netherlands TR46/Y121F/T289A 26.0 21.0 12.0 26.0 9.0 20.0 13.3 9.0 9.0 CF/NL0840 The Netherlands TR46/Y121F/T289A 26.0 21.0 12.0 26.0 9.0 22.0 13.3 9.0 9.0 CF/NL2889 The Netherlands TR46/Y121F/T289A 26.0 21.0 9.0 21.0 13.3 9.0 12.0 26.0 19.0 14-148-2457 France Wildtype 25.0 21.0 12.0 47.0 9.0 11.0 8.0 10.0 5.0 14-148-2448 France Wildtype 23.0 24.0 14.0 34.0 11.0 20.0 13.0 9.0 5.0 14-148-2450 France Wildtype 23.0 24.0 14.0 34.0 11.0 20.0 13.0 9.0 Wildtype 23.0 24.0 14.0 34.0 11.0 20.0 13.0 9.0 5.0 14-148-2447-1 France 8.0 14-148-2458 France Wildtype 23.0 20.0 11.0 10.0 14.0 37.0 7.0 9.0 20.0 2107 Germany TR46/Y121F/T289A 10.0 20.0 8.0 42.0 9.0 10.0 12.0 10.0 20.0 14-148-2460 France TR46/Y121F/T289A 10.0 42.0 9.0 12.0 20.0 8.0 10.0 10.0 20.0 14-105-2468 TR46/Y121F/T289A France 10.0 20.0 8.0 42.0 9.0 10.0 12.0 10.0 9.0 19B The Netherlands TR46/Y121F/T289A 9.0 20.0 13.0 43.0 9.0 10.0 12.0 9.0 Wildtype 14-148-2447-2 France 15.0 20.0 11.0 18.0 10.0 7.0 15.3 10.0 16.1 14-148-2454 15.0 20.0 11.0 10.0 7.0 15.3 10.0 France Wildtype 18.0 16.1 14-148-2455 Wildtype France 15.0 20.0 11.0 18.0 10.0 7.0 15.3 10.0 8.0 14-148-2456 Wildtype 15.0 20.0 11.0 18.0 10.0 7.0 15.3 10.0 France 5.0 14-148-2445 France Wildtype 15.0 20.0 7.0 15.3 11.0 18.0 10.0 10.0

AAC