

Expression of the translocator protein (TSPO) from Pseudomonas fluorescens Pf0-1 requires the stress regulatory sigma factors AlgU and RpoH

Charlène Leneveu-Jenvrin, Emeline Bouffartigues, Olivier Maillot, Marc Feuilloley, Nicole Orange, Pierre Cornelis, Nathalie Connil, Sylvie Chevalier

▶ To cite this version:

Charlène Leneveu-Jenvrin, Emeline Bouffartigues, Olivier Maillot, Marc Feuilloley, Nicole Orange, et al.. Expression of the translocator protein (TSPO) from Pseudomonas fluorescens Pf0-1 requires the stress regulatory sigma factors AlgU and RpoH. Pseudomonas Conference, 2015, Washington, United States. hal-02366398

HAL Id: hal-02366398 https://normandie-univ.hal.science/hal-02366398

Submitted on 15 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Expression of the translocator protein (TSPO) from *Pseudomonas* fluorescens Pf0-1 requires the stress regulatory sigma factors AlgU and RpoH.

Charlène Leneveu-Jenvrin, Emeline Bouffartigues, Olivier Maillot, Marc G. J. Feuilloley, Nicole Orange, Pierre Cornelis, Nathalie Connil and Sylvie Chevalier

Laboratory of Microbiology Signals and Microenvironment, LMSM EA4312, University of Rouen, France

The translocator protein (TSPO), previously designated as peripheral-type benzodiazepine receptor, is an evolutionary conserved protein that is found in many Eukarya, Archae and Bacteria, in which it plays several important functions including membrane biogenesis, signaling and stress response. A *tspo* homologue gene has been identified in several members of the *Pseudomonas* genus, among which the soil bacterium *P. fluorescens* Pf0-1. In this bacterium, the *tspo* gene is located in the vicinity of a putative hybrid histidine kinase-encoding gene.

Since *tspo* has been involved in water stress related response in plants, we explored the effects of hyperosmolarity and temperature on *P. fluorescens* Pf0-1 *tspo* expression using a strategy based on *lux*-reporter fusions.



We show that the two genes PfI01_2810 and tspo are co-transcribed forming a transcription unit. The expression of this operon is growth phase-dependent and is increased in response to high concentrations of NaCl, sucrose and to a D-cycloserine treatment, which are conditions leading to activity of the major cell wall stress responsive extracytoplasmic sigma factor AlgU. The promoter region activity is strongly lowered in a *P. aeruginosa algU* mutant, suggesting that AlgU may be involved at least partly in the molecular mechanism leading to PfI01_2810-tspo expression. In silico analysis of this promoter region failed to detect an AlgU consensus binding site; however, we detect a putative binding site for the heat shock response RpoH sigma factor. Accordingly, the promoter activity of the region containing this sequence is increased in response to high growth temperature and slightly lowered in a *P. aeruginosa rpoH* mutant strain. Taken together, our data suggest that *P. fluorescens tspo* gene may belong at least partly to the cell wall stress response.