

Erratum

Lipid composition of membranes of *Escherichia coli* by liquid chromatography/tandem mass spectrometry using negative electrospray ionisation

Delphine Oursel¹, Corinne Loutelier-Bourhis^{1*}, Nicole Orange², Sylvie Chevalier², Victor Norris³ and Catherine M. Lange¹

¹Spectrométrie de masse Bio-Organique, CNRS UMR 6014, Université de Rouen, 76821 Mont-Saint-Aignan cedex, France

²Laboratoire de Microbiologie du Froid, UPRES 2123, Université de Rouen, 27000 Evreux, France

³Epigenomics Project, genopole[®], 93 rue Henri-Rochefort, 91000 Evry, France and Assemblages Moléculaires: Modélisation et Imagerie SIMS, UMR CNRS 6522, Université de Rouen, 76821 Mont-Saint-Aignan, France¹

Rapid Commun. Mass Spectrom. 2007; **21**: 1721–1728.

DOI: 10.1002/rcm3013

For both PG and PE, the product ions noted as $[M-H - RCHCO - H_2O]^-$ on page 1724 (paragraph 2) and in Table 1 page 1725 should more probably correspond to

$[M-H - RCH_2CO_2H]^-$ ions which have been previously described in the literature by several authors (references 12 and 15 in the text and additional references 39 and 40).

39. Hsu FF, Turk J. *J. Am. Soc. Mass Spectrom.* 2000; **11**: 892.

40. Hsu FF, Turk J. *J. Am. Soc. Mass Spectrom.* 2001; **12**: 1036.