



Seasonal variations of phytoplankton assemblages in relation to environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species

Benlahcen Rijal Leblad, Rachid Amnhir, Sagou Reqia, Ferdaous Sittel, Mouna Daoudi, Mohamed Marhraoui, Mohamed Karim Ouelad Abdellah, Benoît Véron, Hassan Er-Raioui, Mohamed Laabir

► To cite this version:

Benlahcen Rijal Leblad, Rachid Amnhir, Sagou Reqia, Ferdaous Sittel, Mouna Daoudi, et al.. Seasonal variations of phytoplankton assemblages in relation to environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species. Harmful Algae, 2020, 96, pp.101819. 10.1016/j.hal.2020.101819 . hal-02863649

HAL Id: hal-02863649

<https://normandie-univ.hal.science/hal-02863649>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Seasonal variations of phytoplankton assemblages in relation to environmental factors in Mediterranean coastal waters of Morocco, a focus on HABs species

Benlahcen Rijal Leblad^a, Rachid Amnhir^b, Sagou Reqia^b, Ferdaous Sitel^c, Mouna Daoudi^a,
Mohamed Marhraoui^a, Karim Ouelad Abdellah Mohamed^a, Benoît Veron^d, Hassan Er-Raioui^c and
Mohamed Laabir^{e*}

^a*Institut National de Recherche Halieutique – Laboratoire de Surveillance et Suivi du Milieu
Marin, Centre Régional de Tanger. Km 7, Route Ksar Sghir. Malabata. B.P 5268. Dradeb.
Tanger. 90000*

^b*Institut National de Recherche Halieutique, Route de Sidi Abderrahmane, Casablanca*

^c*Université Abdelmalek Saadi, Faculté Sciences, Tanger*

^d*BioMEA BFA – Université de Caen, Basse-Normandie, Caen*

^e*MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France*

*corresponding author, e-mail: mohamed.laabir@umontpellier.fr

Tel. : +33668689436

Fax : +33467143719

ABSTRACT

Studies on phytoplankton and in particular Harmful Algal Blooms (HABs) species in southern Mediterranean waters are scarce. We performed from April 2008 to June 2009 weekly investigations on microphytoplankton community structure and abundance in two contrasted marine ecosystems located in the western Moroccan Mediterranean coast, M'diq Bay and Oued Laou Estuary. Simultaneously, we measured the main physico-chemical parameters. Globally, the two studied areas showed comparable values of the assessed abiotic environmental factors. Temperature and salinity followed seasonal variation with values ranging from 13.5 °C to 21.4 °C and 31 to 36.8, respectively. Average nutrient values in surface water ranged from 0.7 to 45.76 µM for dissolved inorganic nitrogen, 0.02 to 2.10 µM for PO₄ and 0.23 to 17.46 µM for SiO₄ in the study areas. A total of 92 taxa belonging to 8 taxonomic classes were found. The highest number of microphytoplankton abundance reached 1.2 x 10⁶ cells L⁻¹ with diatoms being the most abundant taxa. Factorial Discriminant Analysis (FDA) and Spearman correlation test showed a significant seasonal discrimination of dominant microphytoplankton species. These micro-organisms were associated with different environmental characteristics, in particular temperature and salinity. Numerous HABs species were encountered regularly along the year. Although *Dinophysis* species and *Prorocentrum lima* were present in both sites, no Lipophilic Shellfish Poisoning was detected in the analyzed bivalve mollusks. Domoic acid (DA), produced by toxic species of *Pseudo-nitzschia* was found with concentrations up to 18 µg DA g⁻¹ in the sweet clam *Callista chione*. Data showed that the observed persistent and dramatic paralytic shellfish poisoning (PSP) intoxication of mollusks resulted probably of *Gymnodinium catenatum* proliferations in both studied areas. Contrary to sweet clam *C. chione*, the cockle *Acanthocardia tuberculatum* showed a permanent and extremely high toxicity level during the 15 months survey with up to 7545 µg Equivalent Saxitoxin kg⁻¹ flesh (ten times higher than the sanitary threshold of 800µg eqSTX Kg⁻¹flesh). The present work highlights for the first time the dynamic of microphytoplankton including HABs species and their associated toxin accumulation

in the commercially exploited shellfish in the southern western Mediterranean waters of Morocco. Furthermore, the acquired data will help us to improve the monitoring of HABs species and related toxins in these coastal marine systems.

Keywords: SW Mediterranean, phytoplankton diversity, environmental factors, HABs, toxins.

1. Introduction

The microalgae community has been shown to respond rapidly to environmental changes (Cloern, 2001; Carstensen et al., 2015). Phytoplankton species composition vary with several environmental factors including physical (irradiance, temperature and turbulence), chemical (inorganic and organic nutrients, oligo-elements, vitamins) and biological (competition and predation) (Boyd et al., 2010; Bužančić et al., 2016). To understand the functioning of a marine ecosystem, it is necessary to study its structure, composition and evolution at time and space scales. Phytoplankton is an important trophic component participating to aquatic ecosystem functioning. Phytoplankton community structure is controlled by various environmental conditions (Gailhard, 2003; Trombetta et al., 2019). Knowing which environmental factor control phytoplankton species development helps to understand their evolution and dynamics. Southern coastal Mediterranean marine ecosystems are knowing along the last decades' important perturbations due to massive urbanization, aquaculture and recreative activities and are under the pressure of climate change (Hallegraef 2010; Macias et al., 2015; 2018; Wells et al., 2015; 2018). These anthropogenic factors could influence phytoplankton and thus ecosystem functioning (Bužančić et al., 2016). The studies on the temporal variation of the community structure of phytoplankton in the Moroccan western Mediterranean waters are scarce (El Madani et al., 2011; Daoudi et al., 2012; Rijal Leblad et al., 2013). Two sites were selected for the present study, M'diq Bay and Oued Laou Estuary holding an important socio-economic activity (fishing, shellfish harvesting and recreative activities). M'diq

79 Bay holds aquaculture activity (mussels and fish) and is characterized by an important shellfish
 80 catching (559 tons' year⁻¹) particularly cockle and clam (Rijal Leblad 2012).

81 In the last decades, Harmful Algal Blooms (HABs) are increasing in frequency, intensity
 82 and geographic distribution impacting human health and aquaculture (Hallegraeff, 1993; Vand
 83 Dolah, 2000). Both northern and southern Mediterranean waters were exposed to these noxious
 84 phenomena (Vila et al. 2001a; 2005; Laabir et al., 2011; Abdenadher et al., 2012; Fertouna-
 85 Benlakhhal et al., 2015; Zmerli-Triki et al., 2016; Abadie et al.; 2019; Ben-Rejeb-Jenhani et al.,
 86 2019). Moroccan marine waters are not spared by episodes of toxic microalgae blooms. Paralytic
 87 Shellfish Poisoning (PSP) events have been recorded in Moroccan Atlantic waters since 1969
 88 causing several intoxications leading some times to human death (Tber, 1983; Bourhilli, 1982;
 89 Taleb et al., 2003). Morocco's Mediterranean waters have long known PSP events which caused
 90 closure of shellfish harvesting for several years (Taleb et al., 2001). These PSP events were
 91 associated to *Gymnodinium catenatum* development (Tahri, 1998; Taleb et al., 2001). *Gymnodinium*
 92 *catenatum* was reported for first time in Spanish Mediterranean waters (Delgado, 1990; Bravo et al.,
 93 1990; Gomez 2003). There were indications of the presence of this species in Tunisian (Dammak-
 94 Zouari et al., 2009) and in Algerian coastal waters (Frehi et al., 2007). However information is still
 95 lacking about the dynamic of this neurotoxic dinoflagellate in the Mediterranean marine coastal
 96 waters. In the Western Mediterranean Sea, the occurrence of *Pseudo-nitzschia* was documented in
 97 the coastal waters of Spain (Quijano-Scheggia et al., 2010; Busch et al., 2016), France (Quiroga
 98 2006; Grzebyk et al., 2017), Italy (Cerino et al., 2005; Ruggiero et al., 2015), Tunisia (Sahraoui et
 99 al., 2009; Melliti Ben Garali et al., 2019), Algerian (Illoul et al., 2008) and Morocco (Rijal leblad et
 100 al., 2013). *Dinophysis* and its toxic compounds were reported only few times in Mediterranean
 101 waters (Aissaoui et al. 2014; Garcia-Altares et al. 2016; Bazzoni et al., 2018).

102 The purpose of this study is to investigate the taxonomic composition and abundance of the
 103 microphytoplankton community in two contrasted areas (M'diq Bay and Oued Laou Estuary) of the
 104 western Mediterranean waters with a focus on HABs species. We also aim to analyze the effect of

the main abiotic environmental factors on microphytoplankton succession and in particular on the potentially toxic microalgae. Another objective was to determine to which extent the observed dramatic PSTs intoxications of the cockles and the clams are related to the presence of paralytic toxin producing species in the water column of the studied ecosystems.

2. Material and Methods

2.1. Study sites

Oued Laou Estuary (Station 1; 35°27' 310 N - 05°05' 06 W) and M'diq Bay (Station 2; 35°41'646 N - 05°19'075 W) (Figure. 1), are located in the western Moroccan coast of Mediterranean Sea adjacent to the Gibraltar Strait. M'diq Bay receives terrestrial inputs through a temporal torrential stream (Rijal Leblad, 2012). The hinterland of this area is characterized by a watershed in rocky mountain knowing a high urbanization. Oued Laou Estuary is less urbanized and receives permanent inputs of fresh water through the Oued Laou River, rich in organic matter. The hinterland of this marine coastal area is characterized by irrigated farming and market gardening.

2.2. Phytoplankton and physico-chemical parameters

Sampling was performed weekly in the water column of S1 and S2 (depth of 7-10 m) from April 2008 to June 2009, except when weather conditions did not allowed sailing. Two seawater samples were collected from each site at subsurface (- 0.5 m) of the water column. The first seawater samples were preserved using Lugol's iodine acidic solution for the identification and quantification of microphytoplankton species. Then samples were settled in 25 mL chamber, stored in the dark at ambient temperature, the settling time was 24 hours and samples were analyzed within the same week of sampling. The microphytoplankton species were identified and quantified using a Leica DMIL inverted photonic microscope (Uthermol, 1958). The second

Sea water sample was stored at -20 °C until chemical analyses. Inorganic nutrient (ammonium NH₄, nitrite NO₂, nitrate NO₃, phosphate PO₄ and silicate SiO₄) concentrations were measured using the protocol of Aminot and Chaussepied (1983) by spectrophotometer (Unico SQ4802 UV/VIS double Beam spectrophotometer model). Temperature, practical salinity and pH were measured *in situ* at each station during sampling period with a multi-parameter probe (WTW) 197i. Practical salinity and pH calibrations were performed monthly (see the protocol in [www.geotechenv.com / Manuals / WTW_Manuals / Multi_197i](http://www.geotechenv.com/Manuals/WTW_Manuals/Multi_197i)).

2.3. Domoic acid analyses

The concentrations of DA were determined by High-performance liquid chromatography (HPLC) (Shimadzu 10vp type). This apparatus is composed of a SCL-10vp Controller, a LC-10ADvp Quaternary Pump, a CTO-10vp Colonne Four, a SIL-10ADvp Autosampler, a SPD-M10Avp Photodiode Array Detector, a Vydac C18 column (250 × 4.6 mm, with 5 mm) and the Guard Cartridge (Vydac C18, 5 mm). DA was analyzed monthly and in periods of high *Pseudo-nitzschia* abundance (> 10⁵ cells L⁻¹). It was measured in the whole meat of cockles and sweet clams according to Quilliam's (1995) protocol. Analyses in triplicates were performed using about 100 g of shellfish meat (ten to fifteen individuals were required to have such amount of meat). After being shredded and homogenated, 4 g of meat were added to 16 mL of solvent extraction (methanol-water, 1:1) and then homogenized (Ultra-Turrax for 3 minutes at about 10,000 rpm). The homogenate was centrifuged at least at 4,000 rpm for 10 min to obtain supernatant. The later was analyzed using the following chromatographic conditions: mobile phase flow rate of 1 mL min⁻¹, detector wave length of 242 nm, injection volume of 20 mL and an oven temperature for the column of 40 °C. The determination of DA content in samples was done with a detection limit of 0.3 mg g⁻¹.

2.4. Paralytic shellfish toxicity analyses using bioassay

Commercially exploited bivalve mollusks as cockle (*Acanthocardia tuberculatum*) and the sweet clam (*Callista chione*) were sampled every two weeks at M'diq Bay and Oued Laou Estuary during 2008 and 2009. The seabed over 10-15 m was raked using a dredge and samples brought back using a net. The samples were put in a cooler with ice blocks until they reach the laboratory. PSP toxicity analysis was carried out by mouse bioassay according to the AOAC method (1990). Briefly, 100 g homogenised tissues were mixed with 100 ml 0.1 M chlorydric acid and boiled for 5 mn, pH adjusted to 2-3 and centrifuged for 15 min at 3000 rpm. One milliliter of the supernatant was injected intraperitoneally to three 20 g Albinos mice. The values reported are expressed in $\mu\text{g STXeq kg}^{-1}$ mollusk meat.

2.5. Lipophilic shellfish poisoning analysis

The analyses the lipophilic toxins (Okadaic Acid-group toxins, Pectenotoxin group and Azaspiracid group) were based on the extraction of these toxins using acetone and dichloromethane (DCM) from shellfish samples. The extracts were washed with dichloromethane (DCM) and water. After evaporation of the DCM, the obtained residue was solubilized with a solution of tween 60 at 1%. One milliliter of the extract was injected into the intraperitoneal cavity of Suisse albino mice weighing between 19 and 21 g. A positive result corresponds to the death of 2 or 3 mice injected during a 24-hour observation period. These analyses were conducted according to the method of EURLMB (EU Harmonised SOP MBA Lipophilic Version 4, Vigo, Spain).

2.6. Statistics

Statistical analyses were performed using the software XLSTAT 2011. We used a Spearman correlation and Factorial Discriminant Analysis (FDA) to highlight if there is any seasonal dissimilarities and describe the main patterns of temporal variability of microphytoplakton taxa in relation to environmental factors.

3. Results

3.1. Physico-chemical parameters

Results of physico-chemical parameters are summarized in Table 1 and shown in Figure 2. The lowest seasonal mean temperature values were recorded in winter (14.32 ± 0.67 °C at S1 and 16.07 ± 0.52 °C at S2) and the highest values in summer (20.12 ± 1.68 °C at S1 and 19.50 ± 1.29 °C at S2) (Fig. 2A). Temperature in M'diq Bay was slightly lower than that of Oued Laou Estuary from April to October 2008. The highest seasonal mean salinity values were recorded in summer (36.54 ± 0.09 at S1 and 36.51 ± 0.05 at S2) while the lowest values were observed in winter during rainfall periods with 33.83 ± 2.05 at S1 and 35.06 ± 0.35 at S2 (Fig. 2B, D). Spring and autumn seemed to be transitional periods. However, we observed an important decrease in salinity (up to 31 in November and December 2008 and up 32-33 in April 2009) in Oued Laou estuary (Fig. 2B). This decrease could be related to heavy rainfall events (Fig. 2D) and the subsequent lower salinity of S1 seawater due to freshwater input from Oued Laou river. Along the seasons, pH was slightly basic and oscillated around 8 for both sites. The lowest values were measured during autumn (7.93-7.86) and winter (7.8-7.97) while the maximum values were observed in spring (8.06-8.12) summer (8.11-8.14) in both stations (Fig. 2C). The rainfall episodes were registered during autumnal and winter periods whereas spring and summer were dry. Rainfall level at S2 was generally greater than that recorded at S1 with the highest value ($258 \text{ mm month}^{-1}$) recorded at S2 at October. Summer was characterized by low or no precipitation.

3.2. Nutrients

Data are summarized in Table 1 and shown in Figure 3. The nutrient concentrations fluctuated widely in function of the seasons and were quite similar between S1 and S2. The seasonal mean values of ammonium (NH_4^+) varied widely and ranged from 0.09 to $43.13 \text{ } \mu\text{M}$. The highest mean values were observed in autumn with $24.53 \pm 10.19 \text{ } \mu\text{M}$ at S1 and $17.74 \pm 14.63 \text{ } \mu\text{M}$ at S2. The

lowest mean values were observed in spring with $0.09 \mu\text{M}$ at S1 and $0.12 \mu\text{M}$ at S2. Nitrite concentrations at S1 and S2 station were relatively low. They ranged between 0 (not detected) to $0.19 \mu\text{M}$. The highest values were recorded in autumn and winter with values up to $0.17 \mu\text{M}$ at S1 and $0.19 \mu\text{M}$ at S2. Nitrate ranged between 0.17 to $7.52 \mu\text{M}$ with an average of $1.79 \pm 1.66 \mu\text{M}$ at S1 and $0.89 \pm 0.66 \mu\text{M}$ at S2. At S1, the highest values were recorded at the end of May 2008 ($5.39 \mu\text{M}$), mid-August 2008 ($3.27 \mu\text{M}$) and during the period of November 2008 - early March 2009 ($7.51 \mu\text{M}$) end of April 2009 ($4.63 \mu\text{M}$) and early June 2009 ($4.02 \mu\text{M}$). During the period of October 2008 - March 2009, we observed an increase in the concentration of nitrate at S1. In early October 2008, nitrate concentration did not exceed $1 \mu\text{M}$. Thereafter, it increased gradually to reach the highest registered values ($7.51 \mu\text{M}$ in March 2009). After, there was a drop at the end of March 2009. At S2, the highest levels were recorded in autumn and winter period. The maximum values, 2.59 and $2.37 \mu\text{M}$, were recorded in November 2008 and the second week of February 2009, respectively, with decreasing trend in spring ($0.72 \pm 0.34 \mu\text{M}$) and summer periods ($0.32 \pm 0.21 \mu\text{M}$).

Silicate concentrations at S2 ($3.39 \pm 2.91 \mu\text{M}$) exceeded those measured at S1 ($2.16 \pm 1.54 \mu\text{M}$), the highest values were observed during spring ($4.10 \pm 3.90 \mu\text{M}$) and autumn ($4.12 \pm 2.05 \mu\text{M}$) at S2. Silicate content displayed several fluctuations, the values ranged between 0.23 to $17.46 \mu\text{M}$, with alternation of enrichment and decreasing period. A gradual decrease in silicate was observed from July to September 2008, and an increase from October until March 2009 reaching a maximum value of $17.46 \mu\text{M}$ in March. Another decrease coincided with the massive development of the diatoms genera *Pseudo-nitzschia* and *Leptocylindrus*. After this bloom, silicate contents peaked during mid-April to June 2009, and showed a decline in June 2009.

Phosphate concentrations varied between 0 and $2.10 \mu\text{M}$. Generally, phosphate concentrations at S2 were slightly higher than those recorded at S1. The highest values were recorded during autumn, with $0.94 \pm 0.83 \mu\text{M}$ at S1 and $0.64 \pm 0.65 \mu\text{M}$ at S2. Low values were measured during spring for S1 $0.16 \pm 0.10 \mu\text{M}$ and winter for S2 $0.25 \pm 0.22 \mu\text{M}$. The monthly value of the Si/P

ratio showed large temporal fluctuations (Fig. 4). The highest values (up to 20) of Si/P ratio were recorded at the M'diq bay during 2008 sampling period and thereafter this ratio remained relatively low. Figure 4 showed the correspondence of the peaks of Si/P ratio and the abundances of diatoms.

3.3. Phytoplankton structure

The phytoplankton community composition and abundance in S1 and S2 showed the presence of different classes: Bacillariophyceae, Dinophyceae, Euglenoidea, Dictyochophyceae, Chlorophyceae, Cyanophyceae, Cryptophyceae and Prymnesiophyceae. However, most recorded species belong to two major classes, Bacillariophyceae and Dinophyceae (Fig. 5 and Fig. 6). The total abundance of phytoplankton (cells L⁻¹) showed important seasonal variations (Fig 7.). The highest values were observed in spring (12.10^5 cells L⁻¹) and summer (12.10^5 cells L⁻¹). In contrast, low abundances were observed in autumn (19.10^4 cells L⁻¹) and winter (71.10^3 cells L⁻¹). The average phytoplankton cell abundance recorded in S2 ($18.10^4 \pm 20.10^4$ cells L⁻¹) was slightly higher than that recorded in S1 ($14.10^4 \pm 17.10^4$ cells L⁻¹). Bacillariophyceae was the dominant group followed by Dinophyceae in both sites (Fig 5, 6). At Oued Laou Estuary (Fig.8), the dominant taxa were: *Heterocapsa* spp., *Skeletonema* spp., *Thalassiosira* spp., *Noctiluca scintillans*, *Nitzschia* spp., *Scrippsiella* sp., *Pseudo-nitzschia* spp., *Guinardia striata*, *Leptocylindrus* spp. *Gymnodinium catenatum*, *Euglena* sp., *Plagioselmis* sp., *Chaetoceros* spp. and *Asterionella glacialis*. Except *Pseudo-nitzschia* spp., *Leptocylindrus* spp. and *Chaetoceros* spp. which showed an average relative densities of 29.5%, 24% and 14.5% respectively, the other taxa were poorly represented in this ecosystem; At M'diq Bay (Fig.9), the dominant taxa were: *Thalassiosira* spp., *Thalassionema nitzschioides*, *Prorocentrum triestinum*, *Nitzschia* spp., *Gyrodinium* spp., *Skeletonema* spp., *Scrippsiella* sp., *Leptocylindrus* spp., *Pseudo-nitzschia* spp., *Plagioselmis* sp., *Chaetoceros* spp., and *Asterionella glacialis*. The relative dominance values of these taxa varied from 0% to 90%. The average of the relative densities of *Leptocylindrus* spp.,

Pseudo-nitzschia spp. and *Chaetoceros* spp. were 24%, 23.5% and 20% respectively. These taxa dominated the microphytoplankton community.

3.4. Harmful Algal Blooms species and mollusks intoxications

Results showed the presence of four *Dinophysis* species (*Dinophysis caudata*, *Dinophysis fortii*, *Dinophysis acuminata* and *Dinophysis rotundata*) with the highest densities (up to 560 cells L⁻¹ when counting all *Dinophysis* species cells) registered between the end of May and mid-September 2008. *Dinophysis* species were mostly present and abundant in Oued Laou Estuary (Fig. 10). There was one important peak of DST producing species *Prorocentrum lima* in Oued Laou (1280 cells L⁻¹) on August the 28th 2008. There was no intoxication of the mollusks with LSP during the 15 months' study. The benthic dinoflagellate *Ostreopsis* was found at 4 occasions during the survey with densities not exceeding 80 cells L⁻¹.

Oued Laou Estuary and M'diq Bay showed the development of species belonging to *Pseudo-nitzschia* genus in spring which is related to Amnesic Shellfish Syndrome (ASP). We showed the presence of DA in two analyzed shellfish species, but the sweet clam (*C. chione*) was so far the most contaminated with up to 15.94 µg DA g⁻¹ mollusk meat in M'diq Bay (April 6th 2009) and 16.88 in Oued Laou (September 10th 2008). The cockle contained low DA with concentrations not exceeding 2.06 µg DA g⁻¹ mollusk meat (April 7th 2009) and 4.73 µg DA g⁻¹ mollusk meat (April 6th 2009) in S1 and S2, respectively (Fig. 11).

Alexandrium spp. registered the highest densities in M'diq Bay with up to 5200 cells L⁻¹ in March 3th 2008. This genus was less present at Oued Laou Estuary with concentrations not exceeding 1480 cells L⁻¹ (March 13th 2008). *G. catenatum* which has been related to PSP syndrome showed almost similar distributions in M'diq Bay (up to 2960 cells L⁻¹) and Oued Laou Estuary (maximum of 3960 cells L⁻¹ in January 22st 2008 (Fig. 12). December 2009 registered also an important development of *G. catenatum* with up to 2040 cells L⁻¹ in M'diq Bay. Permanent and high levels of PSTs contamination were observed in Oued Laou Estuary in the cockles (up to 7545 µg STX equiv Kg meat⁻¹, a value almost 10 times higher than sanitary

threshold). Cockles from M'diq Bay were also permanently contaminated but with concentration up to 3720 $\mu\text{g STX equiv Kg meat}^{-1}$ (registered in January 31th 2008). The sweet clams were less contaminated by PST in both sites with concentrations ranging from 0 to 1945 $\mu\text{g STX equiv Kg meat}^{-1}$ and from 0 to 1230 $\mu\text{g STX equiv Kg meat}^{-1}$, in Oued Laou and M'diq Bay, respectively.

3.5. Correlation between phytoplankton and environmental factors

Statistical analyses were applied to highlight any correlation between environmental parameters and species richness and species diversity of phytoplankton in the studied areas (Table 2). Globally, M'diq Bay and Oued Laou Estuary showed the same trend in their correlations with environmental factors (Table 2). Temperature and salinity were correlated positively to species diversity and species richness. This later, was negatively correlated with nitrate (Table 2). Only salinity and pH showed positive correlation with total phytoplankton (Table 2). Three microphytoplankton groups showed a positive correlation with salinity including Prymnesiophyceae, Cryptophyceae and Dictyophyceae and Diatomophyceae (Table 2). Except Raphidophyceae and Chlorophyceae, all the other the phytoplankton groups were positively correlated with pH (Table 2). DIN, NH_4 and SiO_4 were not correlated with any of the examined phytoplankton variables. Nitrate was negatively correlated with Diatomophyceae, Prymnesiophyceae and Euglenoidea, (Table 2). Dictyophyceae was positively correlated with nitrate and nitrite. Dinophyceae and Diatomophyceae were negatively correlated with nitrite. Factorial analyses related to the season indicated that 92.43% (S1) and 95.22% (S2) of the variance was explained by two axes (F1 and F2) (Fig. 13A and Fig. 13C). In relation to the seasons, the variables showed four groups. For both stations, the temporal variability followed a seasonal cycle with a seasonal transition (Fig. 13B and Fig. 13D). Data showed positive correlations between environmental factors and microphytoplankton species on a seasonal basis. For station 1, the correlations were as follows : in winter (NO_2 , NO_3 and *Thalassionema*

nitzschioides), in spring (SiO_4 and *Nitzschia* spp., *Noctiluca scintillans*, *Scrippsiella* sp., *Pseudo-nitzschia* spp., *Thalassiosira* spp., *Euglena* sp., and *Asterionella glacialis*), in summer (pH, temperature, salinity and *Dinophysis* spp., *Leptocylindrus* spp., *Chaetoceros* spp., *Heterocapsa* spp. and *Skeletonema* spp.) and in autumn (PO_4 , NH_4 and *Guinardia delicatula*, *Guinardia striata*, *Ceratium* spp., *Pleurosigma* spp. and *Gymnodinium catenatum*). For station S2, the obtained positive correlations were in winter (NO_2 , NO_3 and *Thalassiosira* spp.), in spring (SiO_4 , pH, *Alexandrium* spp., *Nitzschia* spp., *Scrippsiella* sp., *Prorocentrum triestinum*, *Pseudo-nitzschia* spp., *Chaetoceros* spp., *Thalassionema nitzschioides*, *Skeletonema* spp., *Eucampia zodiacus*, *Gyrodinium* spp., and *Asterionella glacialis*), in summer (Temperature, Salinity, *Leptocylindrus* spp., *Plagioselmis* sp., *Dinophysis* spp. and *Guinardia striata*) and in autumn (PO_4 , NH_4 and *Gymnodinium catenatum*). Factorial analyses according to the season indicated that the discrimination was dominated by temperature and salinity during summer. Winter was characterized by high levels of nitrate and nitrite. During summer, in both stations a gradual decrease of nitrate was observed and coincided with an increase in water temperature and salinity. Autumn was characterized by high values of ammonium and phosphate whereas winter showed high values of nitrite and nitrate which could be explained by nitrification processes during winter season. The highest phytoplankton biomass and diversity were recorded in spring and summer. Spearman analyses were applied to highlight the correlation between the major phytoplankton species and the environmental factors, this is shown in Tables 3 and 4.

4. Discussion

Phytoplankton distribution and diversity in M'diq Bay and Oued Laou Estuary

The inventory of phytoplankton species in Oued Laou Estuary and M'diq Bay showed the dominance of two groups: Bacillariophyceae and Dinophyceae. This finding is in agreement with previous studies showing that these groups are the main phytoplankton components in the Mediterranean (Vilicic et al. 2002, El Madani 2011, Armi et al., 2010, Daoudi et al., 2012; Salhi et al., 2018; Draredja et al., 2019). In our study, in term of cell abundance, the diatoms

dominated largely with *Pseudo-nitzschia*, *Leptocylindrus* and *Chaetoceros* being the major taxa. Species belonging to Bacillariophyceae showed a high diversity in spring whereas dinoflagellates were more diversified during summer season. This corresponds to previous works on the phytoplankton in the Nador lagoon located in the South Western Mediterranean, Morocco (El Madani et al., 2011; Daoudi et al., 2012). According to Smayda (1984), this seasonal succession is typical of temperate ecosystems. The phytoplankton communities in Oued Laou Estuary and M'diq Bay showed a similar seasonal trend. The phytoplankton mean cell abundance was generally slightly higher in M'diq Bay (18×10^4 cells l^{-1}) when compared to that of Oued Laou (14×10^4 cells l^{-1}). This could be the consequence of the relatively low levels of silicate present in Oued Laou Estuary required by diatoms for their growth. The dominance of diatoms in both ecosystems indicates that the Western Moroccan Mediterranean coast is a water mixing area. Margalef *et al.*, (1979) and Gailhard (2003) suggested that the vertical mixing of water column favors the development of diatoms. Hydrological factors such as mixing with nutrient-rich freshwater during rainfall periods are known to impact the development of phytoplankton in a given ecosystem. Physicochemical parameters of water masses together with climatic conditions particularly air temperature fluctuate seasonally in the Mediterranean temperate ecosystems (Dhib et al., 2013; Laanaia et al., 2013; Salhi et al., 2018; Trombetta et al. 2019). Also, the seasonal variability of phytoplankton communities structure and environmental parameters have been demonstrated at weekly and biweekly time scales in various coastal and estuarine marine waters (Jouenne et al., 2007; Lopes et al., 2007; Armi et al., 2010). In our study, temperature seems to be the most important factor which affect the diversity and richness of phytoplankton species in the two investigated areas. The highest phytoplankton abundance was observed in spring and summer. Temperature is correlated to species richness and diversity (table 2). This is in accordance with the recent work of Trombetta et al., (2019) suggesting that water temperature drives phytoplankton blooms in coastal waters. Phytoplankton diversity and dynamic could be impacted by the observed increase in temperature in the Mediterranean (Shen

et al., 2016; Wells et al., 2015; 2019; Kim et al., 2019). To verify this hypothesis, particularly for HABs species, a monitoring of phytoplankton diversity and dynamics is under way in Oued Laou estuary and Mdiq bay with the aim of comparing these data with those acquired 10 years ago. Diatoms were significantly associated with DIN concentrations in winter and with silicates in autumn. The decreasing concentrations of DIN and NO_3 measured in spring and summer could be associated to the observed increase in dinoflagellates.

In our study, the highest phytoplankton densities were recorded in spring and summer corresponding to the increase of temperature and to high nutrient concentrations which were registered in winter. Less important phytoplankton biomass occurred in winter and coincided with rainfall but with low temperature and light intensity levels. The spring and summer seem to be the most favorable periods for phytoplankton development and diversity. In winter, heavy rainfalls allowed nutrients enrichment both ecosystems, particularly ammonium. This was shown in another Mediterranean ecosystem (Alexandria, Egypt) (Abdel-Halim and Khairy 2007). Ammonium turns into nitrite then into nitrate due to bacterial nitrification. Boutaib et al., (2011) showed an increase in bacterial activity during the wet season (autumn and winter) in Moroccan Western Mediterranean coasts. The decrease in nutrient levels was observed during spring and summer. This reduction was probably related to microalgae development during spring. A significant negative correlation between nutrients and microalgae was observed (Tables 2-4). In Oued Laou Estuary, mineral phosphorus concentration is low and very close to the values provided by Redfield et al. (1963) for Mediterranean waters. In contrast, the phosphorus concentrations in M'diq Bay were relatively important probably due to aquaculture activities and to effluent discharges from the three major cities: Ceuta, F'nideq and M'diq with more than 220000 inhabitants. The silicate nutrient is an essential element for the development of Bacillariophyceae and Dictyochophyceae (Richard, 1987). In both ecosystems, silicates were generally abundant in comparison with other nutrients. Generally inorganic N:P ratios were higher than 16, they were $47.61 (\pm 39.87)$ in S1 and $43.34 (\pm 28.25)$ in S2. Inorganic Si:P ratios

in S1 were lower than 20 (12.81 ± 10.65), however in S2 they were slightly higher than 20 (25.35 ± 28.25). Our results were similar to those of Berland et al. (1973) and Daoudi et al. (2012) who suggested that the phosphorus could be considered as the limiting nutrient in Mediterranean Sea.

HABs species and related toxicity in Mdiq Bay and Oued Laou Estuary

This is the first study on the dynamics of several HABs species developing in South Western Mediterranean waters with measurements of environmental factors. Data showed that *Pseudo-nitzschia* genera developed in M'diq Bay and Oued Laou with up to 35×10^4 cells.L⁻¹ in March 30th 2009 with toxic species resulting of DA intoxication with up to $15.94 \mu\text{g DA g}^{-1}$ mollusk meat In M'diq Bay (April 6th 2009) and $16.88 \mu\text{g DA g}^{-1}$ in Oued Laou (September 10th 2008). The first human intoxication with DA after ingestion of mussels was in 1978 in Al-Hoceima Bay, 190 km away from Oued Laou (Mediterranean coast), where patients who had eaten mussels (*Mytilus galloprovincialis*) suffered from loss of memory and disorientation. *Pseudo-nitzschia* blooms have been documented in Moroccan Mediterranean waters by Rijal Leblad et al. (2013) who showed the presence in M'diq Bay waters of eight toxic species: *P. multistriata*, *P. cuspidata*, *P. galaxiae*, *P. multiseries*, *P. pseudodelicatissima*, *P. pungens* var. *aveirensis*, *P. calliantha* and *P. fraudulenta*.

In our study, the blooms of *Pseudo-nitzschia* followed the increase of the Si/P ratio. Silicates were proven to increase *Pseudo-nitzschia* abundance (Thorel et al., 2017). The correlation between DA production and the nutritional status seems to be complex in natural populations, since they are frequently composed of numerous species (Trainer et al. 2009). Studies on *Pseudo-nitzschia* dynamics, diversity and toxicity remain relatively scarce in the South Mediterranean (Sahraoui et al. 2009; Rijal leblad et al., 2013; Melliti Ben Garali et al., 2019). Here, *Pseudo-nitzschia* correlated to salinity and temperature but negatively to nitrate. In Bizerte Bay, (Mediterranea,Tunisia), Melliti Ben Garali et al., 2019 showed a positive correlation

between *Pseudo-nitzschia* abundance and salinity, silicate, phosphorus and urea. Studies are under way to on the biology and ecology of *Pseudo-nitzschia* species responsible for ASP in these fragile marine ecosystems.

G. catenatum in M'diq Bay and Oued Laou showed moderate abundances of up to 3960 cells l⁻¹. However, extremely high levels of PSTs contamination were observed permanently in both investigated ecosystems in the cockles reaching 7545 µg STX equiv Kg meat⁻¹, a value almost 10 times higher than sanitary threshold. The sweet clams were less contaminated by PSTs in both sites with concentrations ranging from 0 to 1945 µg STX equiv Kg meat⁻¹. A peak of *G. catenatum* abundance (>4000 cells L⁻¹) observed in January 2008 corresponded to a maximum level of PSP in mollusks which was not the case of *Alexandrium* cells. This suggest that *G. catenatum* was the main responsible of the mollusks PSP intoxications. This finding based on a bi-weekly survey during two years corroborated the finding of Tahri (1998) and Taleb et al. (2001) who showed that since 1994 (corresponding to the implementation of toxic species and intoxications monitoring in Morocco, RSSL network), PSP outbreaks have been frequently reported and the causative organism have been identified as *G. catenatum*. This dinoflagellate species have been reported to proliferate in Spanish (Fraga, 1996, Busch et al., 2016), Portuguese (Moita et al., 1998, Silva et al., 2015), Algerian (Frehi et al., 2007) and Tunisian (Dammak-Zouari et al., 2009) marine coastal waters. In our study, *G. catenatum* abundance was correlated with ammonium in Oued Laou Estuary and with nitrate and nitrite in M'diq Bay. According Band-Schmidt et al., (2014) toxin production of *G. catenatum* changed with temperature. The growth rate of this neurotoxic species varies with temperature; the highest growth was obtained between 21 and 24 °C (Band-Schmidt et al., 2010; 2014). In our study, *G. catenatum* was abundant in autumn with temperatures of 18-19 °C. However, it was demonstrated that *G. catenatum* is rather an eurythermal and cosmopolitan species tolerating temperatures from 16 to 33 °C (Band-Schmidt et al., 2014) evolving in Pacific, Atlantic and Indian Oceans but also in Mediterranean Sea.

Our data showed the persistence of high level of PSP toxins during the 15 months' survey in the cockles in M'diq Bay and Oued Laou with very high concentrations ($> 3000 \mu\text{g STX equiv Kg meat}^{-1}$) even when *G. catenatum* densities were low (fig 10). The sweet clam presented high PSP levels only in January and March 2008 corresponding to the highest *G. catenatum* densities in the two studied ecosystems. These results reinforce the hypothesis that *G. catenatum* is responsible for the observed PSP intoxication of the mollusks. Taleb et al. (2001) showed the cockle presents PSP toxicity throughout the year while other mollusks like the sweet clam accumulate it seasonally. Sagou et al. (2005) showed that the cockle sequester PSP toxins preferably in non-visceral organs contrary to sweet clam that sequester them in visceral tissues (digestive gland) which could explain the depuration/retention fate of PSTs. Taleb et al., (2001) attributed the high toxicity levels of cockles to the biotransformation of C-toxins (with low specific toxicity) to dc-carbamoylsaxitoxin (dcSTX) with relatively high toxicity. Rijal Leblad et al., (2017) showed throughout laboratory experiments a partial and progressive elimination of PSP toxins in two investigated mollusks, with a slower elimination kinetic in the tuberculate cockle when compared with the sweet clam, needing 120 and 3 days to reach levels of $80 \mu\text{g SXTeq } 100\text{g}^{-1}$ of meat, respectively.

In our study, *Dinophysis caudata* showed the highest abundance, followed by *Dinophysis fortii*, *Dinophysis acuminata* and *Dinophysis rotundata*. Even though *Prorocentrum lima* is known to be a benthic species, it showed abundances of up to $1300 \text{ cells.L}^{-1}$ in the water column of Oued Laou Estuary during August 2008. The other recorded benthic dinoflagellate *Ostreopsis* showed low abundance not exceeding 90 cells.L^{-1} . The occurrence of benthic HABs species have been well documented in Western Mediterranean (Vila et al., 2001; Dhib et al., 2015; Ben Gharbia et al. 2016; 2019) but not yet in Algerian and Moroccan coastal waters. *D. caudata* was shown to produce Dinophysistoxins, PTX-2 and OA responsible for DSP intoxication (Trainer et al., 2013). DSTs are associated to many species of *Dinophysis* species and *P. lima* (Bazzoni et al., 2018). In our study, using the biological method we noted the absence the detection of DST

intoxication in S1 and S2. However, DSTs related to *Dinophysis* species developing have been shown in NW Mediterranean Sea by Bazzoni et al. (2018) and García-Altare et al. (2016) in some in shellfish farming areas of Italian and Spanish waters, respectively. In our study, *Dinophysis* correlated to pH in both sites. Moreover, correlated to salinity and phosphorus in S1 and S2 respectively.

5. Conclusion

For years, PSP events threat the consumer's health and cause dramatic economic lose due to permanent closure of shellfish harvesting areas of the investigated Mediterranean waters, M'diq Bay and Oued Laou Estuary. Our data clearly showed that *G. catenatum* was responsible of PSTs contamination in the study areas and better explained its relationships with environmental factors. The next step should be to isolate several strains of *G. catenatum* from Moroccan waters, to characterize them at the genetic and toxic levels. Ecophysiological studies should help us to better understand how the environmental factors influence the growth and toxin production of this dinoflagellate. ASP was observed but DA did not exceed 18 µg.g⁻¹ meat which remains below the sanitary threshold. However, *Pseudonitzschia* species have to be isolated and their identity determined genetically in addition to the cellular toxin characterization. Concerning DST, the used biological method have to be completed by direct chemical analyses of lipophilic toxins using HPLC-MS/MS technique which have to be implemented in the monitoring program of toxins. Finally, increasing sampling frequency and long term monitoring is important to better known the driving environmental parameters and among them temperature of the dynamic of phytoplankton including HABs species developing in these fragile and exploited coastal marine ecosystems.

Aknowledgements

We thank the staff of the marine environment monitoring station INRH in Tanger. Thanks to IRD (Institut pour la Recherche et le Développement, France) for supporting Moroccan-French collaboration in HABs studies.

Figures and Tables Legend

Fig. 1. Location of the sampled stations (S1 : Oued Laou Estuaty and S2 : M'diq Bay) in the Moroccan Western Mediterranean sea.

Fig. 2. Temporal variation of Rainfall (mm.month⁻¹), pH, temperature (°C) and Salinity in M'diq Bay and Oued Laou Esturay stations located in the Moroccan Western Mediterranean sea.

Fig. 3. Temporal variation of phosphorus (PO₄), Silicate (SiO₄) and Dissolved Inorganic Nitrogen (DIN) expressed in $\mu\text{mole.l}^{-1}$ at M'diq Bay and Oued Laou Estuary

Fig. 4. Temporal variations of Diatom abundance (Cells.L⁻¹), Si/P Ratio and N/P Ratio at M'diq Bay and Oued Laou Estuary

Fig. 5. Contribution in percentage of different taxa to total phytoplankton abundance at the Oued Laou Estuary (A) and M'diq Bay (B)

Fig. 6. Temporal variation of cell densities of the different phytoplankton groups in Oued Laou Estuary (A) and M'diq Bay (B)

Fig. 7. Temporal succession of the phytoplankton abundance, species richness and species diversity values (H') in Oued Laou Estuary and M'diq Bay.

Fig. 8. Annual succession of the dominant Taxa at Oued Laou Estuary from April 2008 to June 2009. The values correspond to relative densities. Astg. : *Asterionella glacialis*, Chae : *Chaetoceros* spp, Plag : *Plagioselmis* sp., Eugl : *Euglena* sp., Gy.ca : *Gymnodinium catenatum*, Lept : *Leptocylindrus* spp, Nitz : *Nitzschia* spp, No.sc. : *Noctiluca scintillans*, Psz. : *Pseudo-nitzschia* spp, Gu.de. : *Guinardia delicatula*, Gu.st. : *Guinardia striata*, Scri : *Scrippsiella* sp, Skel : *Skeletonema* spp, Thal : *Thalassiosira* spp, Thnu : *Thalassionema nitzschioides*).

Fig. 9. Annual succession of the dominant Taxa at M'diq bay from April 2008 to June 2009. The values correspond to relative densities. Astg : *Asterionella glacialis*, Chae : *Chaetoceros* spp, Plag : *Plagioselmis* sp., Gyr : *Gyrodinium* spp., Lept : *Leptocylindrus* spp, Nitz : *Nitzschia* spp, Prtt :

Prorocentrum triestinum, Psz : *Pseudo-nitzschia* spp, Scri : *Scrippsiella* sp, Skel : *Skeletonema* spp, Thal : *Thalassiosira* spp, Th.nu. : *Thalassionema nitzschioides*).

Fig. 10. Temporal variation in cells L⁻¹ of (A) *Dinophysis*. spp., (B) *Prorocentrum lima* and (C) *Ostreopsis* spp. at Oued Laou Estuary and M'diq Bay.

Fig. 11. Temporal variation of (A) *Pseudo-nitzschia* spp. (cells L⁻¹) and domoic acid (µgDA g⁻¹ shellfish flesh) in the cockle and sweet clam at Oued Laou Estuary (B) and M'diq Bay (C).

Fig. 12. Temporal variations of (A) *Alexandrium* spp., (B) *Gymnodinium catenatum*, paralytic shellfish poisoning amount (µg STXeq Kg⁻¹ shellfish flesh) in the cockle and sweet clam at Oued Laou Estuary (C) and M'diq Bay (D).

Fig. 13. Factorial Discriminant Analysis (FDA) for Oued Laou Estuary (A) and M'diq Bay (C). Astg. : *Asterionella glacialis*, Chae : *Chaetoceros* spp, Plag : *Plagioselmis* sp., Eugl : *Euglena* sp., Gy.ca : *Gymnodinium catenatum*, Lept : *Leptocylindrus* spp, Nitz : *Nitzschia* spp, No.sc : *Noctiluca scintillans*, Psz : *Pseudo-nitzschia* spp, Gu.de : *Guinardia delicatula*, Gu.st : *Guinardia striata*, Scri : *Scrippsiella* sp, Skel : *Skeletonema* spp, Thal : *Thalassiosira* spp, Thnu : *Thalassionema nitzschioides*, Din : *Dinophysis* spp, Het : *Heterocapsa* sp., Cer : *Ceratium* spp., Pleu: *Pleurosigma* spp). T°: Temperature, NO₃: nitrate, NO₂ : nitrite, NH₄⁺: ammonium and PO₄: phosphorus.

FDA sample ordination plot considering the seasons; Oued Laou Estuary (B) and M'diq Bay (D).

Table 1. Mean values of physico-chemical environmental parameters at M'diq Bay (Station 2), Mean and Standard Error (SE); n = number of samples; Min = minimum value; Max = maximum value. NH₄: ammonium, NO₂: nitrite, NO₃: nitrate, PO₄: phosphorus and SiO₄: silicate.

Table 2. Spearman correlations linking phytoplankton groups and environmental factors in Mdiq Bay (S2) and Oued Laou Estuary(S1). * Raphidophyceae and Chlorophyceae species were not observed in Mdiq bay.

Table 3. Values of Spearman rank correlation coefficient between dominant species and environmental parameters in Oued Laou Estuary (*: significant). Astg. : *Asterionella glacialis*, Chae : *Chaetoceros* spp, Plag : *Plagioselmis* sp., Eugl : *Euglena* sp., Gy.ca : *Gymnodinium catenatum*, Lept : *Leptocylindrus* spp, Nitz : *Nitzschia* spp, No.sc : *Noctiluca scintillans*, Psz : *Pseudo-nitzschia* spp, Gu.de : *Guinardia delicatula*, Gu.st : *Guinardia striata*, Scri : *Scrippsiella* sp, Skel : *Skeletonema* spp, Thal : *Thalassiosira* spp, Th.nu : *Thalassionema nitzschioides*, Din : *Dinophysis* spp, Het : *Heterocapsa* spp, Cer : *Ceratium*

spp, Pleu: *Pleurosigma* spp). Si: silicate, PO₄: phosphorus, NH₄: ammonium, NO₂: nitrite and NO₃: nitrate.

Table 4. Values of Spearman rank correlation coefficient between dominant species and environmental parameters in M'diq bay (* significant). Astg : *Asterionella glacialis*, Chae : *Chaetoceros* spp, Plag : *Plagioselmis* sp., Gyr : *Gyrodinium* spp., Lept : *Leptocylindrus* spp, Nitz : *Nitzschia* spp, Prrt : *Prorocentrum triestinum*, Psz : *Pseudo-nitzschia* spp, Scri : *Scrippsiella* sp, Skel : *Skeletonema* spp, Thal : *Thalassiosira* spp, Th.nu : *Thalassionema nitzschioides*, Alex: *Alexandrium* spp, Gy. ca: *Gymnodinium catenatum*, Din: *Dinophysis* spp, Euc: *Eucampia zodiacus*, Gu.st : *Guinardia striata*). Si: silicate, PO₄: phosphorus, NH₄: ammonium, NO₂: nitrite and NO₃: nitrate.

References

- Abadie, E., Rhodes, L., Chomérat, N., Chiantella, C., Crottier, A., Laabir, M. 2018. Occurrence of the neurotoxic dinoflagellate *Vulcanodinium rugosum* in Mediterranean: What are the driving environmental factors of its dynamic ?.Harmful Algae. 75, 75-86.
- Abdel-Halim, A.M., Khairy, H.M., 2007. Potential impact of some abiotic parameters on a phytoplankton community in a confined bay of the Eastern Mediterranean Sea: Eastern Harbour of Alexandria, Egypt. Med. Mar. Sc. 8, 49-64.
- Abdenadher, M., Hamza, A., Fekih, W., Hannachi, I., Zouari Bellaaj, A., Bradai, M.N., Aleya, L., 2012. Factors determining the dynamics of toxic blooms of *Alexandrium minutum* during a 10-year study along the shallow south western Mediterranean coasts. Estuar. Coast. Shelf Sci. 106, 102–111.

- 585 Aissaoui, A., Dhib, A., Reguera, B., Ben Hassine, O.K., Turki, S., Aleya, L., 2014. First evidence
586 of cell deformation occurrence during a *Dinophysis* bloom along the shores of the Gulf of
587 Tunis (SW Mediterranean Sea). *Harmful Algae*. 39, 191-201.
- 588
- 589 Aminot, A., Chaussepied, M., 1983. Manuel des analyses chimiques en milieu marin. Brest,
590 CNEXO, 395p.
- 591 AOAC, 1990. Paralytic Shellfish Poisoning, Biological method, final action. In AOAC (Eds.)
592 official methods of analysis. 15th ed. Method n° 959.08 JAOAC, Arlington, VA.881-882.
- 593 Armi, Z, Trabelsi, E., Turki., S., Béjaoui, B., Maïz Naceur, B., 2010. Seasonal phytoplankton
594 responses to environmental factors in a shallow Mediterranean lagoon. *J. Mar. Sc. Tech.* 15,
595 417-426.
- 596 Band-Schmidt, C.J., Bustillos-Guzman, J.J., Hernandez-Sandoval, F.E., Núñez-Vazquez, E.J.,
597 Lopez-Cortes D.J., 2014. Effect of temperature on growth and paralytic toxin profiles in
598 isolates of *Gymnodinium catenatum* (Dinophyceae) from the Pacific coast of Mexico.
599 *Toxicon*. 90, 199-212.
- 600 Band-Schmidt, C.J., Bustillos-Guzman, J.J., Lopez-Cortes, D.J., Garate-Lizarraga, I., Núñez-
601 Vazquez, E.J., Hernandez-Sandoval, F.E., 2010. Ecological and physiological studies of
602 *Gymnodinium catenatum* in the Mexican Pacific: a review. *Mar. Drugs*. 8, 1935-1961.
- 603 Bazzoni, A.M., Mudadu, A.G., Lorenzoni, G., Soro, B., Bardino, N., Arras, I., Sanna, G., Vodret,
604 B., Bazzardi, R., Marongiu, E., 2018. Detection of *Dinophysis* species and associated okadaic
605 acid in farmed shellfish: a two-year study from the western Mediterranean area. *J. Vet. Res.*
606 62, 137-144.
- 607 Ben Gharbia, H., Kéfi-Daly Yahia, O., Amzil, Z., Chomérat, N., Abadie, E., Masseret, E., Sibat, M,
608 Zmerli Triki, H., Nouri, H. and Laabir, M. 2016. Toxicity and Growth Assessments of Three
609 Thermophilic Benthic Dinoflagellates (*Ostreopsis cf. ovata*, *Prorocentrum lima* and *Coolia*
610 *monotis*) Developing in the Southern Mediterranean Basin. *Toxins*. 8, 297-335.
- 611 Ben Gharbia, H., Laabir, M., Ben Mhamed, A., Gueround, S., Daly Yahia, N., Nouri, H., M'Rabeta,
612 C., Shilig, A., Kéfi-Daly Yahia, O. 2019. Occurrence of epibenthic dinoflagellates in relation to
613 biotic substrates and to environmental factors in Southern Mediterranean (Bizerte Bay and Lagoon,

- 614 Tunisia): an emphasis on the harmful *Ostreopsis spp.*, *Prorocentrum lima* and *Coolia monotis*.
 615 Harmful Algae, in press.
- 616 Ben Rejeb Jenhani, A., Fathalli, A., Ben Naceur, H., Hayouni, D., Aouani, J., Romdhane, M.S.,
 617 2019. Screening for alien and harmful planktonic species in the Gulf of Gabes (Tunisia,
 618 Southeastern Mediterranean Sea). Reg. Stud. Mar. Sci, 27
- 619 Berland, B.R., Bonin, D.j., Maestrini, S.Y., Pointier, J.P., 1973. Etude de la fertilité des eaux
 620 marines au moyen de tests biologiques effectués avec des cultures d’algues. II Limitation
 621 nutritionnelle et viabilité de l’inoculum. Int. Revue. Gesanten. Hydrobiol. 58, 203-220.
- 622 Bourhilli, E.H., 1982. Intoxication alimentaire par les moules (51 cas), Agadir, octobre et
 623 novembre 1982. Thèse n°33. Faculté de Médecine et de Pharmacie, Rabat 84 pp.
- 624 Boutaib, R., Marhraoui, M., Oulad Abdellah, M.K., Bouchrif, B., 2011. Comparative study on
 625 faecal contamination and occurrence of *Salmonella spp.* and *Vibrio parahaemolyticus* in two
 626 species of shellfish in Morocco. Open. Environmental. Sciences. 5, 30-37.
- 627 Boyd, P.W., Strzepek, R., Fu, F., Hutchins, D.A., 2010. Environmental control of open-ocean
 628 phytoplankton groups: Now and in the future. Limnol. Ocean. 55(3), 1353–1376.
- 629 Bravo, I., Reguera, B., Martínez, A., Fraga, S., 1990. First report of *Gymnodinium catenatum*
 630 Graham on the Spanish Mediterranean coast. In: Granéli, E., Sundtröm, B., Edler, L.,
 631 Anderson, D.M. (Eds.), Toxic Marine Phytoplankton. Elsevier, New York, pp. 449–452.
- 632 Busch, J.A., Andree K.B., Diogene, J., Fernandez-Tejedor, M., Toebe, K., John, U., Krock, B.,
 633 Urban Tillmann, U., Cembella, A.D., 2016. Toxigenic algae and associated phycotoxins in
 634 two coastal embayments in the Ebro Delta (NW Mediterranean). Harmful Algae. 55, 191–
 635 201.
- 636 Bužančić, M., Gladan, Z.N., Marasović, I., Kušpilić, G., Grbec, B., 2016. Eutrophication influence
 637 on phytoplankton community composition in three bays on the eastern Adriatic coast.
 638 Oceanologia. 58, 302-316.
- 639 Carstensen, J., Klais, R., Cloern, J.E., 2015. Phytoplankton blooms in estuarine and coastal waters:
 640 Seasonal patterns and key species. Estuar Coast Shelf Sci. 162, 98–109.

- 641 Cerino, F., Orsini, L., Sarno, D., Carmela, D.A., Tartaglione L., Zingone, A., 2005. The
642 alternation of different morphotypes in the seasonal cycle of the toxic diatom *Pseudo-*
643 *nitzschia galaxiae*. Harmful Algae. 4, 33–48.
- 644 Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. Mar.
645 Ecol. Progr. Ser. 210, 223 –253.
- 646 Dammak-Zouari, H., Hamza, A., Bouain, A., 2009. Gymnodiniales in the Gulf of Gabes (Tunisia).
647 Cah. Biol. Mar. 50, 153-170.
- 648 Daoudi, M., Serve, L., Rharbi, N., El Madani, F., Vouv  , F., 2012. Phytoplankton distribution in
649 the Nador lagoon (Morocco) and possible risks for harmful algal blooms. Trans. Waters. Bull.
650 6(1), 4-19.
- 651 Delgado, M., 1990. Phytoplankton distribution along the Spanish coast of the Albor  n Sea. Sci.
652 Mar. 54, 169–178.
- 653 Dhib, A., Frossard, V., Turki, S., Aleya, L. 2013. Dynamics of harmful dinoflagellates driven by
654 temperature and salinity in a northeastern Mediterranean lagoon. Environ. Monit. Assess. 185,
655 3369–3382.
- 656 Dhib, A., Fertouna-Bellakhal, M., Turki, S., Aleya, L., 2015. Harmful planktonic and epiphytic
657 microalgae in a Mediterranean Lagoon: the contribution of the macrophyte *Ruppia cirrhosa* to
658 microalgae dissemination. Harmful Algae 45, 1–13.
- 659 Draredja, MA ., Frihi, H. Boualleg, C., Gofart, A., Abadie, E., Laabir, M. 2019. Seasonal variations of
660 phytoplankton community in relation to environmental factors in a protected meso-oligotrophic
661 southern Mediterranean marine ecosystem (Mellah lagoon, Algeria) with an emphasis of HAB
662 species. Environ. Monit. Assess. 191(10).
- 663 EL Madani, F., Chia  r, A., Chafi, A., 2011. Phytoplankton composition and abundance
664 assessment in the Nador lagoon (Mediterranean coast of Morocco). Acta. Botanica. Croatia.
665 70(2), 269-288.
- 666 Fertouna-Bellakhal, M., Dhib, A., Fathalli, A., Bellakhal, M., Chome, N., Masseret, E., Laabir, M.,
667 Turki, S., Aleya, L., 2015. *Alexandrium pacificum* Litaker sp. nov (Group IV): Resting cyst

- 668 distribution and toxin profile of vegetative cells in Bizerte Lagoon (Tunisia, Southern Mediterranean
669 Sea). *Harmful Algae*. 48, 69-82.
- 670 Fraga, S., 1996. Wintering of *Gymnodinium catenatum* Graham in Iberian waters. In. Yasumoto,
671 T., Oshima, Y., Fukuyo, Y. (Eds.), *Harmful and Toxic Algal Blooms*. IOC of UNESCO,
672 Paris, pp. 211-214.
- 673 Frehi, H., Coute, A., Mascarell, G., Perrette-Gallet, C., Ayada, M., Kara, MH., 2007. Harmful and
674 red-tide dinoflagellates in the Annaba bay (Algeria). *Comptes Rend Biol.* 330, 615-628.
- 675 Gailhard, I., 2003. Analyse de la variabilité spatio-temporelle des populations microalgales
676 côtières observées par le « Réseau de surveillance du phytoplancton et des phycotoxines »
677 (REPHY). Thèse de Doctorat d'université. Université d'Aix-Marseille II, Aix-Marseille,
678 France. 187p.
- 679 García-Altare, M., Casanova, A., Fernández-Tejedor, M., Diogène, J., De la Iglesia, P., 2016.
680 Bloom of *Dinophysis* spp. dominated by *D. sacculus* and its related diarrhetic shellfish
681 poisoning (DSP) outbreak in Alfacs Bay (Catalonia, NW Mediterranean Sea): Identification
682 of DSP toxins in phytoplankton, shellfish and passive samplers. *Reg. Stud. Mar. Sc.* 6, 19–28.
- 683 Gobler, C.J., Doherty, O.M., Owen, M., Hattenrath-Lehmann, T.K., Theresa, K., Griffith, A.W.,
684 Kang, Y., Litaker, R.W., 2017. Ocean warming since 1982 has expanded the niche of toxic
685 algal blooms in the North Atlantic and North Pacific oceans. *Proc. National. Acad. Sci. USA*.
686 14(19), 4975-4980.
- 687 Gomez, F., 2003. The toxic dinoflagellate *Gymnodinium catenatum*: an invader in the
688 Mediterranean sea. *Acta. Bot. Croat.* 62, 65–72.
- 689 Grzebyk, D., Audic, S., Lasserre, B., Abadie, E., de Vargas, C., Bec, B., 2017. Insights into the
690 harmful algal flora in northwestern Mediterranean coastal lagoons revealed by
691 pyrosequencing metabarcodes of the 28S rRNA gene. *Harmful Algae*. 68, 1–16.
- 692 Hallegraeff, G.M., 1993. A review of harmful algal blooms and their apparent global increase.
693 *Phycologia* 32, 79–99.

- 694 Hallegraeff, G.M., 2010. Ocean climate change, phytoplankton community responses, and
695 harmful algal blooms: a formidable predictive challenge. *J. Phycol.* 46, 220–235.
- 696 Illoul, H., Maso, M., Fortuno, J.M., Cros, L., Morales-Blake, A., Sériidji, R., 2008. Potentially
697 harmful microalgae in coastal waters of the Algiers area (Southern Mediterranean Sea).
698 *Cryptogamie Algol.* 29, 261-278.
- 699 Jouenne, F., Lefebvre, S., Véron, B., Lagadeuc, Y., 2007. Phytoplankton community structure and
700 primary production in small intertidal estuarine-bay ecosystem (eastern English Channel,
701 France). *Mar. Biol.* 151, 805–825.
- 702 Kim, G.U., Seo, K.H., Chen, D.L., 2019. Climate change over the Mediterranean and current
703 destruction of marine ecosystem. *Scientific Reports*. Vol 9. DOI: 10.1038/s41598-019-55303.
- 704 Laabir, M., Jauzein, C., Genovesi, B., Masseret, E., Grzebyk, D., Cecchi, P., Vaquer, A., Perrin, Y.,
705 Collos, Y., 2011. Influence of temperature, salinity and irradiance on the growth and cell yield of the
706 harmful red tide dinoflagellate *Alexandrium catenella* colonising Mediterranean waters. *J. Plankton*
707 *Res.* 33, 1550-1563.
- 708 Laanaia, N., Vaquer, A., Fiandrino, A., Genovesi, B., Pastoureaud, A., Cecchi, P., Collos, Y. ; 2013.
709 Wind and temperature controls on *Alexandrium* blooms (2000-2007) in Thau lagoon (Western
710 Mediterranean). *Harmful Algae.* 28, 31-36.
- 711 Lopes, C.B., Lillebo, A.I., Dias, J.M., Pereira, E., Vale, C., Duarte, A.C., 2007. Nutrient dynamics
712 and seasonal succession of phytoplankton assemblages in a Southern European Estuary: Ria
713 de Aveiro, Portugal. *Estuar. Coast. Shelf. Sci.* 71, 480–490.
- 714 Macias, D., García-Gorríz, E., Stips, A., 2018. Major fertilization sources and mechanisms for
715 Mediterranean Sea coastal ecosystems. *Limnol. Oceanogr.* 63, 897–914.
- 716 Macias, D.M., Garcia-Gorriz, E., Stips, A., 2015. Productivity changes in the Mediterranean Sea
717 for the twenty-first century in response to changes in the regional atmospheric forcing. *Front.*
718 *Mar. Sci.* 2, 79.
- 719 Margalef, R., Estrada, M., Blasco, D., 1979. Functional morphology of organisms involved in Red
720 Tides, as adapted to decaying turbulence. In Taylor DL, Seliger H.H., (eds) *Toxic Dinoflagel-*

late Blooms, Proc. 2nd Int. Conf., Key Biscayne, Florida, USA, Oct. 31-Nov. 15, 1978: 89-94.

Melliti Ben Garali, S., Sahraoui, I., De la Iglesia, P., Chalghaf, M., Diogène, J., Jamel Ksouri, J., Sakka Hlaili, A., 2019. Factors driving the seasonal dynamics of *Pseudo-nitzschia* species and domoic acid at mussel farming in the SW Mediterranean Sea. Chemistry Ecology. DOI: 10.1080/02757540.2019.1676417

Moita, M.T., Vilarinha, M.G., Palma, A.S., 1998. On the variability of *G. catenatum* Graham blooms in Portuguese waters. In : Reguera, B., Blanco, J., Fernandez, M.L., Wyatt, T. (Eds.), Harmful Algae : Xunta de Galicia and IOC of UNESCO, Spain, pp. 118-121.

Quilliam, M.A., 1995. Chemical methods for domoic acid, the amnesic shellfish poisoning (ASP) toxin. In: G.M. Hallegraeff, D.M. Anderson & A.D. Cembella (Eds.), Manual on Harmful Marine Microalgae, Monographs on Oceanographic Methodology, Vol. 11, Chapter 9. Intergovernmental Oceanographic Commission (UNESCO), Paris, 247-266 (2003)

Quiroga, I., 2006. *Pseudo-nitzschia* blooms in the bay of Banyuls-sur-mer, northwestern Mediterranean Sea. Diatom Res. 21, 91–104.

Redfield, A.C., Ketchum, B.H., Richards, F.A., 1963. The influence of organisms on the composition of seawater. In M. N. Hill (Eds.). The sea, Vol. 2. Interscience. p. 26– 77

Richard, M., 1987. Atlas de phytoplancton marin. Vol. 2, Diatomophycées. CNRS, Paris


Rijal Leblad, B., Lundholm, N., Goux, D., Veron, B., Sagou, R., Taleb, H., Nhala, H., Er-Raioui, H., 2013. *Pseudo-nitzschia* Peragallo (Bacillariophyceae) diversity and domoic acid accumulation in tuberculate cockles and sweet clams in M'diq Bay, Morocco. Acta. Botanica. Croatia. 72 (1), 35–47.


Rijal leblad, B., 2012. Variabilité de la contamination des bivalves (*Callista chione* & *Acanthocardia tuberculata*) par les phycotoxines paralysantes (PSP) et amnésiantes (ASP) dans la frange littorale méditerranéenne marocaine: cas de la baie de M'Diq et de l'estuaire de Oued Laou. MSc Thesis N° 57. Université Abdelmalek Assaadi, Morocco. 176p.


- Rijal Leblad, B., Nhhala1, H., Daoudi, M., Marhraoui, M., Ouelad Abdellah, N.K., Veron, B., Er-
 Raioui, H., 2017. Contamination and depuration of Paralytic Shellfish Poisoning by
Acanthocardia tuberculata cockles and *Callista chione* clams in Moroccan waters. J. Mater.
 Environ. Sci. 8(S), 4634-4641.
- Ruggiero, M.V., Sarno, D., Barra, L., Kooistra, W.H.C.F., Montresor, M., Zingone, A., 2015.
 Diversity and temporal pattern of *Pseudo-nitzschia* species (Bacillariophyceae) through the
 molecular lens. Harmful Algae. 42, 15–24.
- Sagou, R., Amanhir, R., Taleb, H., Vale, P., Blaghen, M., Loutfi, M. 2005. Comparative study on
 differential accumulation of PSP toxins between cockle (*Acanthocardia tuberculatum*) and
 sweet clam (*Callista chione*). Toxicon. 46, 612-618.
- Sahraoui, I., Sakka Hlaili, A., Hadj Mabrouk, H., 2009. Blooms of the diatom genus *Pseudo-*
nitzschia H. Peragallo in Bizerte Lagoon (Tunisia. SW Mediterranean). Diatom Res. 24,175–
 190.
- Salhi, N., Zmerli Triki, H., Molinero, J.C., Laabir, M., Sehli, E., Bellaaj-Zouari, A., Daly Yahia,
 N., Kefi Daly Yahia, O., 2018. Seasonal variability of picophytoplankton under contrasting
 environments in northern Tunisian coasts, southwestern Mediterranean Sea Mar. Pol. Bul.
 129, 866–874.
- Shen, A., Ma, Z., Jiang, K., Li, D., 2016. Effects of temperature on growth, photophysiology,
 Rubisco gene expression in *Prorocentrum donghaiense* and *Karenia mikimotoi*. Ocean Sci. J.
 51 (4), 581–589.
- Silva, T., Caeiro, M.F., Reis Costa, P., Amorim, A., 2015. *Gymnodinium catenatum* Graham
 isolated from the Portuguese coast: Toxin content and genetic characterization. Harmful
 Algae. 48, 94-104.
- Smayda, T.J., 1984. Variations and long-term changes in Narragansett Bay, a phytoplankton-
 based coastal marine ecosystem: relevance to field monitoring for pollution assessment. In


- 772 Concept in marine pollution, pp. 663–679. Ed. By H. H. White. Maryland Seagrass College,
773 University of Maryland
- 774 Tahri, L.J., 1998. *Gymnodinium catenatum* Graham blooms on Moroccan waters. In Reguera, B.,
775 Blanco, J., Fernandez, M.L., Wyatt, T. (Eds.), Harmful Algae. Xunta de Galicia and IOC of
776 UNESCO, Spain, pp. 66-67.
- 777 Taleb, H., Vale, P., Blaghen, M., 2003. Spatial and temporal evolution of PSP toxins along the
778 Atlantic shore of Morocco. *Toxicon*. 41, 199-205.
- 779 Taleb, H., Vale, P., Jaime, E., Blaghen, M., 2001. Study of paralytic shellfish poisoning toxin
780 profile in shellfish from the Mediterranean shore of Morocco. *Toxicon*, 39, 1855-1861.
- 781 Tber, A., 1983. Personal communication. Laboratoire d'analyses des services vétérinaires,
782 Casablanca, Maroc 5p.
- 783 Thorel, M., Claquin P., Schapira, M., Le Gendre, R., Riou, P., Goux, D., LeRoy,
784 B., Raimbault, V., Deton-Cabanillas, A.F., Bazin, P., Kientz-Bouchart, V., Fauchot, J.,
785 2017. Nutrient ratios influence variability in *Pseudo-nitzschia* species diversity and particulate
786 domoic acid production in the Bay of Seine (France). *Harmful Algae*. 68, 192-205.
- 787 Trainer, V.L., Hickey, B.M., Lessard, E.J., Cochlan, W.P., Trick, C.G., Wells, M.L., MacFadyen,
788 A., Moore, S.K., 2009. Variability of *Pseudo-nitzschia* and domoic acid in the Juan de Furca
789 eddy region and its adjacent shelves. *Limnol. Oceanog.* 51, 289-308.
- 790 Trainer, V.L., Moore, L., Bill, B.D., Adams, N.G., Harrington, N., Borchert, J., da Silva, D.A.M.,
791 and Bich-Thuy, L., Eberhart, B.T., 2013. Diarrhetic Shellfish Toxins and Other Lipophilic
792 Toxins of Human Health Concern in Washington State. *Mar. Drugs*. 11, 1815-1835.
- 793 Trombetta, T., Vidussi, F., Mas, S., Parin, D., Simier, M., Mostajir, B., 2019. Water temperature
794 drives phytoplankton blooms in coastal waters. *Plos One*. DOI: 10.1080/02757540.2019.1676417
- 795 Uthermol, H., 1958. Zur vervollkomnung der quantitaven phyto-plankton-methodik. *Mitt. Int.*
796 *Ver. Ther. Angew. Limnol.*, 9

- Van Dolah, F.M., 2000. Marine algal toxins: origins, health effects, and their increased occurrence. *Environ. Health Perspect.* 108, 133–141.
- Vila, M.; Garcés, E.; Masó, M.; Camp, J. 2001a. Is the distribution of the toxic dinoflagellate *Alexandrium catenella* expanding along the NW Mediterranean coast? *Mar. Ecol. Prog. Ser.* 222, 73–83.
- Vila, M., Garcés, E., Masó, M., 2001b. Potentially toxic epiphytic dinoflagellate assemblages on macroalgae in the NW Mediterranean. *Aquat. Microb. Ecol.* 26, 51–60.
- Vila, M., Giacobbe, M.G., Masó, M., Gangemib, E., Sampedro, N., Pennam, A., Azzarob, F., Campa, J., Galluzzic, L., 2005. A comparative study on recurrent blooms of *Alexandrium minutum* in two Mediterranean coastal areas. *Harmful Algae.* 4, 673–695.
- Vilicic, D., Marasovi, I., Miokovi, D., 2002. Checklist of phytoplankton in the eastern Adriatic Sea. *Acta. Botanica. Croatica.* 61, 57–91.
- Wells, M.L., Karlson, B., Wulff, A., Kudela, R., Trick, C., Asnaghi, V., Berdalet, E., Cochlan, W., Davidson, K., De Rijcke, M., Dutkiewicz, S., Hallegraeff, G., Flynn, K., Legrand, C., Paerl, H., Silke, J., Suikkanen, S., Thompson, P., Trainer, V., 2019. Future HAB Science: Directions and Challenges in a Changing Climate. *Harmful Algae.* in press. .
- Wells, M.L., Trainer, V.L., Smayda, T.J., Karlson, B.S., Trick, C.G., Kudela, R.M., Ishikawa, A., Bernard, S., Wulff, A., Anderson, D.M., 2015. Harmful algal blooms and climate change: Learning from the past and present to forecast the future. *Harmful Algae.* 49, 68–93.
- Zmerli-Triki, H., Laabir, M., Moeller, P., Chomérat, N., Kéfi Daly-Yahia, O., 2016. Production of goniodomin A by the dinoflagellate *Alexandrium pseudogonyaulax* developing in southern Mediterranean (Bizerte Lagoon, Tunisia). *Toxicon.* 111, 91–99.












A




B





A




B























1 **Table 1.**

2

Station		T° (°C)	Salinity	pH	[NH ₄] (μM)	[NO ₂] (μM)	[NO ₃] (μM)	[PO ₄] (μM)	[SiO ₄] (μM)	Rainfall (mm/month)
Spring										
S1 (n=19)	Maen (SE)	16.85 (± 1.17)	35.18 (± 1.64)	8.06 (± 0.15)	2.77 (± 2.72)	0.03 (± 0.02)	1.89 (± 1.64)	0.16 (± 0.10)	2.44 (± 1.93)	5.33
	Min – Max	15.5 - 19.7	31 – 36.7	7.78 – 8.24	0.09 – 9.20	0 – 0.06	0.23 – 5.39	0.04 – 0.39	0.28 – 7.46	0 - 16
S2 (n=22)	Maen (SE)	16.46 (± 0.78)	35.73 (± 0.78)	8.12 (± 0.18)	3.75 (± 3.76)	0.03 (± 0.02)	0.72 (± 0.34)	0.30 (± 0.48)	4.1 (± 3.90)	47.58
	Min – Max	15.5 - 18.5	34.1 – 36.7	7.87 – 8.44	0.12 - 17.03	0.01 - 0.12	0.22 - 1.49	0 - 1.77	0.62 - 17.46	1 – 114.4
Summer										
S1 (n=13)	Maen (SE)	20.12 (± 1.68)	36.54 (± 0.09)	8.11 (± 0.13)	9.07 (± 9.85)	0.03 (± 0.03)	0.92 (± 0.82)	0.19 (± 0.06)	2.27 (± 1.39)	0
	Min -Max	17.5 – 22	36.5 – 36.8	7.81 – 8.21	0.86 – 29.97	0.01 – 0.12	0.23 – 3.27	0.04 - 0.27	0.56 – 5.08	
S2 (n=14)	Maen (SE)	19.50 (± 1.29)	36.51 (± 0.05)	8.14 (± 0.17)	6.01 (± 7.80)	0.04 (± 0.02)	0.32 (± 0.21)	0.32 (± 0.56)	2.39 (± 1.69)	3.16
	Min – Max	19.1 – 21.4	36.5 - 36.7	7.81 – 8.40	0.49 – 29.51	0.01 – 0.09	0.17 – 0.45	0.02 -2.06	0.23 – 4.75	0 – 7.5
Autumn										
S1 (n=9)	Maen (SE)	18.39 (± 1.51)	33.83 (± 2.05)	7.96 (± 0.08)	24.53 (± 10.19)	0.07 (±0.05)	1.60 (± 1.02)	0.94 (± 0.83)	1.24 (± 0.78)	148.5
	Min - Max	16 – 20.5	31 - 36.5	7.82 – 8.11	11.56 – 37.91	0.01 – 0.17	0.25 – 2.99	0.10 – 2.10	0.22 – 2.26	38 – 236.5
S2 (n=9)	Maen (SE)	19.43 (± 1.09)	36.06 (± 0.36)	7.93 (± 0.11)	17.74 (± 14.63)	0.05 (± 0.02)	1.38 (± 0.87)	0.64 (± 0.65)	4.12 (± 2.05)	258.83
	Min – Max	17.8 -21.4	35.6 - 36.5	7.78 – 8.04	3.74 - 43.13	0.03 – 0.09	0.37 – 2.59	0.04 – 1.54	1.41 – 7.32	160.5 – 277.5
Winter										
S1 (n=6)	Maen (SE)	14.32 (± 0.67)	35.38 (± 0.83)	7.89 (± 0.08)	3.32 (± 4.27)	0.07 (± 0.05)	3.72 (± 2.37)	0.19 (± 0.05)	2.44 (± 1.29)	83.8
	Min - Max	13.5 – 15.2	33.90 – 36.30	7.80 – 8.01	0.22 – 11.68	0.02 – 0.13	1.02 – 7.52	0.12 – 0.23	1.13 – 4.75	24 – 84.9
S2 (n=9)	Maen (SE)	16.07 (± 0.52)	35.06 (± 0.35)	7.97 (± 0.14)	3.19 (± 4.02)	0.09 (± 0.05)	1.65 (± 0.43)	0.25 (± 0.22)	2.49 (± 1.61)	127.76
	Min – Max	15.5 – 17.2	34.6 – 35.7	7.78 – 8.17	0.22 – 13.26	0.05 – 0.19	1.05 – 2.38	0.10 – 0.82	1.30 – 6.50	48 – 135.8

Table 2. :

Variables	Site Number	Rainfall	T (°C)	Salinity	pH	[PO ₄]	[NO ₂]	[NH ₄]	[NO ₃]	DIN	[SiO ₄]
Species Richness	S1	-0.231	0.296	0.388	0.420	-0.042	-0.217	-0.095	-0.333	-0.094	-0.056
	S2	-0.713	0.339	0.497	0.353	0.049	-0.392	0.125	-0.435	-0.040	0.036
Species Diversity (H')	S1	0.143	-0.014	-0.304	-0.242	0.189	-0.056	0.098	0.184	0.173	0.040
	S2	-0.154	0.325	0.298	-0.193	-0.078	-0.092	0.142	-0.145	0.001	0.055
Prymnosiohyceae	S1	-0.031	0.080	0.170	0.327	0.001	-0.176	-0.126	-0.292	-0.138	0.005
	S2	-0.246	0.005	0.324	0.448	0.088	-0.074	-0.023	-0.274	-0.039	-0.142
Cryptophyceae	S1	-0.277	-0.032	0.186	0.420	0.074	0.042	-0.035	0.018	0.000	0.276
	S2	-0.218	-0.107	0.378	0.371	-0.067	-0.228	-0.115	-0.018	-0.115	-0.029
Euglenophyceae	S1	-0.011	0.059	0.095	0.149	-0.079	-0.176	0.116	-0.313	0.101	-0.173
	S2	-0.084	0.079	0.138	-0.269	-0.138	0.002	0.184	-0.122	0.237	0.205
Dictyochophyceae	S1	0.137	-0.147	-0.107	0.034	0.101	0.229	-0.027	0.035	0.099	-0.130
	S2	0.032	-0.224	-0.291	-0.203	0.177	0.334	0.034	0.298	0.095	0.154
Dinophyceae	S1	-0.378	0.181	0.170	0.386	-0.092	-0.047	-0.025	-0.196	-0.177	0.050
	S2	-0.573	-0.109	0.127	0.433	0.067	-0.388	-0.114	-0.237	-0.228	-0.068
Diatomophyceae	S1	-0.310	0.254	0.544	0.349	-0.263	-0.244	-0.194	-0.261	-0.130	-0.070
	S2	-0.713	0.101	0.362	0.400	-0.062	-0.377	-0.102	-0.445	-0.162	-0.164
Raphidophyceae	S1	-0.134	0.120	0.124	0.005	-0.224	-0.181	-0.250	-0.141	-0.196	-0.217
	S2	*	*	*	*	*	*	*	*	*	*
Chlorophyceae	S1	-0.065	0.174	0.177	0.079	0.273	-0.156	-0.147	-0.068	-0.155	0.152
	S2	*	*	*	*	*	*	*	*	*	*
Total phytoplankton	S1	-0.402	0.196	0.448	0.432	-0.237	-0.068	-0.120	-0.243	-0.244	-0.040
	S2	-0.692	0.064	0.397	0.469	-0.080	-0.455	-0.126	-0.409	-0.181	-0.179

8

9 **Table 3.**

Variable	Gy.Ca	Din.	Het.	Cer.	No.sc.	Scri.	Chae.	Psz.	Lept.	Nitz.	Gu.de	Gu.st.	Skel.	Th.nu	Astg.	Thal.	Plag.	Eugl.	Pleu.
[SiO ₄]	-0.21	0.06	0.26	-0.09	0.12	0.06	-0.05	-0.21	0.00	0.01	-0.00	-0.13	-0.18	-0.12	0.23	0.00	0.32*	-0.13	-0.12
[PO ₄]	0.20	-0.02	0.03	0.18	-0.08	-0.23	-0.20	-0.27	-0.19	-0.31*	0.10	0.06	0.12	-0.26	-0.08	-0.09	0.01	-0.11	0.03
[NH ₄]	0.32*	0.00	-0.18	0.29*	-0.26	-0.09	-0.14	-0.05	-0.18	-0.17	0.20	0.06	0.36*	-0.37*	-0.19	-0.21	-0.09	0.08	-0.13
[NO ₂]	0.26	-0.15	-0.34*	0.01	-0.16	-0.32*	-0.25	-0.19	-0.21	-0.42*	-0.09	0.02	0.10	-0.19	0.05	0.03	0.02	-0.21	0.18
[NO ₃]	0.00	-0.24	-0.17	-0.15	-0.02	-0.26	-0.38*	-0.43*	-0.32*	-0.31*	-0.32*	-0.13	-0.15	-0.16	-0.06	-0.05	0.00	-0.27	0.06
T°	-0.03	0.25	0.29*	0.13	-0.06	0.19	0.36*	0.41*	0.32*	0.23	0.55*	0.40*	0.42*	-0.37*	0.16	-0.21	-0.03	0.05	-0.28
Salinity	-0.28	0.32*	0.27	0.01	-0.00	0.31*	0.56*	0.52*	0.52*	0.43*	0.46*	0.31*	0.25	-0.06	0.33*	0.08	0.18	0.09	-0.26
pH	-0.08	0.37*	0.30*	-0.03	0.04	0.30*	0.36*	0.15	0.27	0.13	0.14	-0.05	-0.05	0.12	0.36*	0.09	0.42*	0.14	-0.06
Rainfull	0.46*	-0.32*	-0.47*	0.15	-0.20	-0.32*	-0.37*	-0.19	-0.33*	-0.43*	-0.36*	-0.10	-0.12	0.17	-0.18	0.16	-0.27	-0.01	0.30*

10

11 **Table 4**

Variable	Alex	Gy. Ca	Din.	Gyr.	Prr.t.	Scri.	Chaet.	Pzs.	Lept.	Euc.	Nitz.	Gu.st.	Skel.	Th.nu.	Astg.	Plag.	Thal.
T°	0.18	-0.03	-0.00	-0.26	-0.14	-0.13	0.34*	0.16	0.11	-0.17	-0.05	0.43*	0.42*	-0.14	-0.18	-0.10	-0.22
Salinity	0.27*	-0.14	0.20	-0.07	0.04	0.07	0.53*	0.26	0.25	-0.09	0.18	0.03	0.23	0.11	-0.05	0.37*	-0.31*
pH	0.28*	-0.13	0.29*	0.50*	0.35*	0.30*	0.39*	0.34*	0.27*	0.27*	0.14	-0.39*	-0.16	0.19	0.03	0.37*	0.01
[NO ₂]	-0.46*	0.34*	-0.22	-0.03	-0.44*	-0.35*	-0.39*	-0.38*	-0.29*	-0.10	-0.28*	0.00	-0.09	0.04	-0.10	-0.23	0.04
[PO ₄]	-0.07	0.21	0.33*	0.27*	-0.03	-0.04	-0.05	-0.04	0.02	0.37*	-0.09	-0.26	-0.10	0.30*	-0.23	-0.07	0.10
[NH ₄]	0.17	0.23	0.01	-0.16	-0.15	-0.13	0.00	-0.01	-0.06	-0.06	-0.1	-0.00	0.31*	0.13	-0.34*	-0.13	-0.11
[NO ₃]	-0.42*	0.39*	-0.16	0.08	-0.25	-0.27*	-0.56*	-0.44*	-0.44*	-0.01	-0.18	-0.22	-0.27*	0.03	-0.06	-0.03	0.13
[SiO ₄]	0.04	-0.01	-0.09	-0.24	0.07	-0.07	-0.17	-0.25	-0.21	-0.24	-0.08	-0.14	0.21	-0.05	-0.01	-0.03	-0.13
Rainfull	-0.27*	0.28*	-0.15	0.14	-0.12	-0.20	-0.50*	-0.31*	-0.40*	0.11	-0.16	-0.35*	-0.14	0.22	-0.06	-0.10	0.098

12