



HAL
open science

Metabolic alterations in Hurley stage 1 Hidradenitis Suppurativa

Maïa Delage, Laure Guenin-Macé, Jean-David Morel, Jean-Marc Doisne, Angèle Schiavo, Marie-Noëlle Ungeheuer, Thi Lam, Olivier Join-Lambert, James Santo, Aude Nassif, et al.

► **To cite this version:**

Maïa Delage, Laure Guenin-Macé, Jean-David Morel, Jean-Marc Doisne, Angèle Schiavo, et al.. Metabolic alterations in Hurley stage 1 Hidradenitis Suppurativa. 9TH CONFERENCE OF THE EUROPEAN HIDRADENITIS SUPPURATIVA FOUNDATION, Feb 2020, Athènes, Greece. hal-02911687

HAL Id: hal-02911687

<https://normandie-univ.hal.science/hal-02911687>

Submitted on 4 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Metabolic alterations in Hurley stage 1 Hidradenitis Suppurativa

Maïa Delage¹, Laure Guenin-Macé², Jean-David Morel², Jean-Marc Doïse³, Angèle Schiavo³, Marie-Noëlle Ungeheuer⁴, Thi Lam¹, Olivier Join-Lambert⁵, James Di Santo³, Aude Nassif¹, Caroline Demangel²


¹ Medical Center, Institut Pasteur, Paris, France; ² Immunobiology of Infection Unit, Institut Pasteur, INSERM U1221, Paris, France; ³ Innate Immunity Unit, Institut Pasteur, INSERM U1223, Paris, France; ⁴ ICAREB Platform, Institut Pasteur, Paris, France; ⁵ Microbiology Department, Caen Hospital and University, France

Introduction: HS pathophysiology is only partially understood. Previous metagenomic studies have identified distinctive anaerobic flora in HS skin, whose expansion correlate with lesion severity. Targeted antibiotic combinations are efficient in treating HS. Dysregulated immune responses have also been identified. In this study, we conducted the first metabolomic analysis of HS skin to gain insights into potential mechanisms linking cutaneous dysbiosis with dysregulated immune responses and HS pathophysiology.

Materials and Methods:



- 20 HS stage 1 lesion
- 20 HS stage 1 peri-lesion
- 20 controls





Skin samples were harvested from early (Stage I) lesions and adjacent healthy skin zones in 19 patients and 20 healthy subjects

500 metabolites reliably identified in all samples

Results:



A metabolomic signature associated with HS lesions:




Our analysis detected a total of 500 metabolites, which distributed differently among skin samples from HC, and from H-HS and L-HS

Principal component analysis clearly discriminated the metabolomes of L-HS and HC skin, while H-HS showed an intermediate signature

Tryptophan metabolites increased in HS lesions:



At a 5% FDR, 228 metabolites were differently modulated between H-HS and L-HS skin, among which 214 were significantly augmented in lesions. The most up-regulated metabolite in L-HS skin was quinolate (more than 10 000 times higher in L- HS lesions than in HC)



Discussion and conclusion:

It has been reported that Tryptophan metabolism can modulate immune response (Harden et al; J Allergy Clin Immunol. 2016). It seems to play a role in Inflammatory Bowel Diseases (Agus et al; Cell Host Microbe. 2018.) This study suggests Tryptophan metabolism has a key role in HS. It thus highlights a new potential pathway in HS and could help us to move forward in the comprehension of HS pathophysiology.