



# Do musicians have better mnemonic and executive performance than actors? Influence of regular musical or theatre practice in adults and in the elderly.

Mathilde Groussard, Renaud Coppalle, Thomas Hinault, Hervé Platel

## ► To cite this version:

Mathilde Groussard, Renaud Coppalle, Thomas Hinault, Hervé Platel. Do musicians have better mnemonic and executive performance than actors? Influence of regular musical or theatre practice in adults and in the elderly.. *Frontiers in Human Neuroscience*, 2020, 10.3389/fnhum.2020.557642 . hal-03113788

**HAL Id: hal-03113788**

**<https://normandie-univ.hal.science/hal-03113788>**

Submitted on 18 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Title:**

**Do musicians have better mnemonic and executive performance than actors? Influence of regular musical or theatre practice in adults and in the elderly.**

**Authors:**

**Groussard, M.<sup>1\*</sup>; Coppalle, R.<sup>1</sup>; Hinault, T.<sup>1</sup>; Platel, H.<sup>1</sup>**

**Affiliations:**

<sup>1</sup> Normandie Univ, UNICAEN, PSL Research University, EPHE, INSERM, U1077, CHU de Caen, Cyceron, Neuropsychologie et Imagerie de la Mémoire Humaine, 14000 Caen, France

**\* Correspondence to :** Mathilde Groussard, PhD, Inserm U1077, GIP Cyceron, Bd Henri Becquerel, BP 5229 14074 Caen cedex.

**E-mail** : mathilde.groussard@unicaen.fr

**Manuscript length :**

**Number of word :** 4598

**Number of tables :** 5

**Number of figures :** 2

**Disclosure statement** : All the authors declare no conflict of interest

**Manuscript contribution to the field :**

The aim of the present study was to investigate the cognitive difference between regular artistic practitioners. Several studies have demonstrated the influence of musical practice on cerebral activity and cognition, but comparing this practice to another artistic domain such as theatre had yet to be done. However, both of these artistic activities require many hours of individual or collective practice in order to reach an advanced level. Music and theatre practice require the interaction between higher-order cognitive functions and several sensory modalities (auditory, verbal, visual and motor), as well as the regular learning of new pieces. Our results are consistent with previous works suggesting that adult musicians outperformed control subjects in standardized cognitive tasks, but a lifespan approach was never adopted in previous studies. Additionally, it highlights for the first time that both musical and theatrical expertise may involve specific cognitive differences across the lifespan. Furthermore, we observed domain-specific cognitive patterns: musical practice was associated with better executive functions and working memory, and theatrical expertise with better long-term verbal memory and verbal fluency.

**Keywords:** Music; Theatre; Practice; Cognition; Aging; Lifespan

## **Abstract**

The effects of musical practice on cognition are well established yet rarely compared with other kinds of artistic training or expertise. This study aims to compare the possible effect of musical and theatre regular practice on cognition across the lifespan. Both of these artistic activities require many hours of individual or collective training in order to reach an advanced level. This process requires the interaction between higher-order cognitive functions and several sensory modalities (auditory, verbal, visual and motor), as well as regular learning of new pieces. This study included participants with musical or theatre training, and healthy controls matched for age (18 to 84 years old) and education. The objective was to determine whether specific expertise in these activities had an effect on cognition across the lifespan, and a protective influence against undesirable cognitive outcomes associated with aging. All participants underwent a battery of cognitive tasks that evaluated processing speed, executive function, fluency, working memory, verbal and visual long-term memories, and non-verbal reasoning abilities. Results showed that music and theatre artistic practices were strongly associated with cognitive enhancements. Participants with musical training were better in executive functioning, working memory and non-verbal reasoning, whereas participants with regular acting practice had better long-term verbal memory and fluency performance. Thus, taken together, results suggest a differential effect of these artistic practices on cognition across the lifespan. Advanced age did not seem to reduce the benefit, so future studies should focus on the hypothetical protective effects of artistic practice against cognitive decline.

## INTRODUCTION

Without a doubt, musical practice has become a model for the study of neuroplasticity in cognitive neuroscience over the past twenty years (Altenmüller, 2008; Schlaug 2015). It is now accepted that musical expertise leads to cerebral reorganizations resulting in changes in the brain anatomy of regions engaged during formal music learning, such as motor (Wan & Schlaug, 2010), auditory perception (Parbery-Clark et al., 2013; Bidelman et al., 2015; Zendel et al., 2019) and memory areas (Groussard et al., 2010, 2014; Fauvel et al. 2014). Musical practice also influences cognitive functioning, involving better performance on tasks that directly call upon skills explicitly learned during formal music learning (*near transfers*) but also with an effect on general cognitive functions (*far transfers*) in musicians (Schellenberg & Weiss 2013, Fauvel et al., 2013, Schlaug 2015). Studies reported better performance for musicians compared to non-musicians mainly in executive functioning, notably working memory, flexibility and verbal fluency (Degé, Kubicek and Schwarzer, 2011; Criscuolo et al. 2019).

Some authors suggested that beyond musical practice, an active, socially engaged, mentally and physically stimulating lifestyle can also have a positive effect on cognitive functioning (Jung et al. 2017). Brain activity and structure are shaped by experience throughout the lifespan, even at an old age. This plasticity has often been demonstrated after long and intensive trainings, where performance in trained activity improves after practice and leads to the building of a cognitive reserve that could explain the interindividual variability regarding aging (Stern, 2009; Chan et al. 2018). This suggests that higher cognitive reserve is associated with compensatory adjustment and could slow down age-related cognitive decline (Hinault and Lemaire, 2020; Kalpouzos et al. 2008). Different factors influence the variability of this reserve among subjects, including levels of education and general lifestyle (diet and physical fitness), but also the quality of social interaction and hobbies (Scarmeas & Stern 2003).

Like music, theatrical practice is an artistic activity that requires many hours of individual or collective training in order to reach an advanced level. This process requires the interaction between higher-order cognitive functions and several sensory-cognitive modalities (auditory, verbal, visual and motor), as well as the regular learning of new pieces (for musicians see Brown et al. 2015). However, only few studies have investigated the positive effect of artistic activities other than music, such as theatre, on cognition in adulthood. While theatrical practice also seems to have potential effect on overall well-being and cognition, its effect on cognitive functions is still poorly understood. To our knowledge, only Noice's team has conducted a series of studies to specify the effect of theatrical practice on cognitive processes. These studies investigated cognitive changes following short-term theatrical interventions in older adults (Noice & Noice 1999, Noice et al. 2004, Banducci et al. 2017, Noice et al. 2014 for review). They compared older participants who received theatre arts training (n=44), visual arts training (n=44) or no-treatment (controls, n=36) during nine 90-minute sessions over a month. The pretest and posttest comparison suggested that the performance of the theatre arts intervention group was better than no-intervention group on word recall, listening span and problem-solving tasks. Compared to the visual arts group, the theatre arts group performed better on problem solving only (Noice et al. 2004). Recent work from this team further expands these results, as Banducci et al. (2017) compared the cognitive benefit of an active acting program including 86 healthy aging versus 93 participants constituting the control group (history of art classroom) for 4 weeks. A cognitive battery was administered before and after intervention, and again in a 4-month follow-up. The participants of the active acting program benefited most relative to the control group in episodic recall

only, with gains still evident up to 4 months after intervention. Both groups were similar in the magnitude of gains in working memory, executive function and processing speed. Due to the scarcity of work on theatre practice compared to music training in the literature, it seems necessary to specify the benefits of theatrical regular practice on cognition and to better understand its effect throughout life. Many factors appear to influence an individual's aging trajectory (Hinault & Lemaire, 2020; Raz and Kennedy, 2009), suggesting that interventions could possibly slow down cognitive decline and promote healthy aging. To this end, various behavioral interventions have been proposed, such as physical activity and cognitive training (Colcombe and Kramer, 2003; Jaeggi et al., 2008; Karbach and Kray, 2009, Fauvel et al., 2013; Sprague et al. 2019 for reviews), and the benefits of arts practices for promoting health have received growing interest. Importantly, while prior studies have undoubtedly shown the association between arts engagement and well-being (Mella et al. 2017 ; Fancourt and Stepoe 2019), objective measurement of the specific cognitive benefit associated with repeated and regular art practices like acting or music across the lifespan, have not been carried out to the best of our knowledge.

The literature is consistent with results obtained by the first study performed on the link between the practice of a musical instrument and cognitive functions in elderly subjects (Hanna-Pladdy & MacKay 2011). These authors observed that elderly musicians outperformed elderly non-musicians on nonverbal working memory, naming and executive function tests. Moreover, their study suggested a correlation between musicians considered to have a high level of expertise (i.e. having at least more than 10 years of practice) and the preservation of cognitive functioning while aging. Contrarily, Mansen et al. 2017 showed that the time spent making music was not the most important criterion with respect to cognitive function compared with other practice characteristics such as current amount of time making music or age of onset of musical practice.

To our knowledge, no study has been carried out among people engaged in theatre practice in order to evaluate their cognitive abilities and possible reserve effects. However, it seems logical to think that many years of theatre practice could influence cognition and especially memory, as actors have to memorize new texts regularly. Similarly, it seems surprising that actors have never constituted a reference group or a comparison group with musicians. This is probably due to the fact that it is more difficult to define equivalent criteria for the level of expertise for actors whose training and practices are more heterogeneous than for musicians educated in music conservatories.

The main objective of this study was to compare the positive influence of musical and theatrical current and regular practice on cognition. Our goal was first to determine, throughout the lifespan, whether people with current musical and theatrical practice could show cognitive differences, and if the number of years of practice influence these modifications. Second, we aimed to study the effect of musical and theatrical practice on cognition in older adults, in order to assess the specific differences in cognition between actors and musicians while aging (e.g. executives and episodic memory processes).

Participants underwent a battery of cognitive tasks evaluating processing speed, executive functioning, fluency, working memory, verbal and visual long-term memories, and non-verbal reasoning abilities. Considering the literature, the main expectation was that both groups (musicians and actors) would perform higher than control subjects without any intensive and regular leisure activity on tasks evaluating both executive functions and memory. We also expected a specific effect of the type of practice on cognition, with better performance for verbal memory in actors and executive functioning and reasoning in musicians. Moreover, we expected that processes involved in reading scores in musicians

would increase their abilities in visuo-spatial memory. We finally expected that these patterns would be maintained in older adults.

## MATERIALS AND METHODS

### Participants

We recruited three groups of healthy subjects differing only in regular and sustained practice of a specific leisure activity (music; theatre, no specific practice for the control group). The dataset was obtained from 146 participants, 50 controls; 50 “musicians” and 46 “actors” matched for age and education. Three participants (1 by group) were excluded because they only partially completed the neuropsychological assessment (Table 1). All participants were native French speakers, with normal hearing and normal or corrected to normal vision without any reported neurological or psychiatric conditions, as assessed by a medical interview. None of them presented signs of cognitive impairment (i.e. two or more scores in two or more cognitive domains below two standard deviations of the norms for their age class). All participants provided informed consent before being tested and all procedures were conducted in agreement with the ethical principles of Declaration of Helsinki.

**Table 1 : Demographic data of participants and practice background information for Musicians and Actors.**

	Controls	Musicians	Actors	Stats	p-value
<b>N of subjects</b>	49	49	45		
<b>Gender</b>	30F +20 M	24F + 26 M	30F + 16 M	$\chi^2=3.096$	0.213
<b>Age</b>	47.47 $\pm$ 17.78 [18-80]	47.84 $\pm$ 18.3 [20-83]	41.58 $\pm$ 18.13 [18-84]	F= 1.748	0.178
<b>Years of education</b>	13.9 $\pm$ 2.946 [9-20]	14.73 $\pm$ 2.139 [9-19]	13.89 $\pm$ 2.648 [7-20]	F= 1.689	0.188
<b>Age onset of practice</b>	n.a	13.02 $\pm$ 12.82 [2-65]	24.9 $\pm$ 15.34 [4-70]	U MW= 506	<.001
<b>Years of practice</b>	n.a	31.51 $\pm$ 16.67 [5-65]	17.13 $\pm$ 10.82 [4-45]	U MW=539.5	<.001
<b>Weekly practice hours</b>	n.a	13.03 $\pm$ 11.03 [1-49]	10.18 $\pm$ 9.83 [2-40]	U MW=1.318	0.191
<b>Number of exhibition /year</b>	n.a	18.14 $\pm$ 24.84 [0-100]	14.43 $\pm$ 22.14 [0-100]	U MW=0.76	0.448

Controls were defined as participants who had practiced any leisure activity regularly (more than 4 hours/week) associated with formal lessons (physical activity or drawing lesson for examples) and had never taken any formal music or acting lessons, that could neither play nor read music.

Participants were included in “Musicians” group when they reported current and regular practice at the moment of the study for more than 3 years of musical instrument without interruption, more than 4 hours/week, and if they had received formal music training. In addition to this, Musicians had to never have practiced theatre. Musicians were recruited from several French conservatories or music schools (no self-educated musicians were included). They played various musical instruments (piano, guitar, trumpet, etc.). To study the influence of instrumental practice and avoid confounding effect between singing and instrumental practices (e.g. Mansens et al. 2017) we excluded participants who previously performed choral singing in all groups.

Participants were included in “Actors” group when they reported current and regular acting practice at the moment of the study for more than 3 years without interruption, more

than 4 hours/week, and if they had taken formal theatre lessons. In addition to this, Actors must had never practiced a musical instrument. All Actors were recruited from theatre companies or cultural associations which offered acting lessons. Without being professional, they learned new texts on a regular basis and had regular live performances (two per month in average).

Musicians' and Actors' background information is provided in Table 1, which includes the age onset of practice, number of years of practice, weekly practice hours, number of exhibitions by year.

We then reduced our sample to individuals who were 50 years and older to study the difference in cognition between older practitioners (Schneider et al. 2019). This sample was composed of 27 Controls, 24 Musicians and 15 Actors (see tables 2).

**Table 2 : Demographic data of participants of 50 years and older Practice background information for Musicians and Actors of 50 years and older**

	Controls	Musicians	Actors	Stats	p-value
<b>N of subjects</b>	27	24	15		
<b>Gender</b>	18F + 9 M	12F + 12 M	9F + 6 M	$\chi^2 = 1.47$	0.48
<b>Age</b>	61.1 $\pm$ 7.29 [51-80]	63.9 $\pm$ 8.81 [50-83]	62.6 $\pm$ 9.73 [52-84]	F= 0.723	0.489
<b>Years of education</b>	14.5 $\pm$ 2.65 [9-19]	14.2 $\pm$ 1.99 [9-17]	12.9 $\pm$ 2.45 [10-20]	F= 2.31	0.108
<b>Age onset of practice</b>	n.a	18.5 $\pm$ 16.6 [2-65]	37.4 $\pm$ 17 [12-70]	U MW= 69.5	<.001
<b>Years of practice</b>	n.a	43 $\pm$ 15 [12-65]	23.7 $\pm$ 13.4 [4-45]	U MW=60.5	<.001
<b>Weekly practice hours</b>	n.a	12.8 $\pm$ 11.00 [3.5-49]	10.5 $\pm$ 11.8 [3-40]	U MW=130	0.152
<b>Number of exhibition /year</b>	n.a	14.3 $\pm$ 18.6 [0-60]	15.9 $\pm$ 20.4 [2-80]	U MW=142	0.27

## **Cognitive functioning**

Cognitive functioning was measured using several assessments covering various cognitive domains (Table 3) that are clearly impacted by normal aging (Salthouse, 2010; Hedden and Gabrieli, 2004), the earliest and most concerned being processing speed, working memory (maintenance and manipulation of information during a short period of time), spatial ability, reasoning, and episodic memory (declarative long-term contextual remembering of personal events or information). The entire test battery was administered in a single session, which lasted about 90 min, and took place in a quiet room.

Cognitive functions assessed	Psychometric tests	Dependent variables (outcomes)
Long-term verbal memory	BEM-144's 12 words (Signoret, 1991)	Total score of the three trials in learning phase (BEM Total) Number of word recalled in the delayed recall (BEM Recall)
	Doors test (Baddeley et al. 1994)	Number of doors recognized (Doors Total)
Long-term visual memory	Rey-Osterrieth complex figure (Rey, 1959)	Score of redraw fidelity (number of details, their completeness and location) (Rey Recall)
		Highest number of digits properly recalled in 2/3 trials (Digit Span)
Working memory	Forward digit span (Godefroy et al. 2008)	Total number of target symbols correctly identified (GZ-f)
Attentional abilities	d2 Test (Brickenkamp, 1981)	Total number of words in 2 min (Phonemic Fluency)
		Total number of words in 2 min (Semantic Fluency)
Executive control and verbal abilities	Phonemic Fluency task (Cardebat et al. 1990)	
	Semantic Fluency task (Cardebat et al. 1990)	
Processing speed	d2 Test (Brickenkamp, 1981)	Number of items processed (GZ)
	Digit-symbol coding subtest (Wechsler, 2000)	Number of correct associations in 2 minutes (Codes)
Non-verbal reasoning	Matrix Reasoning subtests (Wechsler, 2000)	Number of matrices properly completed (Matrix)

### Long-term Memory:

Long-term verbal memory was measured using the 12-word subtest from the Signoret BEM-144 (Signoret 1991). This verbal memory test consists of learning 12 words during 3 sessions. After every trial, participants are asked to recall as many words as possible. Then participants are distracted by performing a non-verbal task for approximately 7 min. After that, they are asked to recall as many words as possible. We used two scores on the 12-word BEM test: the total score of the three trials to evaluate total learning (BEM Total), and the number of words recalled during the delayed recall (BEM Recall) to assess episodic memory.

Long-term visual memory was evaluated using the Baddeley's Doors test (Baddeley et al. 1994). This test is a nonverbal recognition test based on colored photographs of doors composed of two parts (A and B). For each part, 12 doors are shown individually for 3 seconds. Then, the participants are asked to pick out the one out of the four that had been shown before. The score is the number of correct answers of the two parts combined (Doors Total).

The Rey-Osterrieth complex figure (Rey, 1959) was administered and consists of redrawing an abstract geometrical shape from memory that had been copied 3 minutes earlier (Rey Recall). The maximum final score is 36. This test is classically used for evaluating of visuospatial constructional ability and visual episodic memory.

### Executive Functioning

The phonological loop of the working memory, which is the ability to retain verbal information for a short time by mean of mental repetition (Baddeley, 2003; Baddeley et al., 1998), was evaluated using the forward digit span (Godefroy et al 2008). Participants had to immediately recall series of digits in the order they were presented. The score recorded was the size of the forward digit span with 2 successive correct recalls (Digit Span).

We evaluated visual attention using the d2 Test. This test consists of a paper with 14 rows of 47 interspersed "p" and "d" characters. The participant had to cross out as many "d" with two marks above or below them as possible, in any order (target symbols), and had to jump to the next rows every 15 s. The target symbols are relatively similar to the distractors (a



“p” with two marks or a “d” with one or three marks). In this study, we used the overall performance score (GZ-f) corresponding to the total number of target symbols correctly identified (Brickenkamp, 1981).

The phonemic fluency task (Cardebat *et al.* 1990) was used to measure executive functioning. The participant had to name as many words starting with the letter R as possible in two minutes. The score used is the total number of words in two minutes. The semantic fluency was also proposed, in which participants had to name as many fruits as possible in two minutes. Language processing and semantic memory are most the critical components for this task. .

Processing speed was measured using the digit-symbol coding subtest from the Weschler Adult Intelligence Scale (WAIS-III, Weschler, 2000). In this task, each digit (from 1 to 9) was combined with a specific symbol (example 1/- and 9/=) in the upper row. Participants then had 2 minutes to complete the number maximum of symbols corresponding to the digits presented in the lower rows. The score was the number of correct associations performed in 2 minutes (Codes). We also used the number of items processed (Gz) in the d2 test (Brickenkamp, 1981) to evaluate the processing speed.

### ***Reasoning***

We administered the Matrix Reasoning subtests of the WAIS-III (Wechsler, 2000) to estimate participants’ non-verbal reasoning skills. In this test, participants were presented with an unfinished matrix of drawings, and had to choose the drawing that logically completed the matrix. This task is classically associated with fluid intelligence (Matrix) (Carpenter, Just and Shell 1990). We used the number of matrices properly completed as a performance score.

### **Procedure and Statistical Analysis**

To compare all cognitive variables with each other, scores were standardized. Thus, we transformed all neuropsychological measures into z-scores using the mean and standard deviation of the control groups (n=49) as the reference population for each measure due to the lack or poor reliability of French published norms for some assessments. Thus, all variables were on the same scale, with a mean of 0 and a standard deviation of 1 based on the control group. The higher the z-score, the better the performance. This allowed comparing every performance on the same normalized scale.

In order to test for group difference among cognitive tests, we performed multivariate analyses of covariance (MANCOVA), with type of practice (Controls; Musicians; Actors) as between-subjects factor, the cognitive test scores (BEM Total, BEM Recall, Doors Total, Rey Recall, Digit Span, Matrix, Codes, d2 GZ, d2 GZ-f, Phonemic Fluency, Semantic Fluency) as dependent variables and age as confounding variable. As we included more than two dependent variables and because they are intercorrelated, we opted for a MANCOVA. This statistical analysis accounts for the relationship between dependent variables (Warne, 2014). To complete the multivariate analyses and examine group differences for each cognitive variable, univariate analyses were performed. A family-wise Bonferroni’s correction for multiple comparison analyses was carried out 2-by-2 for every significant test.

To further our exploration of the relationship between musical or theatrical practice and cognitive abilities, we performed a second multivariate analysis on cognitive variables restricted to Musicians and Actors and including Age and Years of practice as variables.

In a separate analysis, the same procedure was implemented reducing the sample to people who were 50 years and older to study the effect of expertise on cognition in older adults.

The sample size was based on a power analysis, conducted in G\*Power 3.1. Regarding behavioral interactions between age and cognition, assuming an effect size of Cohen's  $f=0.65$  (derived from Carey et al., 2019), an alpha of .05, and three groups, we determined that a total sample size of at least 15 individuals per study would provide 95% power to detect the effects.

All the statistical analyses were performed with STATISTICA software. The partial Eta square ( $\eta^2_p$ ) was utilized to estimate effect size. Results were considered significant at  $p<.05$ .

## RESULTS

### *Effect of expertise on cognitive variables in adults*


Results from the multivariate tests on the associations between groups of practice (Controls; Musicians; Actors) and cognitive tests adjusted for age exhibited a significant group effect: Wilks's Lambda [ $F_{(22,258)}=3,005$ ,  $p=0.000015$   $\eta^2_{partial}=0.204$ ]. Results showed a significant effect of groups of practice with higher values for Musicians relative to Controls for Rey Recall ( $p=0.024$ ); Codes ( $p=0.002$ ); d2 GZ ( $p=0.023$ ); d2 GZ-f ( $p=0.023$ ); Phonemic Fluency ( $p=0.002$ ), Semantic Fluency ( $p=0.019$ ); relative to Actors for Matrix ( $p=0.006$ ); and compared to both Controls and Actors for Span (*respectively*  $p=0.0002$ ;  $p=0.008$ ), . Results also highlighted significantly higher values for Actors than Controls for BEM Total ( $p=0.002$ ), and Phonemic Fluency ( $p=0.009$ ), and compared to both Controls and Musicians for BEM Recall (*respectively*  $p=0.001$ ;  $p=0,033$ )(Table 4, Figure 1).

The multivariate analysis exhibited an effect of the Age: Wilks's Lambda [ $F_{(11,129)}=5.286$ ,  $p=0.000001$ ], and univariate analyses suggested a significant effect of the Age on BEM Total [ $F_{(1,139)}=12.49$ ,  $p=0.0005$ ], BEM Recall [ $F_{(1,139)}=11.32$ ,  $p=0.001$ ], Rey Recall [ $F_{(1,139)}=12.319$ ,  $p=0.0006$ ], Matrix [ $F_{(1,139)}=13.85$ ,  $p=0.0003$ ], Codes [ $F_{(1,139)}=38.56$ ,  $p<0.001$ ], d2 GZ [ $F_{(1,139)}=17.85$ ,  $p=0.0004$ ], and d2 GZ-f [ $F_{(1,139)}=20.77$ ,  $p=0.0001$ ].

**Table 4 : Statistical results of MANCOVA for each cognitive variable. SD= Standard Deviation**

	Controls	Musicians	Actors	Statistics		
	Mean±SD	Mean±SD	Mean±SD	F	p-value	Post-hoc
<b>BEM Total</b>	25.673 ± 4.451	26.857 ± 4.21	28.778 ± 4.617	<b>4.450</b>	<b>0.0134</b>	Actors > Controls
<b>BEM Recall</b>	8.449 ± 2.39	8.939 ± 2.277	10.067 ± 1.876	<b>5.256</b>	<b>0.006</b>	Actors > Controls Actors > Musicians
<b>Doors Total</b>	17.673 ± 2.593	18.082 ± 2.448	18.40 ± 2.934	0.632	0.533	
<b>Rey Recall</b>	20.806 ± 4.716	23.592 ± 4.745	23.067 ± 6.463	<b>3.763</b>	<b>0.026</b>	Musicians > Controls
<b>Digit Span</b>	5.592 ± 1.29	6.551 ± 1.174	5.822 ± 0.960	<b>9.449</b>	<b>0.0001</b>	Musicians > Controls Musicians > Actors
<b>Matrix</b>	20.531 ± 2.792	21.735 ± 2.564	20.289 ± 2.928	<b>5.328</b>	<b>0.006</b>	Musicians > Actors
<b>Codes</b>	69.184 ± 17.257	78.776 ± 16.37	73.222 ± 12.269	<b>6.906</b>	<b>0.001</b>	Musicians > Controls
<b>d2 GZ</b>	391.306 ± 72.378	431.327 ± 80.379	401.556 ± 79.371	<b>4.729</b>	<b>0.010</b>	Musicians > Controls
<b>d2 GZ-f</b>	376.878± 65.827	413.551 ± 77.785	382.422 ± 71.237	<b>5.285</b>	<b>0.006</b>	Musicians > Controls
<b>Phonemic Fluency</b>	21.408 ± 5.733	25.755 ± 5.445	25.20 ± 7.191	<b>7.619</b>	<b>0.0007</b>	Musicians > Controls Actors > Controls
<b>Semantic Fluency</b>	21.265 ± 4.420	23.796 ± 4.509	23 ± 4.661	<b>3.937</b>	<b>0.022</b>	Musicians > Controls

## Z-scores for each cognitive variable studied



**Figure 1 : median, 1<sup>st</sup> and 3<sup>rd</sup> quartiles, min and max z-scores for each cognitive variable studied. Mean and standard deviation of the control group (n=49) serves as reference population for each measure, as such variables were on the same scale with 0 as the mean and 1 as the standard deviation of the control group. The higher the z-score, the better the performance. (\*) after a test name indicates a significative difference between Controls and Musicians or Actors.**

### *Effect of years of practice on cognitive variable of adult Musicians and Actors:*

The multivariate analysis exhibited a significant effect of groups of practice, Wilks's Lambda [ $F_{(11,80)}=4,058, p=0.0001 \eta^2_{\text{partial}}=0.358$ ], a significant effect of Age, Wilks's Lambda [ $F_{(11,80)}=2,957, p=0.0025 \eta^2_{\text{partial}}=0.289$ ] and no effect of Years of practice on cognitive variables Wilks's Lambda [ $F_{(11,80)}=1,776, p=0.072 \eta^2_{\text{partial}}=0.196$ ]. The univariate analyses on groups of practice confirmed the higher values for Actors relative to Musicians after controlling of Age and Years of practice for BEM Total ( $p=0.039$ ) and BEM Recall ( $p=0.0046$ ) and higher values for Musicians relative to Actors for Span ( $p=0.0023$ ), Matrix ( $p=0,022$ ), Codes ( $p=0,0109$ ), d2 GZ ( $p=0,0404$ ) and d2 GZ-f ( $p=0,0164$ ).

### *Effects of expertise on cognitive variables in older adults*


In a separate analysis, we reduced our sample to participant who were 50 years and older to study the effects of expertise on cognitive aging. Results from the multivariate tests that studied the associations between groups of practice (Controls; Musicians; Actors) and the cognitive tests adjusted for Age exhibited a significant group effect: Wilks's Lambda

[ $F_{(22,104)}=1.732$ ,  $p=0.0347$   $\eta^2_{partial}=0.268$ ]. Results of univariate tests showed a significant effect on groups of practice, with significantly higher values in Musicians relative to Controls for Rey Recall ( $p=0.025$ ), Digit Span ( $p=0.018$ ), Codes ( $p=0.026$ ) and Semantic Fluency ( $p=0.005$ ). No significantly higher values in Actors relative to Controls and difference between Actors and Musicians (Table 5, Figure 2). The multivariate an effect of age, Wilks's Lambda [ $F_{(11,52)}=2.869$   $p=0.005$ ,  $\eta^2=0.378$ ] and univariate analyses suggested a significant effect of age on Doors [ $F_{(1,62)}=12.82$   $p=0.0007$ ], Matrix [ $F_{(1,62)}=5.216$   $p=0.026$ ], Codes [ $F_{(1,62)}=14.45$   $p=0.0003$ ], Phonemic Fluency [ $F_{(1,62)}=4.614$   $p=0.036$ ].

**Table 5 : Statistical results of MANCOVA limited to older adults. SD= Standard Deviation**

	Controls	Musicians	Actors	Statistics		
	Mean±SD	Mean±SD	Mean±SD	F	p-value	Post-hoc
<b>BEM Total</b>	24.8 ± 4.03	26 ± 4.20	26.1 ± 3.81	1.039	0.359	
<b>BEM Recall</b>	8 ± 2.43	8.46 ± 2.32	9 ± 1.69	1.065	0.351	
<b>Doors Total</b>	17.6 ± 2.61	18 ± 2.51	17.9 ± 3.66	0.62	0.541	
<b>Rey Recall</b>	18.8 ± 4.51	22.6 ± 4.26	20.8 ± 7.01	<b>4.413</b>	<b>0.016</b>	Musicians > Controls
<b>Digit Span</b>	5.44 ± 1.19	6.38 ± 1.28	5.87 ± 0.915	<b>4.428</b>	<b>0.016</b>	Musicians > Controls
<b>Matrix</b>	19.6 ± 2.49	20.6 ± 2.18	19.6 ± 3.68	1.541	1.756	
<b>Codes</b>	61.3 ± 15.30	71.1 ± 13	64.3 ± 13.9	<b>5.367</b>	<b>0.007</b>	Musicians > Controls
<b>d2 GZ</b>	380 ± 59.60	401 ± 79.5	366 ± 64.5	1.632	0.204	
<b>d2 GZ-f</b>	365 ± 50.70	383 ± 76.5	347 ± 56.8	1.884	0.161	
<b>Phonemic Fluency</b>	22.6 ± 6.23	26.3 ± 6.81	26.6 ± 8.48	<b>3.261</b>	<b>0.045</b>	
<b>Semantic Fluency</b>	20.1 ± 4.67	24.3 ± 4.44	22.1 ± 4.83	<b>5.754</b>	<b>0.005</b>	Musicians > Controls

## Z-scores for each cognitive variable studied in older participants



**Figure 2 :** median, 1<sup>st</sup> and 3<sup>rd</sup> quartiles, min and max z-scores of the older participants for each cognitive variable studied. Mean and standard deviation of the control group (n=49) serves as reference population for each measure, as such variables were on the same scale with 0 as the mean and 1 as the standard deviation of the control group. The higher the z-score, the better the performance. (\*) after a test name indicates a significative difference between Controls and Musicians or Actors

### *Effect of years of practice on cognitive variables of older Musicians and Actors*

The multivariate analysis exhibited only a significant effect of Age, Wilks's Lambda [ $F_{(11,25)}=2,675$ ,  $p=0.020$   $\eta^2_{\text{partial}}=0.541$ ]. No effect of Years of practice on cognitive variables Wilks's Lambda [ $F_{(11,25)}=1.396$ ,  $p=0.235$   $\eta^2_{\text{partial}}=0.381$ ] and Groups of practice were observed, Wilks's Lambda [ $F_{(11,25)}=0.614$ ,  $p=0.219$   $\eta^2_{\text{partial}}=0.387$ ].

## **DISCUSSION**

The aim of the present study was to investigate possible differences in cognition between different art practitioners ((Musicians and Actors), and also people without artistic training. Several studies have demonstrated the benefits of musical practice on cerebral

activity and cognition, but comparing this practice to another artistic practice such as theatre had yet to be done. Our results are consistent with previous works suggesting that adult musicians outperformed control subjects in standardized cognitive tasks (Fauvel et al. 2013; Schlaug 2015; Sutcliffe et al. 2020) but a lifespan approach was never adopted in anterior studies. In fact, previous work has focused on younger musicians or older musicians whereas our sample had a very wide age range (from 18 to 84 years), allowing us to study practitioners' cognitive differences throughout the lifespan. We observed a superiority of musicians in long-term visual-spatial memory, working memory, processing speed, executive functioning and non-verbal reasoning. Nevertheless, Musicians did not seem to outperform Controls on verbal episodic memory and visual memory. This result appears consistent with result of a meta-analysis performed by Talamini et al. (2017) on memory. In fact, these authors suggested a small effect size for long-term memory and a possible domain-specific stimuli effect in favor of musical stimuli (Talamini et al. 2017).

Nevertheless, we also found a difference in the theatre group on cognition across the lifespan. In line with results of Noice et al. (2004) obtained after a 4-month theatrical intervention, we observed a better long-term verbal memory and verbal fluency in these subjects, compared to controls and musicians. Our results were observed in a large sample of younger to older adults that presented a sustainable practice and intense training. Actors' better performance in verbal episodic memory is consistent with the abilities developed by them while learning a text and retrieving it during performance. In fact, most actors use mnemotechnical strategies to encode and retrieve their scripts (Banducci et al 2017, Noice et al. 2004). Strategies were indeed found to improve memory in both young and older adults (e.g., Hinault et al., 2017a, 2017b). However, Actors did not show a difference on executive functioning, working memory and non-verbal reasoning when compared to Control participants. Thus, in future studies it would seem relevant to consider a more detailed evaluation cognition (including strategy use) in order to confirm whether these effects are limited to verbal and memory aspects and do not influence executive processes, or whether self-monitoring abilities are required in theatre practice (Nettle 2006).

This work is the first to statistically compare these two artistic practices, with the same cognitive assessment battery. It highlights for the first time that both musical and theatrical practices could lead to differences in cognition across the lifespan, confirming previous studies on leisure activities, lifestyle and cognition (Hertzog et al. 2008; Reuter-Lorenz & Park 2014). Furthermore, we observed domain-specific differences, musical practice being associated with better executive functions and reasoning, and theatrical practice with better long-term verbal memory. In fact, Musicians had better performance when compared with Actors on working memory, processing speed, executive functioning, and non-verbal reasoning whereas Actors outperformed Musicians for long-term verbal memory.

In older adults, this pattern seems to be confirmed for musicians, with higher performances on long-term visual-spatial memory, working memory, processing speed and verbal fluency (Hanna-Pladdy and MacKay 2011; Hanna-Pladdy & Gajewski 2012; Amer et al. 2013; Fauvel et al. 2014; Mansens et al. 2018; Ferreri et al. 2019; Criscuolo et al. 2019). The EEG study of Moussard *et al.* (2016) on elderly musicians (currently practicing about 11 hours/week) and non-musicians, confirmed a beneficial effect of musical practice on executive control, and highlighted a more anterior distribution of the P3 wave in musicians, suggesting successful functional reorganization in elderly musicians according to the authors. Moreover, longitudinal studies showed that 6 months of piano lessons given to older non-musicians adults could improve working memory and executive functioning (Bugos et al. 2007, Seinfeld et al. 2013). In older actors, Banducci *et al.* (2017) reported modifications on

verbal long term-memory and fluency after a 4-month theatrical intervention in older adults which suggests a cognitive benefit even after a short period of active art practice. While we reported better performances on verbal long-term memory and fluency for Actors compared with Controls, we could not find any significant difference in the elderly for Actors relative to Musicians. These results must to be taken with caution with regard to the small sample size of the older Actors (n=15) and would require further investigation to confirm stronger verbal cognition associated with theatrical practice in aging.

In line with several studies on musicians that suggested no association between practice time and cognitive functions (e.g., Fauvel et al. 2014; Mansen et al. 2017), our results did not reveal any effect of the number of years of practice on assessed cognitive functioning. Thus, having a regular and current practice appears to better explain the cognitive differences we studied rather than years of practice. These results are interesting, as even a short period of practice can lead to an improvement in cognitive performance in adults across the lifespan. There is a growing consensus toward aging brains remaining plastic and consequently involvement in leisure activities such as music or theatre remains of significant interest since it is possible to start this type of practice at any age (Noice et al. 2014).

These findings are constrained by several limitations that need to be considered in future research. First, our study is essentially descriptive because of its correlational approach and does not allow us to validate causality. Only future interventional or follow-up studies could confirm these results. Second, we partially evaluated the working memory abilities because working memory updating ability was not assessed, as the digit backward span was not among the cognitive assessment. Future studies could aim at specifying music and theatrical practice differential effects on this cognitive process. Third, musicians and actors differed in their average years of practice, musicians showing a longer practice duration than actors. Although this variable was included as covariate in analyses without significant interactions, it could have explained some of the cognitive differences between our groups. Four, global cognitive functioning was not assessed, but no participant was below the passing score in more than one cognitive measure, in line with preserved overall cognitive performance. Furthermore, although power analyses and previous work support the selected sample size, future studies should investigate cognitive differences between older musicians and actors with a larger sample size.

To conclude, our results suggest that artistic practices can account for different individuals' aging trajectories (Raz & Kenedy, 2009), and that regular artistic practice could promote the constitution of cognitive and cerebral reserve (Stern, 2009; Stern et al. 2018). Therefore, promoting access to artistic practice could help people maintaining or even improving their cognition, besides the obvious and well-documented interest such activities have on socialization (Belgrave, 2011); well-being (Noice et al. 2004; Castora-Blincket et al. 2010) and developing creativity (Reynolds et al. 2016; Salimpoor & Zatorre 2013). In line with the evidence reviewed by Sutcliffe et al. (2020) on music training and cognition on aging, our study suggests that musical or theatrical practices, even started late in life, could have an effect on cognitive decline. Ferreira et al. (2015) suggested associations between specific activities and the functioning of individual cognitive domains. Results suggest that cognitive training programs could be individually adjusted to observed cognitive deficits following a neuropsychological assessment, without making it a unique criterion for choosing such activity of course. However, previous works on cognitive interventions for aging and dementia showed mixed results (e.g., Alves et al., 2013). In future clinical researches it would be interesting to determine whether theatrical interventions could improve language and episodic memory processes for people with deficits in these domains. Conversely, as both previous works and our results highlight the possible positive effect of musical practice on

executive functioning (Mansens et al. 2018; Koshimori and Thaut 2019; Platel & Groussard 2020), it seems relevant to specify the effect of musical practice interventions for people with deficits in these processes.

**Acknowledgments:** We would like to thank Baptiste Fauvel and the master students of Caen Normandy University for their help in collecting data and their role in this study. Many thanks to Melanie Ambler for proof reading and reviewing English. The first author thanks Arthur and Robin Pesnel for their interventions and discussions during writing.

**Authors Statement:** MG and HP conceived the study and coordinated the collect of data. MG and TH performed statistical analysis. MG wrote the first draft of the manuscript and RC, TH and HP revised the different version of the manuscript. All authors approved the submitted version.

## REFERENCES

- Altenmüller, E. (2008). Neurology of musical performance. *Clinical Medicine (London, England)*, 8(4), 410–413. <http://www.ncbi.nlm.nih.gov/pubmed/18724609>
- Alves, J. (2013). Non-pharmacological cognitive intervention for aging and dementia: Current perspectives. *World Journal of Clinical Cases*, 1(8), 233. <https://doi.org/10.12998/wjcc.v1.i8.233>
- Amer, T., Kalender, B., Hasher, L., Trehub, S. E., & Wong, Y. (2013). Do Older Professional Musicians Have Cognitive Advantages? *PLoS ONE*, 8(8), e71630. <https://doi.org/10.1371/journal.pone.0071630>
- Baddeley, A. (1994). The magical number seven: Still magic after all these years? *Psychological Review*, 101(2), 353–356. <https://doi.org/10.1037/0033-295X.101.2.353>
- Baddeley A, Gathercole S, Papagno C (1998) The phonological loop as a language learning device. *Psychol Rev* 105, 158–173.
- Baddeley A (2003) Working memory: looking back and looking forward. *Nat Rev Neurosci* 4, 829–839.
- Banducci, S. E., Daugherty, A. M., Biggan, J. R., Cooke, G. E., Voss, M., Noice, T., Noice, H., & Kramer, A. F. (2017). Active Experiencing Training Improves Episodic Memory Recall in Older Adults. *Frontiers in Aging Neuroscience*, 9. <https://doi.org/10.3389/fnagi.2017.00133>
- Bidelman GM, Alain C. (2015) Musical training orchestrates coordinated neuroplasticity in auditory brainstem and cortex to counteract age-related declines in categorical vowel perception. *J Neurosci*, 35(3):1240-1249. doi:10.1523/JNEUROSCI.3292-14.2015
- Belgrave, M. (2011). The Effect of a Music Therapy Intergenerational Program on Children and Older Adults' Intergenerational Interactions, Cross-Age Attitudes, and Older Adults' Psychosocial Well-Being. *Journal of Music Therapy*, 48(4), 486–508. <https://doi.org/10.1093/jmt/48.4.486>
- Brickenkamp, R. (1981). Test d2 - Aufmerksamkeits-Belastungs-Test. *Handbuch*



- 535 *Psychologischer Und Pädagogischer Tests*, 270–273. [http://library.mpib-](http://library.mpib-berlin.mpg.de/toc/z2008_2431.pdf)  
536 [berlin.mpg.de/toc/z2008\\_2431.pdf](http://library.mpib-berlin.mpg.de/toc/z2008_2431.pdf)
- 537 Brown, R. M., Zatorre, R. J., & Penhune, V. B. (2015). *Expert music performance: cognitive,*  
538 *neural, and developmental bases* (pp. 57–86). <https://doi.org/10.1016/bs.pbr.2014.11.021>
- 539 Bugos, J. A., Perlstein, W. M., McCrae, C. S., Brophy, T. S., & Bedenbaugh, P. H. (2007).  
540 Individualized Piano Instruction enhances executive functioning and working memory in  
541 older adults. *Aging & Mental Health*, 11(4), 464–471.  
542 <https://doi.org/10.1080/13607860601086504>
- 543 Cardebat, D ; Doyon, B; Puel, M; Goulet, P; Joannette, Y. (1990). Formal and semantic lexical  
544 evocation in normal subjects. Performance and dynamics of production as a function of  
545 sex, age and educational level. *Acta Neurologica Belgica*, 90(4), 207–217.  
546 <https://europemc.org/article/med/2124031>
- 547 Carpenter, P. A., Just, M. A., & Shell, P. (1990). What one intelligence test measures: a  
548 theoretical account of the processing in the Raven Progressive Matrices Test.  
549 *Psychological Review*, 97(3), 404.
- 550 Castora-Binkley, M., Noelker, L., Prohaska, T., & Satariano, W. (2010). Impact of arts  
551 participation on health outcomes for older adults. *Journal of Aging, Humanities, and the*  
552 *Arts*, 4(4), 352–367. <https://doi.org/10.1080/19325614.2010.533396>
- 553 Chan, D., Shafto, M., Kievit, R., Matthews, F., Spink, M., Valenzuela, M. and Henson, R.  
554 (2018). Lifestyle activities in mid-life contribute to cognitive reserve in late-life,  
555 independent of education, occupation, and late-life activities. *Neurobiology of Aging*, 70,  
556 pp.180-183
- 557 Colcombe, S., & Kramer, A. F. (2003). Fitness Effects on the Cognitive Function of Older  
558 Adults. *Psychological Science*, 14(2), 125–130. [https://doi.org/10.1111/1467-9280.t01-1-](https://doi.org/10.1111/1467-9280.t01-1-01430)  
559 [01430](https://doi.org/10.1111/1467-9280.t01-1-01430)
- 560 Creech, A., Hallam, S., Varvarigou, M., McQueen, H., & Gaunt, H. (2013). Active music  
561 making: a route to enhanced subjective well-being among older people. *Perspectives in*  
562 *Public Health*, 133(1), 36–43. <https://doi.org/10.1177/1757913912466950>
- 563 Criscuolo, A., Bonetti, L., Särkämö, T., Kliuchko, M., & Brattico, E. (2019). On the  
564 association between musical training, intelligence and executive functions in adulthood.  
565 *Frontiers in Psychology*, 10(JULY), 1–12. <https://doi.org/10.3389/fpsyg.2019.01704>
- 566 Degé, F., Kubicek, C., & Schwarzer, G. (2011). Music lessons and intelligence: A relation  
567 mediated by executive functions. *Music Perception*, 29(2), 195–201.  
568 <https://doi.org/10.1525/mp.2011.29.2.195>
- 569 Fancourt, D., & Steptoe, A. (2019). The art of life and death: 14 year follow-up analyses of  
570 associations between arts engagement and mortality in the English Longitudinal Study of  
571 Ageing. *BMJ*, l6377. <https://doi.org/10.1136/bmj.l6377>
- 572 Fauvel, B., Groussard, M., Eustache, F., Desgranges, B., & Platel, H. (2013). Neural  
573 implementation of musical expertise and cognitive transfers: could they be promising in  
574 the framework of normal cognitive aging? *Frontiers in Human Neuroscience*, 7.  
575 <https://doi.org/10.3389/fnhum.2013.00693>
- 576 Fauvel, B., Groussard, M., Mutlu, J., Arenaza-Urquijo, E. M., Eustache, F., Desgranges, B., &  
577 Platel, H. (2014). Musical practice and cognitive aging: two cross-sectional studies point  
578 to phonemic fluency as a potential candidate for a use-dependent adaptation. *Frontiers in*  
579 *Aging Neuroscience*, 6, 227. <https://doi.org/10.3389/fnagi.2014.00227>

- 580 Ferreira, N., Owen, A., Mohan, A., Corbett, A., & Ballard, C. (2015). Associations between  
581 cognitively stimulating leisure activities, cognitive function and age-related cognitive  
582 decline. *International Journal of Geriatric Psychiatry*, 30(4), 422–430.  
583 <https://doi.org/10.1002/gps.4155>
- 584 Ferreri, L., Moussard, A., Bigand, E., Schellenberg, E. G., Ferreri, L., Moussard, A., Bigand,  
585 E., & Schellenberg, E. G. (2019). Music and the Aging Brain. *The Oxford Handbook of*  
586 *Music and the Brain*, November, 622–644.  
587 <https://doi.org/10.1093/oxfordhb/9780198804123.013.25>
- 588 Godefroy, O., Jeannerod, M., Allain, P., & Le Gall, D. (2008). Frontal lobe, executive  
589 functions and cognitive control. *Revue Neurologique*, 164(SUPPL. 3), 119–127.  
590 [https://doi.org/10.1016/S0035-3787\(08\)73302-2](https://doi.org/10.1016/S0035-3787(08)73302-2)
- 591 Godefroy, Olivier, Gibbons, L., Diouf, M., Nyenhuis, D., Roussel, M., Black, S., &  
592 Bugnicourt, J. M. (2014). Validation of an integrated method for determining cognitive  
593 ability: Implications for routine assessments and clinical trials. *Cortex*, 54(1), 51–62.  
594 <https://doi.org/10.1016/j.cortex.2014.01.016>
- 595 Groussard, M., La Joie, R., Rauchs, G., Landeau, B., Chételat, G., Viader, F., Desgranges, B.,  
596 Eustache, F., & Platel, H. (2010). When Music and Long-Term Memory Interact: Effects  
597 of Musical Expertise on Functional and Structural Plasticity in the Hippocampus. *PLoS*  
598 *ONE*, 5(10), 8.  
599 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2950159&tool=pmcentrez&](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2950159&tool=pmcentrez&rendertype=abstract)  
600 [endertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2950159&tool=pmcentrez&rendertype=abstract)
- 601 Groussard, M., Viader, F., Landeau, B., Desgranges, B., Eustache, F., & Platel, H. (2014).  
602 The effects of musical practice on structural plasticity: The dynamics of grey matter  
603 changes. *Brain and Cognition*, 90, 174–180. <https://doi.org/10.1016/j.bandc.2014.06.013>
- 604 Hanna-Pladdy, B., & Gajewski, B. (2012). Recent and Past Musical Activity Predicts  
605 Cognitive Aging Variability: Direct Comparison with General Lifestyle Activities.  
606 *Frontiers in Human Neuroscience*, 6(July), 198.  
607 <https://doi.org/10.3389/fnhum.2012.00198>
- 608 Hanna-Pladdy, B., & MacKay, A. (2011). The relation between instrumental musical activity  
609 and cognitive aging. *Neuropsychology*, 25(3), 378–386.  
610 <https://doi.org/10.1037/a0021895>
- 611 Hedden, T., & Gabrieli, J. D. E. (2004). Insights into the ageing mind: A view from cognitive  
612 neuroscience. In *Nature Reviews Neuroscience* (Vol. 5, Issue 2, pp. 87–96).  
613 <https://doi.org/10.1038/nrn1323>
- 614 Hertzog, C., Kramer, A. F., Wilson, R. S., & Lindenberger, U. (2008). Enrichment Effects on  
615 Adult Cognitive Development. *Psychological Science in the Public Interest*, 9(1), 1–65.  
616 <https://doi.org/10.1111/j.1539-6053.2009.01034.x>
- 617 Hinault, T., Lemaire, P., & Touron, D. (2017a). Aging effects in sequential modulations of  
618 poorer-strategy effects during execution of memory strategies. *Memory*, 25(2), 176–186.  
619 <https://doi.org/10.1080/09658211.2016.1146300>
- 620 Hinault, T., Lemaire, P., & Touron, D. (2017b). Strategy combination during execution of  
621 memory strategies in young and older adults. *Memory*, 25(5), 619–625.  
622 <https://doi.org/10.1080/09658211.2016.1200626>
- 623 Hinault, T., & Lemaire, P. (2020). Aging effects on brain and cognition: What do we learn  
624 from a strategy perspective? In A. Gutchess, & A. Thomas (Eds.), *The Cambridge*

- 625 *Handbook of Cognitive Aging: A life course perspective*. Cambridge: Cambridge  
626 University Press.
- 627 Jaeggi, S. M., Buschkuhl, M., Jonides, J., & Perrig, W. J. (2008). Improving fluid  
628 intelligence with training on working memory. *Proceedings of the National Academy of*  
629 *Sciences*, 105(19), 6829–6833. <https://doi.org/10.1073/pnas.0801268105>
- 630 Jung, M. S., Kim, H., Lee, Y., Kim, M., & Chung, E. (2017). Different effects of cognitive  
631 and non-exercise physical leisure activities on cognitive function by age in elderly  
632 Korean individuals. *Osong Public Health and Research Perspectives*, 8(5), 308–317.  
633 <https://doi.org/10.24171/j.phrp.2017.8.5.04>
- 634 Kalpouzos, G., Eustache, F., & Desgranges, B. (2008). Réserve cognitive et fonctionnement  
635 cérébral au cours du vieillissement normal et de la maladie d'Alzheimer [Cognitive  
636 reserve and neural networks in normal aging and Alzheimer's disease]. *Psychologie &*  
637 *Neuropsychiatrie Du Vieillessement*, 6(2), 97–105.  
638 <https://doi.org/10.1684/pnv.2008.0120>
- 639 Karbach, J., & Kray, J. (2009). How useful is executive control training? Age differences in  
640 near and far transfer of task-switching training. *Developmental Science*, 12(6), 978–990.  
641 <https://doi.org/10.1111/j.1467-7687.2009.00846.x>
- 642 Koshimori, Y., & Thaut, M. H. (2019). New Perspectives on Music in Rehabilitation of  
643 Executive and Attention Functions. *Frontiers in Neuroscience*, 13(November), 1–6.  
644 <https://doi.org/10.3389/fnins.2019.01245>
- 645 Mansens, D., Deeg, D. J. H., & Comijs, H. C. (2018). The association between singing and/or  
646 playing a musical instrument and cognitive functions in older adults. *Aging and Mental*  
647 *Health*, 22(8), 964–971. <https://doi.org/10.1080/13607863.2017.1328481>
- 648 Mella, N., Grob, E., Döll, S., Ghisletta, P., & de Ribaupierre, A. (2017). Leisure activities and  
649 change in cognitive stability: A multivariate approach. *Brain Sciences*, 7(3), 8–11.  
650 <https://doi.org/10.3390/brainsci7030027>
- 651 Moussard, A., Bermudez, P., Alain, C., Tays, W., & Moreno, S. (2016). Life-long music  
652 practice and executive control in older adults: An event-related potential study. *Brain*  
653 *Research*, 1642, 146–153. <https://doi.org/10.1016/j.brainres.2016.03.028>
- 654 Nettle, D. (2006). Psychological profiles of professional actors. *Personality and Individual*  
655 *Differences*, 40(2), 375–383. <https://doi.org/10.1016/j.paid.2005.07.008>
- 656 Noice, H., & Noice, T. (1999). Long-term Retention of Theatrical Roles. *Memory*, 7(3), 357–  
657 382. <https://doi.org/10.1080/096582199387977>
- 658 Noice, H., Noice, T., & Staines, G. (2004). A Short-Term Intervention to Enhance Cognitive  
659 and Affective Functioning in Older Adults. *Journal of Aging and Health*, 16(4), 562–  
660 585. <https://doi.org/10.1177/0898264304265819>
- 661 Noice, T., Noice, H., & Kramer, A. F. (2014). Participatory arts for older adults: A review of  
662 benefits and challenges. *Gerontologist*, 54(5), 741–753.  
663 <https://doi.org/10.1093/geront/gnt138>
- 664 Parbery-Clark A, Strait DL, Hittner E, Kraus N. (2013) Musical training enhances neural  
665 processing of binaural sounds. *J Neurosci*, 33(42):16741-16747.  
666 doi:10.1523/JNEUROSCI.5700-12.2013
- 667 Platel H, Groussard M, (2020) Benefits and limits of musical interventions in pathological  
668 aging. In Cuddy L., Belleville S., Moussard A. “Music and the aging brain”, Academic Press.

669 Raz, N., and Kennedy, K. M. (2009). “A systems approach to the aging brain: neuroanatomic  
670 changes, their modifiers and cognitive correlates,” in *Imaging the Aging Brain*, eds W.  
671 Jagust and M. D’Esposito (New York, NY: Oxford University Press), 43–70.

672 Reuter-Lorenz, P. A., & Park, D. C. (2014). How Does it STAC Up? Revisiting the  
673 Scaffolding Theory of Aging and Cognition. *Neuropsychology Review*, 24(3), 355–370.  
674 <https://doi.org/10.1007/s11065-014-9270-9>

675 Rey,A.(1959). Manuel de la Figure Complexe de Rey. Paris: Editions du Centre de  
676 Psychologie Appliquée .vel. *Acta Neurol.Belg.* 90,207–217.

677 Reynolds, J., Bernard, M., Rezzano, J., & Rickett, M. (2016). Ageing, Drama, and Creativity:  
678 Translating Research Into Practice. *Gerontology and Geriatrics Education*, 37(3), 307–  
679 328. <https://doi.org/10.1080/02701960.2016.1152267>

680 Salimpoor, V. N., & Zatorre, R. J. (2013). Neural interactions that give rise to musical  
681 pleasure. *Psychology of Aesthetics, Creativity, and the Arts*, 7(1), 62–75.  
682 <https://doi.org/10.1037/a0031819>

683 Salthouse, T. A. (2010). Selective review of cognitive aging. *Journal of the International*  
684 *Neuropsychological Society*, 16(5), 754–760.  
685 <https://doi.org/10.1017/S1355617710000706>

686 Scarmeas, N., & Stern, Y. (2003). Cognitive Reserve and Lifestyle NIH Public Access. In *J*  
687 *Clin Exp Neuropsychol* (Vol. 25, Issue 5).

688 Schellenberg, E. G., & Weiss, W. M. (2013). 12 - Music and Cognitive Abilities. In *The*  
689 *Psychology of Music (Third Edition)* (3rd ed., Issue September 2016, pp. 499–550).  
690 Elsevier. <https://doi.org/10.1016/B978-0-12-381460-9.00012-2>

691 Schlaug, G. (2015). Musicians and music making as a model for the study of brain plasticity.  
692 *Progress in Brain Research*, 217, 37–55. <https://doi.org/10.1016/bs.pbr.2014.11.020>

693 Schneider, C. E., Hunter, E. G., & Bardach, S. H. (2019). Potential Cognitive Benefits From  
694 Playing Music Among Cognitively Intact Older Adults: A Scoping Review. *Journal of*  
695 *Applied Gerontology*, 38(12), 1763–1783. <https://doi.org/10.1177/0733464817751198>

696 Seinfeld, S., Figueroa, H., Ortiz-Gil, J., & Sanchez-Vives, M. V. (2013). Effects of music  
697 learning and piano practice on cognitive function, mood and quality of life in older  
698 adults. *Frontiers in Psychology*, 4(NOV). <https://doi.org/10.3389/fpsyg.2013.00810>

699 Signoret,J.(1991). Batterie d’Efficienc e Mnésique. Paris:Elsevier.

700 Sprague, B. N., Freed, S. A., Webb, C. E., Phillips, C. B., Hyun, J., & Ross, L. A. (2019). The  
701 impact of behavioral interventions on cognitive function in healthy older adults: A  
702 systematic review. *Ageing Research Reviews*, 52(July 2018), 32–52.  
703 <https://doi.org/10.1016/j.arr.2019.04.002>

704 Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, 47(10), 2015–2028.  
705 <https://doi.org/10.1016/j.neuropsychologia.2009.03.004>

706 Stern, Y., Arenaza-Urquijo, E. M., Bartrés-Faz, D., Belleville, S., Cantilon, M., Chetelat, G.,  
707 Ewers, M., Franzmeier, N., Kempermann, G., Kremen, W. S., Okonkwo, O., Scarmeas,  
708 N., Soldan, A., Udeh-Momoh, C., Valenzuela, M., Vemuri, P., Vuoksimaa, E., Arenaza  
709 Urquijo, E. M., Bartrés-Faz, D., ... Vuoksimaa, E. (2018). Whitepaper: Defining and  
710 investigating cognitive reserve, brain reserve, and brain maintenance. *Alzheimer’s &*  
711 *Dementia*. <https://doi.org/10.1016/j.jalz.2018.07.219>

- Sutcliffe, R., Du, K., & Ruffman, T. (2020). Music Making and Neuropsychological Aging: A Review. *Neuroscience & Biobehavioral Reviews*.  
<https://doi.org/10.1016/j.neubiorev.2020.03.026>
- Talamini, F., Altoè, G., Carretti, B., & Grassi, M. (2017). Musicians have better memory than nonmusicians: A meta-analysis. *PLoS ONE*, 12(10), 1–21.  
<https://doi.org/10.1371/journal.pone.0186773>
- Verghese, J., LeValley, A., Derby, C., Kuslansky, G., Katz, M., Hall, C., Buschke, H., & Lipton, R. B. (2006). Leisure activities and the risk of amnesic mild cognitive impairment in the elderly. *Neurology*, 66(6), 821–827.  
<https://doi.org/10.1212/01.wnl.0000202520.68987.48>
- Verghese, J., Lipton, R. B., Katz, M. J., Hall, C. B., Derby, C. A., Kuslansky, G., Ambrose, A. F., Sliwinski, M., & Buschke, H. (2003). Leisure Activities and the Risk of Dementia in the Elderly. *New England Journal of Medicine*, 348(25), 2508–2516.  
<https://doi.org/10.1056/NEJMoa022252>
- Wan, C. Y., & Schlaug, G. (2010). Music Making as a Tool for Promoting Brain Plasticity across the Life Span. *The Neuroscientist*, 16(5), 566–577.  
<https://doi.org/10.1177/1073858410377805>
- Warne, R. T. (2014). A primer on multivariate analysis of variance (MANOVA) for behavioral scientists. *Practical Assessment, Research and Evaluation*, 19(17), 1–10.
- Wechsler, D. (2000). Echelle d'intelligence de Wechsler pour adultes, troisième édition. *Manuel*. Paris: Centre de Psychologie Appliquée.
- Zendel BR, West GL, Belleville S, Peretz I. Musical training improves the ability to understand speech-in-noise in older adults. *Neurobiol Aging*. 2019;81:102-115.  
[doi:10.1016/j.neurobiolaging.2019.05.015](https://doi.org/10.1016/j.neurobiolaging.2019.05.015)



