



Faba bean root exudates alter pea root colonization by the oomycete *Aphanomyces euteiches* at early stages of infection

Yohana Laloum, Christophe Gangneux, Bruno Gügi, Arnaud Lanoue, Thibaut Munsch, Adrien Blum, Adrien Gauthier, Isabelle Trinsoutrot-Gattin, Isabelle Boulogne, Maïté Vicré, et al.

► To cite this version:

Yohana Laloum, Christophe Gangneux, Bruno Gügi, Arnaud Lanoue, Thibaut Munsch, et al.. Faba bean root exudates alter pea root colonization by the oomycete *Aphanomyces euteiches* at early stages of infection. *Plant Science*, 2021, 312, pp.111032. 10.1016/j.plantsci.2021.111032 . hal-03352845

HAL Id: hal-03352845

<https://normandie-univ.hal.science/hal-03352845>

Submitted on 16 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **ORIGINAL RESEARCH ARTICLE**

2 **Title**

3 **Faba bean root exudates ~~prevent~~ alter pea root colonization by the oomycete**
4 ***Aphanomyces euteiches* at early stages of infection**

5
6 Yohana Laloum^{1,2,#}, Christophe Gangneux¹, Bruno Gügi², Arnaud Lanoue³, Thibaut Munsch³,
7 Adrien Blum¹, Adrien Gauthier¹, Isabelle Trinsoutrot-Gattin¹, Isabelle Boulogne², Maïté
8 Vicré², Azeddine Driouich², Karine Laval¹ and Marie-Laure Follet-Gueye^{2*}.

9
10 ¹AGHYLE research unit, UP 2018.C101, UniLaSalle Rouen 3 rue du tronquet CS 40118,
11 76134 Mont Saint Aignan, France.

12 ² Normandie Univ, UNIROUEN, Glyco-MEV, EA4358, SFR NORVEGE FED 4277, I2C
13 Carnot, IRIB, 76000 Rouen, France

14 ³Université de Tours, EA 2106 «Biomolécules et Biotechnologies Végétales», UFR des
15 Sciences Pharmaceutiques, 31 Av. Monge, F37200 Tours, France

16 Present address :

17 [#]CTIFL, Centre opérationnel de Lanxade, 28 Route des Nébouts F-24130 Prignonrieux, France

18 **Running title: Control of *A. euteiches* pea root rot agent**

19 *Corresponding authors:

20 Marie-Laure Follet-Gueye

21 marie-laure.follet-gueye@univ-rouen.fr

22 ORCID : 0000-0003-3199-6659

23 Tel: + 33(0)2 35 14 00 39

24

Faba bean root exudates ~~prevent~~ alter pea root colonization by the oomycete *Aphanomyces euteiches* at early stages of infection

Abstract: *Aphanomyces euteiches* is an oomycete pathogen that causes the pea root rot. We investigated the potential role of early belowground defense in pea (susceptible plant) and faba bean (tolerant plant) at three days after inoculation. Pea and faba bean were inoculated with *A. euteiches* zoospores. Root colonization was examined. Root exudates from pea and faba bean were harvested and their impact on *A. euteiches* development were assessed by using *in vitro* assays. *A. euteiches* root colonization and the influence of the oomycete inoculation on specialized metabolites patterns and arabinogalactan protein (AGP) concentration of root exudates were also determined. In faba bean root, *A. euteiches* colonization was very low as compared with that of pea. Whereas infected pea root exudates have a positive chemotaxis index (CI) on zoospores, faba bean exudate CI was negative suggesting a repellent effect. ~~A protective effect of faba bean on pea was suggested by co-culture experiments.~~ While furanoacetylenic compounds were only detected in faba bean exudates, AGP concentration was specifically increased in pea. This work showed that early in the course of infection, host susceptibility to *A. euteiches* is involved *via* a plant-species specific root exudation opening new perspectives in pea root rot disease management.

Key words: *Aphanomyces euteiches*, *Pisum sativum*, specialized metabolites; root colonization, root exudates, *Vicia faba*

Abbreviations: AGP: arabinogalactan protein; *A. euteiches*: *Aphanomyces euteiches*; CI: chemotaxis index; dpi: days post-inoculation; RET: root extracellular trap.

1. Introduction

Aphanomyces euteiches Drechsler is a soil-borne oomycete pathogen that causes root rot disease in several legume species including pea and faba bean [1–4]. The disease is a major limitation to pea crop production worldwide [5]. Except prophylactic procedures that are mainly based on crop rotation and biodetection of potential inoculum sources in the soil, no efficient methods are currently available to control the pea root rot disease [6–9].

A. euteiches is a diploid, homothallic oomycete, which belongs to Telonemids, Stramenopiles, Alveolates, and Rhizaria (TSAR) clade [10]. Its life cycle includes both sexual and asexual reproduction stages by the succession of oospore and zoospore formation. Oospores can survive in the soil for more than ten years without host crops [11–13]. Under favorable conditions and at close vicinity of a host plant, oospores produce a mycelium that can directly infect root tissues or form a zoosporangium [14]. The zoosporangium releases a large number of zoospores [15] that are chemo-attracted by root exudates [16–18]. At the root surface, the zoospores pass from a mobile state to an immobile state. The zoospores encyst preferentially at the elongation zone within 30 min after reaching the root [19–21]. The cysts then germinate and form coenocytic hyphae that penetrate the intercellular spaces of root cortex within hours [22]. Within a few days of infection, *A. euteiches* mycelium produces oogonia, which are fertilized by antheridia resulting in the formation of new oospores [6,23]. Infected roots become soft, water-soaked and turn blackish-brown. In severe cases, seed production is reduced and plants die prematurely [5].

Although several resistance genetic studies have been conducted, no resistant pea cultivars have been identified so far [24–26]. According to Moussart *et al.*, (2008) [27] , partial levels of resistance have been reported for common vetch, faba bean, clover and other

1 species with a very high level of resistance observed in chickpea. Knowledge on the defense
2 mechanisms established by legumes in response to *A. euteiches* infection are essentially based
3 on transcriptional data obtained from the model plant *Medicago truncatula* [28–31]. Using
4 RNA-sequence analysis, Gaulin *et al.*, (2008) [32] have shown that *A. euteiches* pathogenicity
5 is correlated with the production of a specialized secretome including carbohydrate-active
6 enzymes.

7 Comprehension of *A. euteiches* life cycle during the first stage of infection could have
8 an important role in limiting root infection and enhancing plant resistance. At early stages of
9 infection, pea root tip remains free of colonization [21]. This has been attributed to the
10 presence of highly secretory cells at the root apex called border cells [33–36] or root-
11 associated, cap-derived cells (AC-DCs) [37]. These cells are the most mature cells of the root
12 cap, which detach when the root apex is placed in water [34]. At the soil-root interface, AC-
13 DCs play a key role in plant-microorganisms interactions in the rhizosphere and ensure root
14 apical meristem protection in response to biotic and abiotic stresses. [21] Cannesan *et al.*
15 (2011) demonstrated that *A. euteiches* infection induced an increase of pea root AC-DCs
16 production and altered their morphology. Root AC-DCs secrete large amounts of mucilage
17 containing polysaccharides and proteins, combined with extracellular DNA and other
18 metabolites [21,34,38–42]. The mucilage, together with root AC-DCs, form a protective web-
19 like structure named the “Root Extracellular Trap” (RET) [37,43]. It is commonly accepted
20 that over half of the root exudates originate from the RET namely by the secretion of AC-DCs
21 [44,45]. Root exudates are defined as the suite of chemical compounds released by diffusion
22 or secretion from the whole plant roots into the rhizosphere [46]. They contain both low- and
23 high molecular weight compounds. The low-molecular weight categories include amino or
24 organic acids, sugars and specialized metabolites, while the high-molecular weight
25 compounds consist mostly of polysaccharides and proteins [47] . The production of root

exudates is constitutive [48]; however under the influence of biotic and abiotic factors [49] their composition varies qualitatively and quantitatively [50] driving interactions with soil communities [51].

A few studies relating to both the impact of legume root exudates on *A. euteiches* behavior and the effect of *A. euteiches* infection on root exudate composition have been carried out [21,52–54]. The flavonoid, prunetin (5,4'-dihydroxy-7-methoxy-isoflavone) from pea root exudate is involved in *A. euteiches* zoospore attraction [52]. Pea root exudates contain specific antimicrobial compounds such as the phytoalexin pisatin whose level increases in response to *A. euteiches* [21,53]. The presence of the arabinogalactan proteins (AGPs) in the RET of pea attracts zoospores, induce their encystment and prevent the cyst germination [42].

In this study, early stages of root infection by *A. euteiches* were explored in pea (*Pisum sativum* L.) and faba bean (*Vicia faba* L.). Pea is the most susceptible host [55] whereas french faba bean was described as less sensitive and even tolerant [27]. Given that first injuries of root rot appear in the root between 3 and 4 days after inoculation [11,55], root colonization was examined at 3 days post inoculation. Pea and faba bean root exudates were harvested to investigate the early belowground responses. The effect of root exudates on *A. euteiches* development was assessed. Influence of the oomycete inoculation on specialized metabolite and glyco-polymer contents of root exudates were also determined. Our data show that faba bean root exudates may have a repellent effect on *A. euteiches* zoospores. ~~Moreover, faba bean root exudates present a protective effect on pea root against *A. euteiches* during the first three days of infection.~~ This study highlights promising future applications in pea crop management.

2. Materials and methods

2.1. Plant material

Pea (*Pisum sativum*, AVENGER variety) and faba bean (*Vicia faba*, FANFARE variety) seeds were surface sterilized with 70% ethanol (v/v) and 0.9% sodium hypochlorite solution (v/v) for 10 min. Seeds were immersed in sterile water for 6 hours at 24°C and placed in a sterile culture box (Eco box 2, Dutscher) with 1.2% agar at 24°C and 16 hours photoperiod. Pea and faba bean seedlings with 3 cm root length were used for optical microscopy analyses and infection assays.

2.2. *Aphanomyces euteiches* strain culture

Aphanomyces euteiches isolate RB84 (the French reference strain for pea) was supplied by Anne Moussart (INRAE). Isolate was grown and maintained on Potato Dextrose Agar (PDA) plates in the dark at 24°C. Zoospores were produced according to [42] Cannesan *et al.* (2012) and adjusted at a concentration of 10^5 zoospores ml⁻¹ water.

2.3. Histochemical staining and light microscopy

Pea and faba bean roots were inoculated with 10^5 zoospores ml⁻¹ during two and half hours in dark at 24°C. Seedlings were grown with sterile source water at 24°C and 16 hours photoperiod. Seedlings were harvested 1, 2 and 3 days post-inoculation (dpi), fixed in 100 % methanol and conserved at 4°C. *A. euteiches* was labeled with a fluorescein isothiocyanate–wheat germ agglutinin (FITC–WGA) conjugate, according to [21] Cannesan *et al.* (2011). Infected roots were observed using a Leica DMI 6000B inverted microscope using epifluorescence mode (excitation filter: 480 nm; emission filter: 527 nm).

2.4. *Aphanomyces euteiches* DNA quantification in root infection bioassay

Inoculated pea and faba bean roots were harvested at 1, 2 and 3 dpi for total DNA extraction. Non-inoculated seedlings were used as controls. Three independent biological replicates were

1 used. Total genomic DNA was extracted from roots using the PowerPlant ® Pro DNA
2 isolation kit (MoBio Laboratories, USA) according to the manufacturer's instructions, and
3 eluted in 50 µl and stored at -20°C. DNA was quantified by fluorimetry using the Fluorescent
4 DNA quantitation Kit (Hoechst 33258, Bio-Rad, CA). Total genomic DNA of *A. euteiches*
5 was extracted from mycelium. *A. euteiches* isolate was grown on 60 ml of Sabouraud
6 Dextrose broth 30 % (w/v) in a cell culture flask during 7 days at 24°C in the dark. Total
7 genomic DNA was extracted from 50 mg crushed lyophilized mycelium using the E.N.Z.A ®
8 HP Fungal DNA Kit (Omega bio-tek, USA). Total DNA was eluted in 50 µl and was
9 quantified using the Eppendorf BioPhotometer. DNA extracts were stored at -20°C. Real-time
10 quantitative PCR was performed using the LightCycler® 480 Instrument II Real Time PCR
11 system (Basel, Switzerland, Roche) in a total volume of 25 µl. The qPCR mix was prepared as
12 follows: 4 ng of plant DNA, 12.5 µl of Thermo Scientific ABsoluteqPCR Mix no rox
13 (Thermo Fischer Scientific), 0.25 mM bovin serum albumin (GeneOn, DE) and 250 nM of
14 each primer Ae_ITS1_39F and Ae_ITS1_167R targeting the ITS region, described by [8]
15 Gangneux *et al.* (2014). Influence of plant DNA and component co-extracted from roots was
16 previously assessed. A first range with serial dilution from 4 ng to $4 \cdot 10^{-5}$ ng of *A. euteiches*
17 DNA was performed in triplicate as control. A second range was achieved in triplicate with
18 serial dilution from 4 ng to $4 \cdot 10^{-5}$ ng of *A. euteiches* DNA, add to 10 ng of plant DNA
19 extracted from pea or faba bean roots. Amplification results were compared between the two
20 ranges. After an initial enzyme activation step of 15 min at 95°C, 50 cycles of PCR were
21 performed as follows: 30 s at 95°C, 30 s at 60°C, and 30 s at 72°C. In the last step, melting
22 curves were added with the following conditions: 10 s at 95°C, 30 s at 60°C and slowly
23 heating to 97°C, with 1.1°C increase every 10 seconds with continuous measurement of
24 fluorescence at 520 nm.

2.5. Root exudate collection from seedling roots

Seedlings with a root length of 3 cm were placed in an ecobox within 200 ml of source water. One hundred seedlings were used per condition. Infection assays were carried out by inoculating roots with a solution of 10^5 zoospores ml^{-1} during two and half hours in the dark at 24°C. Control conditions were done by maintaining seedlings two and half hours in water. Roots were then rinsed and placed in 200 ml of water. Both control and inoculated seedlings were incubated at 24°C and 16 hours photoperiod during 3 days. Root exudates from pea and faba bean were performed by adapting the method previously described by [56] Carreras *et al.* (2019). Root incubation medium was centrifugated at 9 000 rpm for 15 min. Root exudates (supernatant) were freeze dried and weighed for subsequent biological assay and biochemical analyses.

2.6. Chemotaxis bioassay on *Aphanomyces euteiches* zoospores

Chemotaxis tests were performed by adapting the method previously described by [42] Cannesan *et al.* (2012). Zoospore attraction and germination were explored on a microscopy wells-slide (Fig. 3A). A droplet of zoospores containing 500 zoospores (20 μl) was added in the middle well, source water (20 μl) was added into the right well and test compounds into the left well (20 μl). Water was used as control and Gum Arabic (GA) (0.02 mg) as standards, whereas pea or faba bean root exudates from inoculated or non-inoculated seedlings are tested (4 mg). A bridge was made between the wells. The three wells were connected to each other by water. Only the zoospores, which moved out of the middle well were counted after 4 hrs. A chemotaxis index (CI) was calculated using equation based on previous studies [57–60] as:
$$\text{CI} = \frac{(\text{Number of zoospores within the Z1} - \text{Number of zoospores within the Z2})}{(\text{Number of zoospores which made a choice Z1+Z2})}$$
 Slides were incubated 4 hours in the dark at room

temperature. The percentage of germinated encysted zoospores was calculated as: [(Number of germinated cysts per zone / Total number of attracted cysts per zone) X100]. Observations were made with a Leica DM IL with a 10x objective.

2.7. *Aphanomyces euteiches* mycelial growth bioassay

Pea and faba bean root exudates were evaluated for their effect on *A. euteiches* mycelium growth. A mycelium plug was cut from the margin of 7 days-old on PDA, deposited on the Petri plate center containing PDA medium and 50 µg ml⁻¹ of ampicillin and 12.5 µg ml⁻¹ of streptomycin. A droplet of 50 µl containing 4 mg of pea or faba bean exudates was deposited at 1.5 cm from the mycelium plug. Water was used as control. Petri plate was placed at 24°C in the dark. Mycelium growth was measured at 3, 4, 5, 6 and 7 days post-incubation.

2.8. UPLC-MS analyses

Freeze dried root exudates were resuspended in methanol before UPLC-MS analyses. The system consisted in an ACQUITY™ Ultra Performance Liquid Chromatography system coupled to a photo diode array detector (PDA) and a Xevo TQD mass spectrometer (Waters, Milford, MA) equipped with an electrospray ionization (ESI) source controlled by Masslynx 4.1 software (Waters, Milford, MA). Analyte separation was achieved by using a Waters Acquity HSS T3 C18 column (150 × 2.1 mm, 1.8 µm) with a flow rate of 0.4 ml min⁻¹ at 55°C. The injection volume was 5 µl. The mobile phase consisted of solvent A (0.1% formic acid in water) and solvent B (0.1% formic acid in acetonitrile). Chromatographic separation was achieved using an 18-min linear gradient from 5 to 70% (v/v) solvent B. MS detection was performed in both positive and negative modes. The capillary voltage was 3,000 V and sample cone voltages were 30 and 50 V. The cone and desolvation gas flow rates were 60 and 800 l h⁻¹. A list of mass-to-charge (m/z) ratios corresponding to molecular ions was

established for both plant species, based on total ion current chromatograms from switching positive (ES^+) and negative modes (ES^-). Identification of analytes was based on retention times, m/z values, and UV spectra and by comparison with commercial standards or data from literature when no authentic standards were available (Supplementary Table S1 and S2). UPLC-MS analyses were achieved using selected ion monitoring (SIM) mode of the targeted molecular ions. SIM chromatograms were integrated using the subroutine QuanLynx 4.1 for data mining. Peak integration was performed using the ApexTrack algorithm with a mass window of 0.1 Da and relative retention time window of 1min followed by Savitzky–Golay smoothing (iteration = 1 and width = 1). The resulting pairs of m/z values and retention times were also manually controlled. Tryptophan, *p*-coumaric acid, *t*-ferulic acid and formononetin were purchased from Sigma Chemical Company (St. Louis, MO) and pisatin from Apin Chemicals Ltd (Abingdon,UK).

2.9.Preparation of alcohol insoluble residues (AIR) and monosaccharide analysis

Root exudates was incubated with 96% ethanol (4:1) 24 hours at 4°C. Samples were centrifuged at 4 000 g during 15 min at 4°C and precipitated molecules (AIR of root exudates) was dried and stored at -20°C. Two mg of AIR of root exudates from pea and faba bean were treated in 2 M trifluoroacetic acid (TFA) for 2 hours at 110°C to release monosaccharides. TFA from samples was removed twice with 50% isopropanol: water solution washings. After freeze-drying, monosaccharide hydrolysates were converted in methyl-glycosides in 1 M HCl/methanol (Supelco) for 16 hours at 80°C. Samples were washed twice with methanol and were treated with hexamethyldisilazane (HMDS): trimethylchlorosilane (TMCS): pyridine solution (3: 1: 9) (Supelco), for 20 min at 80°C. The resulting trimethylsilyl methyl glycosides were dried, resuspended in 1 ml of cyclohexane and injected in the 3800 Varian GC-FID system equipped with a CP-Sil5-CB capillary column (Agilent Technologies) and Helium was

the gas vector. Elution was performed with the following method: 40°C for 3 min, 15°C min⁻¹ from 40°C to 160°C, 1.5°C min⁻¹ from 160°C to 220°C and finally 20°C min⁻¹ from 220°C to 280°C. Relative quantification of each monosaccharide was carried out using external monosaccharide standards for their respective retention times and response factors and using inositol as internal standard [61].

2.10. *Arabinogalactan protein (AGP) quantification by rocket gel electrophoresis*

Yariv reagent (β -d-Glc Y) [62–64] was used to detect free or attached type II arabinogalactan (AGII) to AGPs in root exudates using rocket electrophoresis. AGPs were quantified in root exudates precipitated (AIR of exudates) or not by ethanol. For the two plant species, the analyses were applied on extracts from seedlings inoculated or not by *A. euteiches*. Gels were composed of 1% agarose containing 90 mM Tris (pH 8.4), 90 mM boric acid, 2 mM EDTA, and 20 mg.ml⁻¹ of β -d-Glc Y reagent according to van Holst and Clarke (1986). Root exudate fractions (1 mg) were loaded into each well and run for 16 h at 200 V/5 mA/10 W. Gum Arabic from acacia (Fisher Scientific) was used as a standard. 1 mg of *A. euteiches* mycelium and 1 mg of mycelium exudates were also loaded as control. After migration, gels were rinsed with 2% NaCl (w/v). Quantification of AGP was estimated using peak areas of Gum Arabic as standard. Peak areas of AGII were from the sample fractions and Gum Arabic control were calculated using ImageJ software.

2.11. *Statistical analysis*

GraphPad Prims 7.0 software was used for statistical analysis. Nonparametric tests (Kruskal-Wallis and Mann-Withney) were performed with 5% level of significance.

3. Results

3.1. Root colonization by *A. euteiches* during the first three days of infection

Pea root surface colonization was assessed by *A. euteiches* staining at 1, 2 and 3 dpi (Fig. 1). The colonization of pea roots was much stronger than that of faba bean roots. At 1 dpi, encysted zoospores and germinated cysts were localized over both the elongation and maturation zones (Fig1B). Mycelial hyphae were mostly observed longitudinally at the maturation zone. At 2 dpi, mycelium was extended and grown on the root elongation and maturation zones. At 3 dpi, it fully surrounded the root tip and elongation zones (Fig.1B). On faba bean root surface, at 1 dpi, a lower number of cysts was observed. Cysts were distributed throughout the root and a few cysts germinated preferentially above the meristematic zone. At 2 dpi, cyst germination was weak. At 3 dpi, a few cysts had produced a germ tube on both elongation and maturation zone root parts.

3.2. Effect of pea and faba bean co-cultures on the colonization of the roots by the pathogen

We monitored the colonization of roots of both species by the pathogen in a co-culture by quantifying *A. euteiches* DNA from roots of both species at 1, 2, and 3 dpi (Fig. 2 and Fig. S2). We also assessed the effect of faba bean root exudates on the colonization of pea root by the pathogen. Three experimental conditions were used. To test the impact of co-culture on zoospore encystment, both species were co-inoculated with zoospores and co-cultivated during 3 days (condition 1). In condition 2, the plants were inoculated separately and then co-cultivated in order to evaluate the effect of co-culture on mycelium growth. Pea plants were inoculated and then cultured in the presence of faba bean root exudates in condition 3. Controls corresponding to peas and faba beans inoculated and cultured separately have been made (Supplementary Fig. S1). Major data are first that there is more pathogen DNA in pea

than in faba bean roots (Fig. 2A, 2B and Fig. S1) and secondly that *A. euteiches* DNA content decreased in both plants when they are co-cultivated (Fig. 2A, 2B). While DNA content of *A. euteiches* is strictly reduced in faba bean at 1 dpi, it is significantly altered in pea at 3 dpi. Third, the addition of faba bean root exudates in the pea culture medium induced a significant reduction of infection (measured by the content of the pathogen DNA) (Fig. 2C). For pea, there are no significant difference between the three conditions (Fig. S2). The comparison of the number of ITS copies between *A. euteiches* DNA control and *A. euteiches* DNA supplemented by plant DNA did not reveal any significant differences (Supplementary Table S1).

3.3. Effect of pea and faba bean root exudates on *A. euteiches*

To study the effect of root exudates on zoospore behavior, we have performed chemotaxis and germination *in vitro* assays using the experimental device shown in Fig 3A. We have assessed the effect of bean and pea root exudates as well as Gum Arabic (Fig.3B, 3C). Our data show that root exudates from non-inoculated peas had no chemoattractive effect on zoospores whereas those from inoculated plants presented a significant positive CI of 0.31 ± 0.03 . Interestingly, we found that root exudates from both non-inoculated and inoculated faba bean plants had a significant negative CI equal to -0.52 ± 0.12 and -0.49 ± 0.06 (Fig. 3B). Remarkably, among the root exudates tested, only those from non-inoculated peas promoted the germination of cysts ($13.64 \pm 2.84\%$) as compared with the controls $1.67 \pm 1.11\%$ (Fig. 3C). Gum Arabic, had a strong attractive effect (CI = 0.82 ± 0.14) but did not promote cyst germination. Chemotaxis index corresponding to the neutral condition (water vs water) was 0.17 ± 0.19 and the percentage of germinated cysts in Z1 and Z2 areas were $7.25 \pm 4.57\%$ and $5.02 \pm 3.15\%$ (Fig. 3C).

The impact of root exudates from non-inoculated plants on *A. euteiches* mycelium growth was determined (Fig. 3D). Except a slight positive effect of pea root exudates on mycelium growth at days 4 and 5, the growth of *A. euteiches* mycelium was neither inhibited nor stimulated by the exudates of both plants. Also, no alteration of mycelium growth was observed in the presence of antibiotics in the PDA medium (Supplementary Table S2).

3.4. UPLC-MS based analyses of root exudates

UPLC-DAD-MS-based metabolite profiling was carried out on pea and faba bean root exudates. Major peaks were annotated and assigned according to their retention time, UV and mass spectra by comparison with standards or data from the literature. MS experiments were performed in both positive and negative ionization modes. For the two species, while the metabolic profiles were not qualitatively altered by infection, the concentration of metabolites often increased (tables 1 and 2; Fig. S3).

In pea root exudate extracts, five peaks were assigned to tryptophan, *p*-coumaric acid, trans-ferulic acid, formononetin and pisatin by comparison with authentic standards (Table 1 and Fig. S3A). All these metabolites have been previously described in pea [65,66]. At a retention time of 5.38 min, a peak showed a molecular ion [M-H]⁻ at m/z 609 and [M+H]⁺ at m/z 611. In negative mode this analysis produced a characteristic fragment at m/z 301 [M-H-glucose-rhamnose]⁻ corresponding to a quercetin moiety. Therefore, this compound was identified as quercetin-3-rutinoside as previously described in pea seed coat extracts [67]. A peak at 7.10 min produced a molecular ion [M-H]⁻ at m/z 302. The corresponding UV spectrum and MS fragmentation features allowed to annotate this peak as DMDI (7,2'-dihydroxy-4',5'-methylene-dioxyisoflavanol). In the same way, the peak at 10.22 min showing molecular ions [M-H]⁻ at m/z 297 and [M+H]⁺ at m/z 299 as well as specific UV spectrum was assigned as

DMD (7,2'-dihydroxy-4',5'-methylenedioxyisoflavone). These two last compounds are known to be biosynthetic intermediates in the pathway of pisatin a phytoalexin of pterocarpan type with formononetin as precursor [66]. In faba bean exudates, tryptophan and formononetin were undoubtedly identified by comparison with pure standards (Table 2 and Fig. S3B). The peak at retention time 6.32 min, showing a molecular ion [M-H]⁻ at m/z 385 and a fragment ion 223 [M-H-162]⁻ corresponding to the loss of a glucose moiety was assigned to sinapoyl-glucoside, a hydroxycinnamic acid derivative previously described in faba bean leaf extracts (Neugart *et al.*, 2015). Peak at 8.11 min produced a [M-H]⁻ molecular ion at m/z 273 and fragments ions at m/z 224 and 183. It was assigned to wyerone epoxide as reported in faba bean (Hargreaves *et al.* 1976). Peak at 10.41 min gave a molecular ion [M-H]⁻ at m/z 261 producing two fragments at m/z 217 and 191, and was assigned to dihydrowyerol. This metabolite was previously reported in *Vicia faba* [68]. Peak at 10.57 min showed a [M-H]⁻ ion at m/z 243 and a fragment ion at m/z 225 and was attributed to wyerone acid as previously described [68]. These three last compounds belong to furanoacetylene family, a non-flavonoid class of phytoalexins. The peak at 12.09 min showed a molecular ion [M-H]⁻ at m/z 269 and was assigned to the pterocarpan medicarpin as previously described in broad bean [69].

3.5. AGPs content in root exudates

Given that AGPs impacted *A. euteiches* development [42] and that the relative levels of Ara and Gal monosaccharides were modulated following inoculation (Fig. S4; Table S3), we focused on AGPs analysis. A semi-quantitative analysis of type II arabinogalactans (AG-II), a typical side chain of AGPs, was performed (Table 3). Rocket gel electrophoresis of root exudates (AIR or raw fractions) detected AGP in pea and faba bean extracts (Supplementary Fig. S5). The amount of AGP strongly increased in extracts from infected pea roots as compared with non-infected roots (Table 3). In contrast, no significant variations were

observed in extracts from faba bean roots. It is important to note that AGPs, were detected neither in *A. euteiches* mycelium nor in the mycelium culture medium (Supplementary Fig. S5).

4. Discussion

Although the life cycle of *A. euteiches* is characterized by rapid colonization of the root system within hosts, better understanding of the early stages of the disease development is necessary. We studied the *A. euteiches* colonization, during the first three days of infection, within pea and faba bean roots respectively sensitive and tolerant to this root rot agent. ~~To the best of our knowledge, this work reveals for the first time that pea root colonization by *A. euteiches* can be significantly reduced by faba bean root exudates.~~

4.1. Host specificity of *A. euteiches* infection at early stages

Compared with faba bean, pea roots are more intensely colonized by *A. euteiches* during the first three days of infection. Microscopical and qPCR data showed that colonization of pea root is high at 3 dpi. These results are in line with those reported by Billard et al. (2019) [54] showing that root necrosis is observed much earlier in pea than in faba bean roots. We show that host sensitivity is involved in the oomycete infectious process at early stages and attest that *A. euteiches* strain RB84, is preferentially aggressive on the susceptible plant. As previously observed [19,21], we show that pea root tips remained free of infection. These results highlight the protective function of root cap-derived border cells and their secretions toward the root apical meristem at early stages of infection. The primary function of motile zoospores is to connect oomycetes to the host [70]. Several host-specific molecular attractants have been reported such as isovaleraldehyde, which attracts *Phytophthora palmivora* [71], prunetine a pea compound that attracts *A. euteiches* zoospores [52] or indole-3-aldehyde

1 isolated from cabbage that promotes *Aphanomyces raphani* zoospores attraction [72]. Root
2 exudates from pea induce a higher rate of oospore germination than those from vetch and faba
3 bean [54]. Although, pea root exudates from non-inoculated plants affect zoospore
4 germination it has little effect on zoospore attraction. The chemotaxis index (CI) of exudates
5 originated from non-inoculated peas is nearly null. In contrast, the CI is positive for exudates
6 originated from inoculated peas. Kong and Hong (2010) [73] reported that *Phytophthora*
7 *nicotianae* zoospores secrete a fluid able to promote successful infection in *Arabidopsis*
8 *thaliana*. Exudates from zoospores of *Phytophthora* and *Pythium* species act as a *quorum-*
9 *sensing* signals impacting zoospore homing and triggering infection [73–75]. Herein, exudates
10 from *A. euteiches* zoospores and/or mycelium may be present in harvested pea exudates and
11 would promote zoospore attraction. The CI for root exudates from non-inoculated or
12 inoculated faba beans was negative which suggests that faba bean exudates had a repellent
13 effect towards *A. euteiches* zoospores. Host repellent signals have been mostly demonstrated
14 in plant-nematode interactions [76–79]. As for oomycete-host interactions, a phenolic extract
15 from the root of the non-host plant *Portulaca oleracea*, was shown to exhibit a repellent
16 activity on *Aphanomyces cochlioides* zoospores [80]. Bazghaleh *et al.*, (2018) [81] showed
17 that, in response to *A. euteiches* infection, the root polyphenolic composition of lentil is
18 dependent on the plant genotypes. The metabolic profiling of pea and faba bean root exudates
19 based on UPLC-MS analyses revealed the presence of two main classes of plant specialized
20 metabolites, phenolic compounds and furanoacetylenic derivatives. Among phenolic
21 compounds, flavonoids are ubiquitously found in the plant kingdom, whereas isoflavones are
22 restricted to the subfamily of Papilionoideae [82]. Within Fabaceae, almost all species of the
23 subfamily Papilionoideae accumulate isoflavones and derivatives, including phytoalexins of
24 the pterocarpan type [82]. Here, the presence of pterocarpan biosynthetic pathway was
25 observed in both legumes confirming previous chemotaxonomy observations in whole plants

[66,69]. Focusing on the root exudate composition, the presence of the phytoalexin pisatin in pea root exudates was already reported [21], however the presence of pterocarpan-type phytoalexin (medicarpin) in faba bean root exudates has never been reported in earlier studies. Although different susceptibility levels toward *A. euteiches* were observed, pterocarpan-type phytoalexins were identified in both legumes suggesting that they might not be strongly involved in the susceptibility/tolerance during early stages of infection by *A. euteiches*. A search for the distribution of furanoacetylene in the metabolite-plant species database Knapsack indicates a presence restricted to the two genus *Vicia* and *Lens* [83]. Furanoacetylenic derivatives were described as markers of resistance of faba bean to chocolate spot disease [84]. The ability of *Botrytis fabae* and not *B. cinerea* to metabolize the phytoalexin wyerone acid was of primary significance in the pathogenicity towards faba bean [68]. Therefore, we suggest that the presence of furanoacetylenic in root exudates of faba bean might contribute to tolerance mechanisms of faba bean to the pathogen *A. euteiches*.

Coupled to small molecules, root exudates contain macromolecules like cell wall polymers [85,86] (Chaboud, 1983; Knee *et al.*, 2001). Hinch and Clarke (1980) [87] demonstrated that zoospore adhesion of *Phytophthora cinnamomi* depends on carbohydrate components present at the *Zea mays* L. root surface. Pea root exudate monosaccharide profiles are quite similar to those previously described [86,88,89]. For the two legumes, Ara and Gal are among the major sugars detected suggesting the presence of arabinogalactan motifs. The precipitation by the β -glucosyl Yariv reagent assessed that AG-II are present in the root exudates. AG-II are the major side chain of arabinogalactan proteins (AGP). AGP is a family of non-enzymatic cell surface hydroxyproline-rich glycoproteins found in plant cell walls, in the plasma membrane or in plant secretions [63,90] especially the root cap-associated mucilage and root exudates [42,56,91]. These proteoglycans control cell morphology and growth, and play a major role in

1 plant-micro-organism interactions [38,91–93]. Here, a high CI was recorded with the Gum
2 Arabic (an AGP-like molecule) whereas the cyst germination was not detectable. These
3 observations are in line with the results described by Cannesan *et al.* (2012) [42] using
4 purified pea root AGPs that show that AGPs stimulate zoospore attraction and encystment
5 but also reduce cyst germination. The present results revealed an increase of AGP content in
6 root exudates concomitantly to infection in pea. Exudates from non-inoculated peas were
7 favorable to cyst germination while those from inoculated seedlings had no impact. This
8 could be linked to the increase of AGP in pea root exudates after infection.

9 ~~Attraction of zoospores and their encystment do not necessarily guarantee successful~~
10 ~~infection and mycelium colonization mostly depends on their capacity to penetrate host~~
11 ~~tissues [94]. Van West *et al.* (2003) [70] suggest that host specificity is involved during the~~
12 ~~penetration phase. The estimation of *A. euteiches* root content, indicate that mycelium~~
13 ~~developed more importantly in pea root as compared with faba bean showing a better ability~~
14 ~~to pea root penetration. Expression of pea genes involved in cell wall modifications such as~~
15 ~~those encoding callose synthase, pectin esterase and pectin esterase inhibitor are modulated~~
16 ~~during *A. euteiches* infection [31]. Gaulin *et al.* (2018) [95] have also shown that *A. euteiches*~~
17 ~~pathogenicity towards *Medicago truncatula* is correlated with the production of specialized~~
18 ~~secretome including carbohydrate active enzymes such as cellulase, cellobiohydrolase,~~
19 ~~xylanase, polygalacturonase and pectin lyase. During penetration and colonization stages,~~
20 ~~*A. euteiches* is likely to secrete specific cell wall degrading enzymes that facilitate mycelium~~
21 ~~penetration and development within the host [96–98].~~

22 4.2. Pea susceptibility towards *A. euteiches* is reduced by faba bean

Moussart *et al.* (2013) [99] indicated that faba bean resistant cultivars contribute to reduce the inoculum potential of soils naturally infested by *A. euteiches*. We revealed that *A. euteiches* colonization in pea root at 3 dpi is significantly reduced when pea is co-cultured with faba bean or when pea is inoculated and cultured in the presence of faba bean root exudates. As a significant decrease of *A. euteiches* DNA content is strictly recorded at 1dpi in faba bean and at 3dpi in pea, we hypothesize that co-inoculation and co-culture experiments have distinct effect in the two plants. Co-culture have a minor impact on zoospore attraction but affect the mycelium colonization in pea. Conversely, we could postulate that co-culture prevents zoospore attraction in faba bean. As a whole, these results suggest that faba bean might have a protective effect on pea. This protective effect of faba bean could be related to the repulsive or biocide effects of its exudates, and /or to another effect such as an activation of pea root defense responses. Our data reveal that faba bean exudates have no biocide effect on mycelium and their repulsive effect is suggested by chemotaxis *in vitro* assay. In response to oligogalacturonide elicitor in pea, Sameh et al. in 2017, [100] showed an upregulation of genes involved either in the basal defense (salicylic acid and reactive oxygen species genes) and either in the antifungal defensins, lignans and the phytoalexin pisatin pathways. Expression of pea genes involved in cell wall modifications such as those encoding callose synthase, pectin esterase and pectin esterase inhibitor are also reported to be modulated during *A. euteiches* infection [31]. Supplementary experiments such as monitoring of plant defense genes or marker enzymatic activities could be performed to investigate pea defense responses to *A. euteiches* infection at early stages in presence of faba bean exudates. To assess the protective effect of faba bean towards pea it also would be interesting to explore it throughout the plant's life.

Root exudates directly influence root-root interaction in the rhizosphere [51,53,100–102]. Regarding plant species communication, Wentzell and Kliebenstein (2008) [103] showed that glucosinolate activation in *Arabidopsis* was regulated by the density of neighboring plants. More recently, Chen *et al.* (2019) [104] showed that tobacco plants respond to the presence of belowground neighbors by activating different defense pathways.

Conclusions

The present study provides new findings related to *A. euteiches* colonization at early stages of infection in pea and faba bean, respectively susceptible and tolerant hosts. We revealed that faba bean root exudates presented a negative CI towards zoospores. We showed that at three days after zoospore inoculation, *A. euteiches* was more abundant in pea compared with faba bean root. We also reveal that pea root colonization by *A. euteiches* can be significantly reduced by faba bean root exudates. These provide promising perspectives for new pest management toward *Aphanomyces* root rot in the field pea crops.

6. Author Contributions

Y.L, M-L.F-G and C.G planned and designed the research. Y.L performed experiment and wrote the first version of the manuscript with the assistance of M-L.F-G, A. G. and A.D. B.G contributed in GC-FID experiment, A.B contributed in biological assays in Petri plate, A.L and T.M performed the phenolic composition of root exudates by UPLC-MS analyses. All authors read, provided critical feedback and approved the final version of the manuscript.

Funding

This work was supported by the research network SFR NORVEGE of Normandy region in the Give Peas a Chance project, the French ministry of Research and Higher Education and

the University of Rouen Normandy. YL was supported by a PhD fellowship from the Normandy Region.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgments

The authors are thankful to Anne Moussart (INRAE) for providing the *A. euteiches* strain RB84 and Philippe Declerck from RAGT for providing pea and faba bean seeds

References

[1] F.R. Jones, C. Drechsler, Root Rot of Peas in the United States Caused by *Aphanomyces Euteiches* nsp, J. Agric. Res. 30 (1925) 293-325 Plates 1–6.

[2] W.F. Pfender, D.J. Hagedorn, *Aphanomyces euteiches* f.sp. *phaseoli*, a causal agent of bean root and hypocotyl rot., (1982). <https://www.semanticscholar.org/paper/Aphanomyces-euteiches-f.sp.-phaseoli%2C-a-causal-of-Pfender-Hagedorn/493ecf0811bf76810d72cc0d3340f2c5436d18c2> (accessed April 12, 2020).

[3] L. Lamari, Etiology of seedling blight and root rot of faba bean (*Vicia faba*) in Manitoba, (1982). <https://mspace.lib.umanitoba.ca/xmlui/handle/1993/3465> (accessed July 26, 2017).

[4] E.B. Holub, C.R. Grau, J.L. Parke, Evaluation of the *forma specialis* concept in *Aphanomyces euteiches*, Mycol. Res. 95 (1991) 147–157. [https://doi.org/10.1016/S0953-7562\(09\)81004-6](https://doi.org/10.1016/S0953-7562(09)81004-6).

[5] L. Wu, K.-F. Chang, R.L. Conner, S. Strelkov, R. Fredua-Agyeman, S.-F. Hwang, D. Feindel, *Aphanomyces euteiches*: A Threat to Canadian Field Pea Production, Engineering. 4 (2018) 542–551. <https://doi.org/10.1016/j.eng.2018.07.006>.

[6] E. Gaulin, C. Jacquet, A. Bottin, B. Dumas, Root rot disease of legumes caused by *Aphanomyces euteiches*, Mol. Plant Pathol. 8 (2007) 539–548. <https://doi.org/10.1111/j.1364-3703.2007.00413.x>.

[7] Anne Moussart, Development of an efficient screening test for pea resistance to *Aphanomyces euteiches*., (2001). <http://onlinelibrary.wiley.com/insb.bib.cnrs.fr/doi/10.1046/j.1365-3059.2001.00590.x/epdf> (accessed July 18, 2017).

- 1 [8] C. Gangneux, M.-A. Cannesan, M. Bressan, L. Castel, A. Moussart, M. Vitré-
2 Gibouin, A. Driouich, I. Trinsoutrot-Gattin, K. Laval, A Sensitive Assay for Rapid Detection
3 and Quantification of *Aphanomyces euteiches* in Soil, *Phytopathology*. 104 (2014) 1138–
4 1147. <https://doi.org/10.1094/PHYTO-09-13-0265-R>.
- 5 [9] L. Wu, K.-F. Chang, S.-F. Hwang, R. Conner, R. Fredua-Agyeman, D. Feindel, S.E.
6 Strelkov, Evaluation of host resistance and fungicide application as tools for the management
7 of root rot of field pea caused by *Aphanomyces euteiches*, *Crop J.* 7 (2019) 38–48.
8 <https://doi.org/10.1016/j.cj.2018.07.005>.
- 9 [10] P.J. Keeling, F. Burki, Progress towards the Tree of Eukaryotes, *Curr. Biol.* 29 (2019)
10 R808–R817. <https://doi.org/10.1016/j.cub.2019.07.031>.
- 11 [11] G.C. Papavizas, W.A. Ayers, *Aphanomyces* Species and Their Root Diseases in Pea
12 and Sugarbeet: A Review, Agricultural Research Service, U.S. Department of Agriculture,
13 1974.
- 14 [12] P. W.f, H. D.j, Disease progress and yield loss in *Aphanomyces* root rot of peas
15 [*Aphanomyces euteiches* f. sp. *pisii*, *Pisum sativum*]., *Phytopathology*. (1983).
16 <http://agris.fao.org/agris-search/search.do?recordID=US19840071497> (accessed April 12,
17 2020).
- 18 [13] T.L. Willsey, S. Chatterton, M. Heynen, A. Erickson, Detection of interactions
19 between the pea root rot pathogens *Aphanomyces euteiches* and *Fusarium* spp. using a
20 multiplex qPCR assay, *Plant Pathol.* 67 (2018) 1912–1923.
21 <https://doi.org/10.1111/ppa.12895>.
- 22 [14] H. Shang, C.R. Grau, R.D. Peters, Oospore Germination of *Aphanomyces euteiches* in
23 Root Exudates and on the Rhizoplanes of Crop Plants, *Plant Dis.* 84 (2000) 994–998.
24 <https://doi.org/10.1094/PDIS.2000.84.9.994>.
- 25 [15] D.K. Malvick, C.R. Grau, J.A. Percich, Characterization of *Aphanomyces euteiches*
26 strains based on pathogenicity tests and random amplified polymorphic DNA analyses,
27 *Mycol. Res.* 102 (1998) 465–475. <https://doi.org/10.1017/S0953756297005029>.
- 28 [16] J.L. CUNNINGHAM, Attraction of *Aphanomyces euteiches* zoospores to pea and
29 other plant roots., *Phytopathology*. 52 (1962) 616–618.
- 30 [17] H. Sekizaki, R. Yokosawa, C. Chinen, H. Adachi, Y. Yamane, Studies on zoospore
31 attracting activity. II. Synthesis of isoflavones and their attracting activity to *Aphanomyces*
32 *euteiches* zoospore, *Biol. Pharm. Bull.* 16 (1993) 698–701.
- 33 [18] J.W. Deacon, S.P. Donaldson, Molecular recognition in the homing responses of
34 zoosporic fungi, with special reference to *Pythium* and *Phytophthora*, *Mycol. Res.* 97 (1993)
35 1153–1171. [https://doi.org/10.1016/S0953-7562\(09\)81278-1](https://doi.org/10.1016/S0953-7562(09)81278-1).

- 1 [19] J.L. Cunningham, D.J. Hagedorn, Penetration and infection of pea roots by zoospores
2 of *Aphanomyces euteiches*, *Phytopathology*. 52 (1962) 827–834.
- 3 [20] J.W. Deacon, G. Saxena, Germination triggers of zoospore cysts of *Aphanomyces*
4 *euteiches* and *Phytophthora parasitica*, *Mycol. Res.* 102 (1998) 33–41.
5 <https://doi.org/10.1017/S0953756297004358>.
- 6 [21] M.A. Cannesan, C. Gangneux, A. Lanoue, D. Giron, K. Laval, M. Hawes, A.
7 Driouich, M. Vitré-Gibouin, Association between border cell responses and localized root
8 infection by pathogenic *Aphanomyces euteiches*, *Ann. Bot.* 108 (2011) 459–469.
9 <https://doi.org/10.1093/aob/mcr177>.
- 10 [22] I. Badreddine, C. Lafitte, L. Heux, N. Skandalis, Z. Spanou, Y. Martinez, M.-T.
11 Esquerre-Tugaye, V. Bulone, B. Dumas, A. Bottin, Cell Wall Chitosaccharides Are Essential
12 Components and Exposed Patterns of the Phytopathogenic Oomycete *Aphanomyces*
13 *euteiches*, *Eukaryot. Cell*. 7 (2008) 1980–1993. <https://doi.org/10.1128/EC.00091-08>.
- 14 [23] W.W. Scott, A monograph of the genus *Aphanomyces*, Virginia Agricultural
15 Experiment Station, Blacksburg, Va., 1961.
- 16 [24] M.L. Pilet-Nayel, F.J. Muehlbauer, R.J. McGee, J.M. Kraft, A. Baranger, C.J. Coyne,
17 Consistent Quantitative Trait Loci in Pea for Partial Resistance to *Aphanomyces euteiches*
18 Isolates from the United States and France, *Phytopathology*. 95 (2005) 1287–1293.
19 <https://doi.org/10.1094/PHTO-95-1287>.
- 20 [25] C. Lavaud, A. Lesné, C. Piriou, G.L. Roy, G. Boutet, A. Moussart, C. Poncet, R.
21 Delourme, A. Baranger, M.-L. Pilet-Nayel, Validation of QTL for resistance to *Aphanomyces*
22 *euteiches* in different pea genetic backgrounds using near-isogenic lines, *Theor. Appl. Genet.*
23 128 (2015) 2273–2288. <https://doi.org/10.1007/s00122-015-2583-0>.
- 24 [26] A. Desgroux, V. L'Anthoëne, M. Roux-Duparque, J.-P. Rivière, G. Aubert, N. Tayeh,
25 A. Moussart, P. Mangin, P. Vetel, C. Piriou, R.J. McGee, C.J. Coyne, J. Burstin, A. Baranger,
26 M. Manzanares-Dauleux, V. Bourion, M.-L. Pilet-Nayel, Genome-wide association mapping
27 of partial resistance to *Aphanomyces euteiches* in pea, *BMC Genomics*. 17 (2016) 124.
28 <https://doi.org/10.1186/s12864-016-2429-4>.
- 29 [27] A. Moussart, M.N. Even, B. Tivoli, Reaction of genotypes from several species of
30 grain and forage legumes to infection with a French pea isolate of the oomycete
31 *Aphanomyces euteiches*, *Eur. J. Plant Pathol.* 122 (2008) 321–333.
32 <https://doi.org/10.1007/s10658-008-9297-y>.
- 33 [28] F. Colditz, H.-P. Braun, C. Jacquet, K. Niehaus, F. Krajinski, Proteomic profiling
34 unravels insights into the molecular background underlying increased *Aphanomyces*
35 *euteiches*-tolerance of *Medicago truncatula*, *Plant Mol. Biol.* 59 (2005) 387–406.
36 <https://doi.org/10.1007/s11103-005-0184-z>.

- 1 [29] F. Colditz, K. Niehaus, F. Krajinski, Silencing of PR-10-like proteins in *Medicago*
2 *truncatula* results in an antagonistic induction of other PR proteins and in an increased
3 tolerance upon infection with the oomycete *Aphanomyces euteiches*, *Planta*. 226 (2007) 57–
4 71. <https://doi.org/10.1007/s00425-006-0466-y>.
- 5 [30] C. Ameline-Torregrosa, B. Dumas, F. Krajinski, M.-T. Esquerre-Tugaye, C. Jacquet,
6 Transcriptomic approaches to unravel plant–pathogen interactions in legumes, *Euphytica*. 147
7 (2006) 25–36. <https://doi.org/10.1007/s10681-006-6767-1>.
- 8 [31] S. Hosseini, M. Elfstrand, F. Heyman, D. Funck Jensen, M. Karlsson, Deciphering
9 common and specific transcriptional immune responses in pea towards the oomycete
10 pathogens *Aphanomyces euteiches* and *Phytophthora pisi*, *BMC Genomics*. 16 (2015).
11 <https://doi.org/10.1186/s12864-015-1829-1>.
- 12 [32] E. Gaulin, M.-A. Madoui, A. Bottin, C. Jacquet, C. Mathé, A. Couloux, P. Wincker, B.
13 Dumas, Transcriptome of *Aphanomyces euteiches*: new oomycete putative pathogenicity
14 factors and metabolic pathways, *PloS One*. 3 (2008) e1723.
15 <https://doi.org/10.1371/journal.pone.0001723>.
- 16 [33] X. Zhao, I.J. Misaghi, M.C. Hawes, Stimulation of border cell production in response
17 to increased carbon dioxide levels, *Plant Physiol*. 122 (2000) 181–188.
- 18 [34] M.C. Hawes, U. Gunawardena, S. Miyasaka, X. Zhao, The role of root border cells in
19 plant defense, *Trends Plant Sci*. 5 (2000) 128–133.
- 20 [35] U. Gunawardena, M. Rodriguez, D. Straney, J.T. Romeo, H.D. VanEtten, M.C.
21 Hawes, Tissue-specific localization of pea root infection by *Nectria haematococca*.
22 Mechanisms and consequences, *Plant Physiol*. 137 (2005) 1363–1374.
23 <https://doi.org/10.1104/pp.104.056366>.
- 24 [36] M.C. Hawes, G. Curlango-Rivera, Z. Xiong, J.O. Kessler, Roles of root border cells in
25 plant defense and regulation of rhizosphere microbial populations by extracellular DNA
26 ‘trapping,’ *Plant Soil*. 355 (2012) 1–16. <https://doi.org/10.1007/s11104-012-1218-3>.
- 27 [37] A. Driouich, C. Smith, M. Ropitiaux, M. Chambard, I. Boulogne, S. Bernard, M.-L.
28 Follet-Gueye, M. Vicié, J. Moore, Root extracellular traps versus neutrophil extracellular
29 traps in host defence, a case of functional convergence?, *Biol. Rev.* 94 (2019) 1685–1700.
30 <https://doi.org/10.1111/brv.12522>.
- 31 [38] M. Vicié, C. Santaella, S. Blanchet, A. Gateau, A. Driouich, Root Border-Like Cells
32 of *Arabidopsis*. Microscopical Characterization and Role in the Interaction with
33 Rhizobacteria, *Plant Physiol*. 138 (2005) 998–1008. <https://doi.org/10.1104/pp.104.051813>.
- 34 [39] F. Wen, H.D. VanEtten, G. Tsaprailis, M.C. Hawes, Extracellular Proteins in Pea Root
35 Tip and Border Cell Exudates, *Plant Physiol*. 143 (2007) 773–783.
36 <https://doi.org/10.1104/pp.106.091637>.

- 1 [40] F. Wen, G.J. White, H.D. VanEtten, Z. Xiong, M.C. Hawes, Extracellular DNA is
2 required for root tip resistance to fungal infection, *Plant Physiol.* 151 (2009) 820–829.
3 <https://doi.org/10.1104/pp.109.142067>.
- 4 [41] C. Durand, M. Vitré-Gibouin, M.L. Follet-Gueye, L. Duponchel, M. Moreau, P.
5 Lerouge, A. Driouich, The organization pattern of root border-like cells of *Arabidopsis* is
6 dependent on cell wall homogalacturonan, *Plant Physiol.* 150 (2009) 1411–1421.
7 <https://doi.org/10.1104/pp.109.136382>.
- 8 [42] M.A. Cannesan, C. Durand, C. Burel, C. Gangneux, P. Lerouge, T. Ishii, K. Laval, M.-
9 L. Follet-Gueye, A. Driouich, M. Vitré-Gibouin, Effect of arabinogalactan proteins from the
10 root caps of pea and *Brassica napus* on *Aphanomyces euteiches* zoospore chemotaxis and
11 germination, *Plant Physiol.* 159 (2012) 1658–1670. <https://doi.org/10.1104/pp.112.198507>.
- 12 [43] A. Driouich, M.-L. Follet-Gueye, M. Vitré-Gibouin, M. Hawes, Root border cells and
13 secretions as critical elements in plant host defense, *Curr. Opin. Plant Biol.* 16 (2013) 489–
14 495. <https://doi.org/10.1016/j.pbi.2013.06.010>.
- 15 [44] G.J. Griffin, M.G. Hale, F.J. Shay, Nature and quantity of sloughed organic matter
16 produced by roots of axenic peanut plants, *Soil Biol. Biochem.* 8 (1976) 29–32.
17 [https://doi.org/10.1016/0038-0717\(76\)90017-1](https://doi.org/10.1016/0038-0717(76)90017-1).
- 18 [45] M.C. Hawes, G. Curlango-Rivera, F. Wen, G.J. White, H.D. Vanetten, Z. Xiong,
19 Extracellular DNA: the tip of root defenses?, *Plant Sci. Int. J. Exp. Plant Biol.* 180 (2011)
20 741–745. <https://doi.org/10.1016/j.plantsci.2011.02.007>.
- 21 [46] E. Oburger, D.L. Jones, Sampling root exudates – Mission impossible?, *Rhizosphere.*
22 6 (2018) 116–133. <https://doi.org/10.1016/j.rhisph.2018.06.004>.
- 23 [47] H.P. Bais, T.L. Weir, L.G. Perry, S. Gilroy, J.M. Vivanco, The role of root exudates in
24 rhizosphere interactions with plants and other organisms, *Annu. Rev. Plant Biol.* 57 (2006)
25 233–266. <https://doi.org/10.1146/annurev.arplant.57.032905.105159>.
- 26 [48] D.V. Badri, J.M. Vivanco, Regulation and function of root exudates, *Plant Cell*
27 *Environ.* 32 (2009) 666–681. <https://doi.org/10.1111/j.1365-3040.2009.01926.x>.
- 28 [49] A. Jousset, L. Rochat, A. Lanoue, M. Bonkowski, C. Keel, S. Scheu, Plants Respond
29 to Pathogen Infection by Enhancing the Antifungal Gene Expression of Root-Associated
30 Bacteria, *Mol. Plant-Microbe Interactions®.* 24 (2010) 352–358.
31 <https://doi.org/10.1094/MPMI-09-10-0208>.
- 32 [50] A. Lanoue, V. Burlat, G.J. Henkes, I. Koch, U. Schurr, U.S.R. Röse, De novo
33 biosynthesis of defense root exudates in response to *Fusarium* attack in barley, *New Phytol.*
34 185 (2010) 577–588. <https://doi.org/10.1111/j.1469-8137.2009.03066.x>.
- 35 [51] N. Eisenhauer, A. Lanoue, T. Strecker, S. Scheu, K. Steinauer, M.P. Thakur, L.

- 1 Mommer, Root biomass and exudates link plant diversity with soil bacterial and fungal
2 biomass, *Sci. Rep.* 7 (2017) 1–8. <https://doi.org/10.1038/srep44641>.
- 3 [52] R. Yokosawa, S. Kuninaga, H. Sekizaki, *Aphanomyces euteiches* zoospore attractant
4 isolated from pea root: Prunetin, *Jpn. J. Phytopathol.* 52 (1986) 809–816.
5 <https://doi.org/10.3186/jjphytopath.52.809>.
- 6 [53] U. Baetz, E. Martinoia, Root exudates: the hidden part of plant defense, *Trends Plant*
7 *Sci.* 19 (2014) 90–98. <https://doi.org/10.1016/j.tplants.2013.11.006>.
- 8 [54] E. Billard, A. Quillévéré-Hamard, C. Lavaud, M.L. Pilet-Nayel, C.L. May, Testing of
9 life history traits of a soilborne pathogen in vitro: Do characteristics of oospores change
10 according the strains of *Aphanomyces euteiches* and the host plant infected by the pathogen?,
11 *J. Phytopathol.* 167 (2019) 313–320. <https://doi.org/10.1111/jph.12799>.
- 12 [55] E. Wicker, M. Hullé, F. Rouxel, Pathogenic characteristics of isolates of
13 *Aphanomyces euteiches* from pea in France, *Plant Pathol.* 50 (2001) 433–442.
14 <https://doi.org/10.1046/j.1365-3059.2001.00590.x>.
- 15 [56] A. Carreras, S. Bernard, G. Durambur, B. Gügi, C. Loutelier, B. Pawlak, I. Boulogne,
16 M. Vicié, A. Driouich, D. Goffner, M.-L. Follet-Gueye, In vitro characterization of root
17 extracellular trap and exudates of three Sahelian woody plant species, *Planta.* 251 (2019) 19.
18 <https://doi.org/10.1007/s00425-019-03302-3>.
- 19 [57] O. Margie, C. Palmer, I. Chin-Sang, C. elegans Chemotaxis Assay, *J. Vis. Exp. JoVE.*
20 (2013). <https://doi.org/10.3791/50069>.
- 21 [58] N.D. Warnock, L. Wilson, C. Patten, C.C. Fleming, A.G. Maule, J.J. Dalzell,
22 Nematode neuropeptides as transgenic nematocides, *PLOS Pathog.* 13 (2017) e1006237.
23 <https://doi.org/10.1371/journal.ppat.1006237>.
- 24 [59] T.R. Fleming, A.G. Maule, C.C. Fleming, Chemosensory Responses of Plant Parasitic
25 Nematodes to Selected Phytochemicals Reveal Long-Term Habituation Traits, *J. Nematol.* 49
26 (2017) 462–471.
- 27 [60] T.N. Shivakumara, T.K. Dutta, U. Rao, A novel in vitro chemotaxis bioassay to assess
28 the response of *Meloidogyne incognita* towards various test compounds, *J. Nematol.* 50
29 (2018) 487–494. <https://doi.org/10.21307/jofnem-2018-047>.
- 30 [61] F.L. Mauff, C. Loutelier-Bourhis, M. Bardor, C. Berard, A. Doucet, M.-A. D’Aoust,
31 L.-P. Vezina, A. Driouich, M.M.-J. Couture, P. Lerouge, Cell wall biochemical alterations
32 during *Agrobacterium*-mediated expression of haemagglutinin-based influenza virus-like
33 vaccine particles in tobacco, *Plant Biotechnol. J.* 15 (2017) 285–296.
34 <https://doi.org/10.1111/pbi.12607>.
- 35 [62] G.J. van Holst, A.E. Clarke, Organ-Specific Arabinogalactan-Proteins of *Lycopersicon*

1 peruvianum (Mill) Demonstrated by Crossed Electrophoresis, *Plant Physiol.* 80 (1986) 786–
2 789. <https://doi.org/10.1104/pp.80.3.786>.

3 [63] E. Nguema-Ona, S. Coimbra, M. Vitré-Gibouin, J.-C. Mollet, A. Driouich,
4 Arabinogalactan proteins in root and pollen-tube cells: distribution and functional aspects,
5 *Ann. Bot.* 110 (2012) 383–404. <https://doi.org/10.1093/aob/mcs143>.

6 [64] K. Kitazawa, T. Tryfona, Y. Yoshimi, Y. Hayashi, S. Kawauchi, L. Antonov, H.
7 Tanaka, T. Takahashi, S. Kaneko, P. Dupree, Y. Tsumuraya, T. Kotake, β -Galactosyl Yariv
8 Reagent Binds to the β -1,3-Galactan of Arabinogalactan Proteins, *Plant Physiol.* 161 (2013)
9 1117–1126. <https://doi.org/10.1104/pp.112.211722>.

10 [65] M. Dueñas, I. Estrella, T. Hernández, Occurrence of phenolic compounds in the seed
11 coat and the cotyledon of peas (*Pisum sativum* L.), *Eur. Food Res. Technol.* 219 (2004) 116–
12 123. <https://doi.org/10.1007/s00217-004-0938-x>.

13 [66] R.M. Celoy, H.D. VanEtten, (+)-Pisatin biosynthesis: From (–) enantiomeric
14 intermediates via an achiral 7,2'-dihydroxy-4',5'-methylenedioxyisoflav-3-ene,
15 *Phytochemistry*. 98 (2014) 120–127. <https://doi.org/10.1016/j.phytochem.2013.10.017>.

16 [67] A. Troszyńska, I. Estrella, M.L. López-Amóres, T. Hernández, Antioxidant Activity of
17 Pea (*Pisum sativum* L.) Seed Coat Acetone Extract, *LWT - Food Sci. Technol.* 35 (2002)
18 158–164. <https://doi.org/10.1006/fstl.2001.0831>.

19 [68] J.W. Mansfield, D.A. Widdowson, The metabolism of wyerone acid (a phytoalexin
20 from *Vicia faba* L.) by *Botrytis fabae* and *B. cinerea*, *Physiol. Plant Pathol.* 3 (1973) 393–404.
21 [https://doi.org/10.1016/0048-4059\(73\)90012-X](https://doi.org/10.1016/0048-4059(73)90012-X).

22 [69] I.M. Abu-Reidah, M. del M. Contreras, D. Arráez-Román, A. Fernández-Gutiérrez, A.
23 Segura-Carretero, UHPLC-ESI-QTOF-MS-based metabolic profiling of *Vicia faba* L.
24 (Fabaceae) seeds as a key strategy for characterization in foodomics, *ELECTROPHORESIS*.
25 35 (2014) 1571–1581. <https://doi.org/10.1002/elps.201300646>.

26 [70] P. van West, A.A. Appiah, N.A.R. Gow, Advances in research on oomycete root
27 pathogens, *Physiol. Mol. Plant Pathol.* 62 (2003) 99–113. [https://doi.org/10.1016/S0885-5765\(03\)00044-4](https://doi.org/10.1016/S0885-5765(03)00044-4).

29 [71] J.N. Cameron, M.J. Carlile, Binding of isovaleraldehyde, an attractant, to zoospores of
30 the fungus *Phytophthora palmivora* in relation to zoospore chemotaxis, *J. Cell Sci.* 49 (1981)
31 273–281.

32 [72] R. Yokosawa, S. Kuninaga, *Aphanomyces raphani* zoospore attractant isolated from
33 cabbage: indole-3-aldehyde, *Nihon Shokubutsu Byori Gakkaiho Ann. Phytopathol. Soc. Jpn.*
34 (1979). <http://agris.fao.org/agris-search/search.do?recordID=US201302810275> (accessed July
35 19, 2017).

- 1 [73] P. Kong, B.M. Tyler, P.A. Richardson, B.W.K. Lee, Z.S. Zhou, C. Hong, Zoospore
2 interspecific signaling promotes plant infection by *Phytophthora*, *BMC Microbiol.* 10 (2010)
3 313. <https://doi.org/10.1186/1471-2180-10-313>.
- 4 [74] P. Kong, J.M. McDowell, C. Hong, Zoospore exudates from *Phytophthora nicotianae*
5 affect immune responses in *Arabidopsis*, *PLoS ONE*. 12 (2017).
6 <https://doi.org/10.1371/journal.pone.0180523>.
- 7 [75] 2014 Meeting | Compounds from Zoospore Exudate Serve as a Signal to Promote
8 Zoosporic Germination and Infection of *Phytophthora erythroseptica*, (n.d.).
9 https://www.apsnet.org/meetings/Documents/2014_meeting_abstracts/aps2014abO161.htm
10 (accessed April 12, 2020).
- 11 [76] X. Zhao, M. Schmitt, M.C. Hawes, Species-dependent effects of border cell and root
12 tip exudates on nematode behavior, *Phytopathology*. 90 (2000) 1239–1245.
13 <https://doi.org/10.1094/PHYTO.2000.90.11.1239>.
- 14 [77] R.H.C. Curtis, Plant-nematode interactions: environmental signals detected by the
15 nematode's chemosensory organs control changes in the surface cuticle and behaviour,
16 *Parasite Paris Fr.* 15 (2008) 310–316. <https://doi.org/10.1051/parasite/2008153310>.
- 17 [78] R.N. Perry, M. Moens, J.L. Starr, *Root-knot Nematodes*, CABI, 2009.
- 18 [79] G. Yang, B. Zhou, X. Zhang, Z. Zhang, Y. Wu, Y. Zhang, S. Lü, Q. Zou, Y. Gao, L.
19 Teng, Effects of Tomato Root Exudates on *Meloidogyne incognita*, *PloS One*. 11 (2016)
20 e0154675. <https://doi.org/10.1371/journal.pone.0154675>.
- 21 [80] M. Mizutani, Y. Hashidoko, S. Tahara, Factors responsible for inhibiting the motility
22 of zoospores of the phytopathogenic fungus *Aphanomyces cochlioides* isolated from the non-
23 host plant *Portulaca oleracea*, *FEBS Lett.* 438 (1998) 236–240.
- 24 [81] N. Bazghaleh, P. Prashar, R.W. Purves, A. Vandenberg, Polyphenolic Composition of
25 Lentil Roots in Response to Infection by *Aphanomyces euteiches*, *Front. Plant Sci.* 9 (2018)
26 1131. <https://doi.org/10.3389/fpls.2018.01131>.
- 27 [82] M. Wink, F. Botschen, C. Gosmann, H. Schäfer, P.G. Waterman, Chemotaxonomy
28 Seen from a Phylogenetic Perspective and Evolution of Secondary Metabolism, in: *Annu.*
29 *Plant Rev. Vol. 40 Biochem. Plant Second. Metab.*, John Wiley & Sons, Ltd, 2010: pp. 364–
30 433. <https://doi.org/10.1002/9781444320503.ch7>.
- 31 [83] F.M. Afendi, T. Okada, M. Yamazaki, A. Hirai-Morita, Y. Nakamura, K. Nakamura,
32 S. Ikeda, H. Takahashi, M. Altaf-Ul-Amin, L.K. Darusman, K. Saito, S. Kanaya, KNApSACk
33 family databases: integrated metabolite-plant species databases for multifaceted plant
34 research, *Plant Cell Physiol.* 53 (2012) e1. <https://doi.org/10.1093/pcp/pcr165>.
- 35 [84] H.F. Nawar, J.O. Kuti, Weyerone Acid Phytoalexin Synthesis and Peroxidase Activity

- 1 as Markers for Resistance of Broad Beans to Chocolate Spot Disease, *J. Phytopathol.* 151
2 (2003) 564–570. <https://doi.org/10.1046/j.1439-0434.2003.00732.x>.
- 3 [85] A. Chaboud, Isolation, purification and chemical composition of maize root cap slime,
4 *Plant Soil.* 73 (1983) 395–402. <https://doi.org/10.1007/BF02184316>.
- 5 [86] E.M. Knee, F.C. Gong, M. Gao, M. Teplitski, A.R. Jones, A. Foxworthy, A.J. Mort,
6 W.D. Bauer, Root mucilage from pea and its utilization by rhizosphere bacteria as a sole
7 carbon source, *Mol. Plant-Microbe Interact. MPMI.* 14 (2001) 775–784.
8 <https://doi.org/10.1094/MPMI.2001.14.6.775>.
- 9 [87] J.M. Hinch, A.E. Clarke, Adhesion of fungal zoospores to root surfaces is mediated by
10 carbohydrate determinants of the root slime, *Physiol. Plant Pathol.* 16 (1980) 303–IN2.
11 [https://doi.org/10.1016/S0048-4059\(80\)80001-4](https://doi.org/10.1016/S0048-4059(80)80001-4).
- 12 [88] F. Xie, A. Williams, A. Edwards, J.A. Downie, A plant arabinogalactan-like
13 glycoprotein promotes a novel type of polar surface attachment by *Rhizobium*
14 *leguminosarum*, *Mol. Plant-Microbe Interact. MPMI.* 25 (2012) 250–258.
15 <https://doi.org/10.1094/MPMI-08-11-0211>.
- 16 [89] J. Mravec, X. Guo, A.R. Hansen, J. Schückel, S.K. Kračun, M.D. Mikkelsen, G.
17 Mouille, I.E. Johansen, P. Ulvskov, D.S. Domozych, W.G.T. Willats, Pea Border Cell
18 Maturation and Release Involve Complex Cell Wall Structural Dynamics, *Plant Physiol.* 174
19 (2017) 1051–1066. <https://doi.org/10.1104/pp.16.00097>.
- 20 [90] E. Nguema-Ona, M. Vicié-Gibouin, M. Gotté, B. Plancot, P. Lerouge, M. Bardor, A.
21 Driouich, Cell wall O-glycoproteins and N-glycoproteins: aspects of biosynthesis and
22 function, *Front. Plant Sci.* 5 (2014). <https://doi.org/10.3389/fpls.2014.00499>.
- 23 [91] A.S. Koroney, C. Plasson, B. Pawlak, R. Sidikou, A. Driouich, L. Menu-Bouaouiche,
24 M. Vicié-Gibouin, Root exudate of *Solanum tuberosum* is enriched in galactose-containing
25 molecules and impacts the growth of *Pectobacterium atrosepticum*, *Ann. Bot.* 118 (2016)
26 797–808. <https://doi.org/10.1093/aob/mcw128>.
- 27 [92] E. Nguema-Ona, A. Bannigan, L. Chevalier, T.I. Baskin, A. Driouich, Disruption of
28 arabinogalactan proteins disorganizes cortical microtubules in the root of *Arabidopsis*
29 *thaliana*, *Plant J.* 52 (2007) 240–251. <https://doi.org/10.1111/j.1365-313X.2007.03224.x>.
- 30 [93] E. Nguema-Ona, M. Vicié-Gibouin, M.-A. Cannesan, A. Driouich, Arabinogalactan
31 proteins in root-microbe interactions, *Trends Plant Sci.* 18 (2013) 440–449.
32 <https://doi.org/10.1016/j.tplants.2013.03.006>.
- 33 [94] M. Larousse, E. Galiana, Microbial Partnerships of Pathogenic Oomycetes, *PLoS*
34 *Pathog.* 13 (2017) e1006028. <https://doi.org/10.1371/journal.ppat.1006028>.
- 35 [95] E. Gaulin, M.J.C. Pel, L. Camborde, H. San-Clemente, S. Courbier, M.-A. Dupouy, J.

Lengellé, M. Veyssiere, A. Le Ru, F. Grandjean, R. Cordaux, B. Moumen, C. Gilbert, L.M. Cano, J.-M. Aury, J. Guy, P. Wincker, O. Bouchez, C. Klopp, B. Dumas, Genomics analysis of *Aphanomyces* spp. identifies a new class of oomycete effector associated with host adaptation, *BMC Biol.* 16 (2018) 43. <https://doi.org/10.1186/s12915-018-0508-5>.

[96] Ayers, Factors affecting the pectolytic activity of *Aphanomyces euteiches* *in vitro* and in infected tissue, (n.d.).

[97] Ayers, Purification and properties of the endopolygalacturonase of *Aphanomyces euteiches*, (n.d.).

[98] R. Kjølner, S. Rosendahl, Enzymatic Activity of the Mycelium Compared with Oospore Development During Infection of Pea Roots by *Aphanomyces euteiches*, *Phytopathology*. 88 (1998) 992–996. <https://doi.org/10.1094/PHTO.1998.88.9.992>.

[99] A. Moussart, M.N. Even, A. Lesné, B. Tivoli, Successive legumes tested in a greenhouse crop rotation experiment modify the inoculum potential of soils naturally infested by *Aphanomyces euteiches*, *Plant Pathol.* 62 (2013) 545–551. <https://doi.org/10.1111/j.1365-3059.2012.02679.x>.

[100] S. Selim, J. Sanssené, S. Rossard, J. Courtois, Systemic induction of the defensin and phytoalexin pisatin pathways in pea (*Pisum sativum*) against *Aphanomyces euteiches* by acetylated and nonacetylated oligogalacturonides, *Molecules*. 22 (2017) 1017. <https://doi.org/10.3390/molecules22061017>.

~~[100] K. Steinauer, A. Chatzinotas, N. Eisenhauer, Root exudate cocktails: the link between plant diversity and soil microorganisms?, *Ecol. Evol.* 6 (2016) 7387–7396. <https://doi.org/10.1002/ece3.2454>.~~

~~[101] J. Sasse, E. Martinoia, T. Northen, Feed Your Friends: Do Plant Exudates Shape the Root Microbiome?, *Trends Plant Sci.* 23 (2018) 25–41. <https://doi.org/10.1016/j.tplants.2017.09.003>.~~

~~[102] K. Zhalnina, K.B. Louie, Z. Hao, N. Mansoori, U.N. da Rocha, S. Shi, H. Cho, U. Karaoz, D. Loqué, B.P. Bowen, M.K. Firestone, T.R. Northen, E.L. Brodie, Dynamic root exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial community assembly, *Nat. Microbiol.* 3 (2018) 470–480. <https://doi.org/10.1038/s41564-018-0129-3>.~~

~~[103] A.M. Wentzell, D.J. Kliebenstein, Genotype, age, tissue, and environment regulate the structural outcome of glucosinolate activation, *Plant Physiol.* 147 (2008) 415–428. <https://doi.org/10.1104/pp.107.115279>.~~

~~[104] B.J.W. Chen, R. Hajiboland, S. Bahrami Rad, N. Moradtalab, N.P.R. Anten, Presence of Belowground Neighbors Activates Defense Pathways at the Expense of Growth in Tobacco Plants, *Front. Plant Sci.* 10 (2019). <https://doi.org/10.3389/fpls.2019.00751>.~~

1 **Figures**




Figure 1

2

3 **Fig.1. A. euteiches** colonization in pea and faba bean roots at early stage of infection. at

4 1, 2 and 3 dpi by epifluorescence microscopy. dpi, days post-inoculation; non-germinated

1 cysts (white triangles), germinated cyst (white arrows) and mycelium (arrows heads). Scale

2 bars: 500 μ m (A); 100 μ m (B).




Figure 2

Fig. 2 Quantification of *A. euteiches* DNA in pea and faba bean roots

A, DNA quantification of *A. euteiches* by qPCR in pea and faba bean roots co-inoculated and co-cultivated (condition 1). **B**, DNA quantification of *A. euteiches* in pea and faba bean roots inoculated separately and co-cultivated (condition 2). **C**, DNA quantification of *A. euteiches* in pea roots inoculated and cultivated in the presence of faba bean root exudates from non-inoculated seedlings (condition 3). For each condition, control is pea (black colored histograms) or faba bean (grey colored histograms) inoculated and cultivated separately. Each histogram represents the mean of three biological replicates ($n=3$) and error bars indicate the standard error. Statistical analysis was performed using Mann-Whitney test ($*P<0.05$) (5% level of significance). dpi, days post-inoculation.

10




Figure 3

Fig. 3. Effect of pea and faba bean root exudates on *A. euteiches* behavior

A, Experimental device for chemotaxis and cyst germination assays. Zoospores were put in the middle well. Water, used as control, was put in the right. Water, Gum Arabic or root exudates were been put in the left. Chemotaxis index was calculated as: $[(\text{number of zoospores within the Z1} - \text{number of zoospores within the Z2}) / (\text{number of zoospores which made a choice Z1+Z2})]$. The percentage of cysts that germinate is determined for each zones as : $[(\text{Number of germinated cysts per zone} / \text{Total number of cysts per zone}) \times 100]$.

B, *A. euteiches* zoospores chemotaxis response. **C**, Germination of cysts in zones 1 and 2. **B**, **C**, Statistical analysis was performed using Mann-Whitney test comparing the test compounds with water control (* $P < 0.05$; ** $P < 0.01$) (5% level of significance). Each histogram represents the mean of five biological replicates ($n=5$) excepted for GA ($n=3$). Error bars indicate the standard error.

D, Effect of root exudates from non-inoculated seedlings on mycelium growth. Surfaces of mycelium were compared with water control condition. Error bars indicate the standard error of five biological replicates ($n=5$) excepted for water control conditions ($n=3$). Statistical analysis was performed using Mann-Whitney test comparing the test compounds with water control (* $P < 0.05$) (5% level of significance). Ae, *A. euteiches* ; GA, Gum Arabic ; FRE, faba bean root exudates from non-inoculated seedlings ; PRE, pea root exudates from non inoculated seedlings ; FRE +Ae, faba bean root exudates from inoculated seedlings ; PRE + Ae , pea root exudates from inoculated seedlings ; Z1, zone 1; Z2, zone 2.

Tables

Table 1: List of polyphenols identified in pea root exudates.

RT (min)	Compound assignment	Compound class	MW	[M-H]-	[M+H]+	Fragments (ES-)	Fragments (ES+)	References
3.39	tryptophane ^a	amino acid	204	203	205	188; 146		standard
5.38	quercetin-3-rutinoside	flavonol	610	609	611	131	301 [M+H-gluc-rha]+ 287; 141	Mullen et al. 2003
5.76	<i>p</i> -coumaric acid ^a	phenolic	164	163	165	119	147; 119	standard
6.32	trans-ferulic acid ^a	phenolic	194	193	195	178; 134	177	standard
7.1	DMDI ^a	isoflavanol	302	301		273; 163		Celoy and VanEtten. 2013
10.22	DMD ^b	isoflavone	298	297	299		321 [M+Na]+	Celoy and VanEtten. 2013
11.19	formononetin ^a	isoflavone	268	267	269	252		standard
11.76	pisatin ^a	pterocarpan	314	313		299; 181		standard

^a7,2'-dihydroxy-4',5'-methylene-dioxyisoflavanol

^b7,2'-dihydroxy-4',5'-methylenedioxyisoflavone

Table 2: List of polyphenols identified in faba bean root exudates.

RT (min)	Compound assignment	Compound class	MW	[M-H]-	[M+H]+	Fragments (ES-)	Fragments (ES+)	References
3.39	tryptophane ^a	amino acid	204	203	205		188.1; 146.0	standard
6.32	sinapoyl-glucoside	phenolic	386	385		269; 223 [M-H-Gluc]-; 171		Neugart et al. 2015
8.11	wyerone epoxide	furanoacetylene	274	273		224; 183		Hargreaves et al. 1976
10.41	dihydrowyerol	furanoacetylene	262	261	263	217; 191	207	Mansfield et al. 1979
10.57	wyerone acid	furanoacetylene	244	243	245	225	227	Mansfield and Widdowson. 1973
11.15	formononetin ^a	isoflavone	268	267	269	252		standard
12.09	medicarpin	pterocarpan	270	269	271	254; 163	165; 137	Abu-Reidah et al. 2014

4

5

Table 3. AGP quantification in root exudates at three days after inoculation

To quantify AGP, the quantification of β -glucosyl Yariv precipitate in AIR of root exudates for peas and faba beans was assessed. Controls consist in peas or faba beans non-inoculated. Each histogram represents the mean of four technical replicates ($n=4$) $\pm$ the standard error. For statistical analysis Mann-Whitney nonparametric test (5% level of significance) was performed ($*P<0.05$). Ae, *A. euteiches*; AGP, arabinogalactan protein.

	AGPs quantification (mg ml ⁻¹)	
	Control	Plant + <i>Ae</i>
Pea	0.15 ± 0.09	0.42 ± 0.13 *
Faba bean	0.14 ± 0.15	0.05 ± 0.09 ns

1

2