



Decreased susceptibility to didecyldimethylammonium chloride among clinical strains of *Pseudomonas aeruginosa*, a ten years retrospective study, at the University Hospital of Caen, France.

Marine Pottier, François Gravey, Sophie Castagnet, Michel Auzou, Leduc Guillaume, Marion Demuynck, Albertine Léon, Simon Le Hello

► To cite this version:

Marine Pottier, François Gravey, Sophie Castagnet, Michel Auzou, Leduc Guillaume, et al.. Decreased susceptibility to didecyldimethylammonium chloride among clinical strains of *Pseudomonas aeruginosa*, a ten years retrospective study, at the University Hospital of Caen, France.. The 31st European Congress of Clinical Microbiology & Infectious Diseases (ECCMID), Jul 2021, online, Austria. hal-03418585

HAL Id: hal-03418585

<https://normandie-univ.hal.science/hal-03418585>

Submitted on 7 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Decreased susceptibility to didecyldimethylammonium chloride among clinical strains of *Pseudomonas aeruginosa*, a ten years retrospective study, at the University Hospital of Caen, France.

Marine Pottier^{1, 2, 3}, François Gravey¹, Sophie Castagnet², Michel Auzou⁴, Guillaume Leduc⁴, Marion Demuynck¹, Albertine Léon^{2, 3, *}, Simon Le Hello^{1, 4, 5, *}

¹ Normandie Univ, UNICAEN, GRAM 2.0 - Caen (France), ² Laboratoire LABÉO, Saint Contest - Caen (France), ³ Normandie Univ, UNICAEN, U2RM - Caen (France), ⁴ Service de microbiologie, CHU - Caen (France), ⁵ Service d'hygiène hospitalière, CHU - Caen (France), ^{*} Equivalent coordination

INTRODUCTION

- Didecyldimethylammonium chloride (DDAC) is a quaternary ammonium compound (QACs) used in several disinfection process in hospital and in veterinary environments.
- Throughout 2018, numerous strains of *Pseudomonas aeruginosa* (PSAE) isolated in the University Hospital (UH) of Caen showed a **decreased susceptibility to DDAC**, according to the **NF EN-13727+A2 guidelines** (1).
- Mechanisms** involved are **not fully understood**, but first studies suggested the **involvement of the resistance-nodulation-division efflux pump MexAB-OprM** (2).
- Decreased susceptibility to disinfectant agents is a **major concern to public health** as it allows pathogens to **persist in their environment**.

AIM

To assess the emergence of DDAC decreased susceptibility among clinical strains of PSAE isolated in a French UH over the last decade.

METHOD

Over the period 2011 – 2020 :

Random inclusion of **10 strains / years** of clinical PSAE strains (100 in total) from the microbiology department of the UH of Caen.


Determination of **Minimum inhibitory concentrations (MICs)**, in triplicate by broth dilution method.

According to the International Organization for Standardization (3).

Decrease susceptibility cutoff corresponding to the **concentration of DDAC in a disinfectant solution** in regard of the manufacturer's instructions.

MIC < 62.9 mg.L⁻¹
Susceptible strain (S)

MIC > 62.9 mg.L⁻¹
Decrease susceptible strain (DS)



RESULTS

Table I: Repartition of strains origins.

Isolation year	Strain origin					
	Respiratory	Urinary	Blood culture/Catheter	Digestive	Surgical pus	Others ^a
2011	1	1	4	-	2	2
2012	2	-	6	-	2	-
2013	3	1	2	2	1	1
2014	2	4	-	1	2	1
2015	2	3	2	-	2	1
2016	2	3	-	4	1	-
2017	2	6	-	-	-	2
2018	6	1	-	3	-	-
2019	3	-	-	2	2	3
2020	2	2	1	2	1	2
Total	25	21	15	14	13	12

Notes

^a The following sample types were categorized as "Others": biopsy, ocular, auricular, genital, wound and unknown origin.

Among the 100 strains, the three main origins were (Table I):

- 1) **respiratory** samples → 25 strains
- 2) **urinary** samples → 21 strains
- 3) **blood culture or catheter** samples → 15 strains

CONCLUSIONS

- The decreased susceptibility to DDAC is **not a new phenomenon** but steady over time and still present.
- As **nearly one-third** of the tested strains have a decreased sensitivity to CDMA, this could explain, in part, why PSAE is a **persistent bacteria in hospital**.
- Studies are ongoing to 1) Test PSAE strains isolated from **hospital environment** samples and from **animals** 2) **Explore the mechanism(s) involved** in DS to DDAC.

- For **each year** (Figure I) and **each type of sample tested** (Figure II) at least **one DS strain** was found.
- Overall, **28 of all the tested strains** presented a DS to DDAC.
- MIC range for DS strains was : **64 mg.L⁻¹** or **128 mg.L⁻¹**.




Figure I: Temporal evolution of MICs to DDAC for PSAE strains isolated from patients (n=100). Red dot: strains with DDAC decreased susceptibility; green dot: DDAC susceptible strains; dot size: headcount.

- The annual average rate of DS to DDAC was 28 % (Figure I).
- This annual rate varied:
 - from 10 % for years 2015, 2017 and 2020,
 - to 60 % for 2014.
- No significant variation** of MICs **between** and **throughout years** (Kruskal-Wallis test, *p*-value > 0,1).




Figure II: Dispersion of MICs to DDAC according to strains origin (n=100). Red: strains with DDAC decreased susceptibility; green: DDAC susceptible strains; dot size: headcount.

- For a given **sample type**, the average rate of DS to DDAC was 27 % (Figure II).
- DS to DDAC was the mainly found in:
 - 1) **respiratory** samples (n= 8/25 → **32 %**),
 - 2) **surgical pus** samples (n= 4/13 → **31 %**),
 - 3) **urinary** samples (n= 6/21 → **29 %**) and **digestive** samples (n= 4/14 → **29 %**).

REFERENCES

- (1) **Association française de normalisation**. Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity in the medical area - Test method and requirements (phase 2, step 1). *NF EN 13727+A2*, 2015. p. 56.
- (2) Dreier J, Ruggerone P. Interaction of antibacterial compounds with RND efflux pumps in *Pseudomonas aeruginosa*. *Front Microbiol* [Internet]. 2015 Jul 8;6. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4495556/>
- (3) **International Organization for Standardization**. Clinical laboratory testing and *in vitro* diagnostic test systems - Susceptibility testing of infectious agents and evaluation of performance of antimicrobial susceptibility test devices - Part 1: Reference method for testing the *in vitro* activity of antimicrobial agents against rapidly growing aerobic bacteria involved in infectious diseases. *ISO 20776-1* (2006), 2006.

ACKNOWLEDGEMENTS



CONTACT INFORMATION

Mme Marine Pottier (PhD-student)

Mail: marine.pottier@laboratoire-labeo.fr

Address: EA2656 GRAM2.0 – CHU de Caen, service de microbiologie, partie recherche, 3^e étage