



**HAL**  
open science

## Biofilm formation of *Staphylococcus lugdunensis* strains is associated with clonal lineage and dependent on iron availability

L Destruel, S Dahyot, X Argemi, M Grand, G Prevost, M Pestel-Caron

### ► To cite this version:

L Destruel, S Dahyot, X Argemi, M Grand, G Prevost, et al.. Biofilm formation of *Staphylococcus lugdunensis* strains is associated with clonal lineage and dependent on iron availability. 32nd European Congress of Clinical Microbiology & Infectious Diseases ECCMID, Apr 2022, Lisbonne, France. hal-03706333

**HAL Id: hal-03706333**

**<https://normandie-univ.hal.science/hal-03706333>**

Submitted on 27 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Biofilm formation of *Staphylococcus lugdunensis* strains is associated with clonal lineage and dependent on iron availability

L. Destruel<sup>1</sup>, S. Dahyot<sup>1,2</sup>, X. Argemi<sup>3,4</sup>, M. Grand<sup>1</sup>, G. Prevost<sup>3</sup>, M. Pestel-Caron<sup>1,2</sup>

<sup>1</sup>Normandie Univ, UNIROUEN, UNICAEN, Inserm UMR 1311 DYNAMICURE, 76000 Rouen, France, <sup>2</sup>Normandie Univ, UNIROUEN, UNICAEN, Inserm UMR 1311 DYNAMICURE, CHU Rouen, Department of microbiology, F-76000 Rouen, France, <sup>3</sup>VBP EA7290, Fédération de Médecine Translationnelle de Strasbourg, Institut de Bactériologie, Centre Hospitalier Régional Universitaire, Université de Strasbourg, Strasbourg, France, <sup>4</sup>Maladies Infectieuses et Tropicales, Hôpitaux Universitaires de Strasbourg, Strasbourg, France

## Background

*Staphylococcus lugdunensis* is a virulent coagulase-negative staphylococcal species<sup>1</sup>. The ability to form biofilm appears to be one of its main virulence factors<sup>2</sup>, and is promoted by iron-limitation condition<sup>3</sup>. Nevertheless, little is known about biofilm-associated genotypes.

In this context, the objective of this study was to compare the impact of iron restriction on **fitness** and **biofilm formation** of strains belonging to all clonal complexes (CCs) defined by Multilocus sequence typing<sup>4</sup> to date.

## Materials

### Strains

- The study included **49** clinical strains from **different clonal complexes** and **geographical origins** (French and Swedish cities) (Table 1)

Clonal complex	CC1	CC2	CC3	CC4	CC5	CC6	CC7	Singleton*
Number of strains	11	5	11	4	5	5	4	4
Geographical origins	Strasbourg, Rouen, Tours, Kronenberg, Nancy	Strasbourg, Montpellier, Nantes, Nancy, Rouen	Rouen, Strasbourg	Strasbourg, Rouen, Bordeaux	Strasbourg, Bordeaux, Rouen	Kronenberg, Nantes, Strasbourg, Rouen	Strasbourg, Rouen	Rouen, Strasbourg

Table 1: *S. lugdunensis* strains categorized by clonal complexes and geographical origins. \*Singleton Sequence Type (ST)

### Fitness

- Growth kinetics** over 24 hours performed in triplicates for all strains (Tecan spark®):

✓ In **rich media** (Tryptic Soy Broth [TSB])

✓ In **iron-restricted condition**

(RPMI 1640 with 1% casamino acids)

✓ Determination of **generation time** ( $G = \ln 2 / \mu$ ) and **maximal growth rate** ( $\mu$ ) (Figure 1)




Figure 1 : Diagram of a growth curve

### Biofilm formation




Figure 2: Biofilm formation on polystyrene microtiter plates

- 24h biofilm formation** studied using **crystal violet staining** method on 96-well polystyrene microtiter plates in quadruplicates:

✓ In **rich media** (TSB with 1% glucose) for all strains

✓ In **iron-restricted condition** for 34 strains (Figure 2)

✓ Using *S. epidermidis* ATCC 35984 as biofilm positive control

## Results

### Fitness

	Generation time (G)	Maximal growth rate ( $\mu$ )
<b>Rich medium (TSB)</b>	37.34 minutes $\pm$ 4.13	1.139 $\pm$ 0.132
<b>Iron restricted medium (RPMI + 1% CAS)</b>	80.72 minutes $\pm$ 33.14	0.634 $\pm$ 0.20

Table 2: Generation time and maximal growth rate of 49 *S. lugdunensis* strains in rich and iron restricted media

- No difference** in growth either according to the strains or according to the CCs, whatever the medium (Table 2)

### Biofilm formation

#### In rich medium




Figure 3: Biofilm formation of 49 *S. lugdunensis* strains after a 24h incubation at 37°C in rich medium (TSB + 1% glucose) according to the CC (OD mean values). Error bars represent the standard deviation within each CC.

- All strains were biofilm producers ( $DO_{590nm} > 0,1$ ) in rich medium after 24h
- CC2** strains produced significantly **more biofilm** than **CC4** ( $p = 0.0203$ ), **CC5** ( $p = 0.0215$ ) and **singleton STs** ( $p = 0.0045$ ) strains (Figure 3)

#### In iron-restricted conditions




Figure 4: Biofilm formation of 34 *S. lugdunensis* strains after a 24h incubation at 37°C in rich medium (TSB + 1% glucose [full bars]) or iron-restricted medium (RPMI + 1% casamino acids [hatched bars]) according to the CC (OD mean values). Error bars represent the standard deviation within each CC.

- In **iron-restricted conditions**, **CC1** ( $p = 0.0079$ ), **CC2** ( $p = 0.0079$ ) and **CC3** ( $p = 0.026$ ) strains produced significantly **less biofilm** than in rich medium
- On the contrary, **CC6** ( $p = 0.0079$ ) strains were significantly **more productive** (Figure 4)

## Conclusions

- This study shows for the first time that the **biofilm-forming ability** of *S. lugdunensis* strains is strongly **correlated with CC affiliation**.
- In addition, two different **CC-dependent behaviors** are observed in an **iron-restricted environment**. These results are to be confirmed with a larger number of strains.