

Sensory Prediction and Repetition Suppression in the tactile modality as early markers of executive attention development at preschool age

Marie Anquetil, Victoria Dumont, Anne-Lise Marais, Nadege Roche-Labarbe,

Sandrine Rossi

▶ To cite this version:

Marie Anquetil, Victoria Dumont, Anne-Lise Marais, Nadege Roche-Labarbe, Sandrine Rossi. Sensory Prediction and Repetition Suppression in the tactile modality as early markers of executive attention development at preschool age. Flux Congress 2022, Sep 2022, Paris, France. 2022. hal-03770333

HAL Id: hal-03770333 https://normandie-univ.hal.science/hal-03770333

Submitted on 6 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License



INTRODUCTION

- The tactile modality, the first to develop, is the foundation of cognitive development.
- Sensory Prediction (SP) is a core mechanism sustained by our perceptions and allows the brain to anticipate future stimulations on the basis of previous sensory inputs (Friston, 2005). Repetition Suppression (RS) refers to the reduction of activity in the brain when a predicted stimulus is repeated or becomes irrelevant.
- Sensory Prediction in the tactile modality was suggested as an early precursor of cognitive development (Dumont et al., 2022).
- Predictive mechanisms may form the basis of executive attention development by sustaining self-regulation processes such as habituation or anticipation (Bubic et al., 2010).

The aim of this study is to investigate how tactile prediction and other tactile processing phenomena are related to executive attention development at preschool age.





• Frontal activity and behavioral performance at the executive attention task of the Child-ANT (all participants)



METHODS

Children between 48 and 60 months perform the following tasks in random order while 128-electrodes electroencephalography (Magstim EGI) is being acquired:

• an adapted version of the Child-Attention Network Task (ANT, a measure of executive attention) (Rueda et al., 2004)





CAYLAR

• a tactile oddball-omission paradigm (vibrotactile stimuli are presented on the forearm randomly interspersed with deviant and omitted stimuli).

Familiarization	Oddball-omission protocol	Familiarization	
		control	

• Executive attention as a function of group, and as a function of inhibition



- The N2 component is larger for both conditions in children with atypical development or born prematurely.
- The peak amplitude of N2 in the incongruent condition is positively related with inhibitory control performance at the Stroop task.
- Tactile sensory processing (all participants)







Children also perform the following behavioral tasks in random order :

- Movement Assessment Battery for Children 2nd edition (MABC-2)
- Stroop task adapted for non-reading children (Wright et al., 2003)
- Head-Toes-Knees-Shoulders (HTKS) task (McClelland et al., 2014)

Parents are also invited to complete 3 questionnaires : Dunn's Full Sensory Profile, BRIEF-P and ESSENCE 2-5.

We investigate the relationship between evoked potentials for executive attention (via the Child-ANT) and tactile sensory processing (via the oddball paradigm).

- For executive attention, we investigate the modulation of the N2 component, between congruent and incongruent conditions.
- For tactile sensory processing, we assess **repetition suppression**, *i.e.*, the decrease in somatosensory evoked potential (SEP) between the **familiarization** and the **control sequence** during the oddball omission-paradigm in the somatosensory cortex. We also assess potentials evoked by **deviant** stimulations.

We also investigate the relationship between brain evoked potentials and behavioral measurements.

CONCLUSION AND PERSPECTIVES

- During the Child-ANT, we observe differences of N2 amplitude for atypical profiles of children, associated with their performance on inhibitory control at the Stroop.
- We also demonstrate RS and deviance detection in the tactile modality.

- **Repetition suppression** : the amplitude of SEP at N140 decreases across the sequence.
- **Deviance detection** : the amplitude of SEP at N140 during **deviant** stimulations is higher compared to an equal number of standard stimulations.
- Tactile processing as a function of executive attention



- Tactile repetition suppression as a function of executive functions
- N140 amplitude for the control tactile stimuli increases with N2 amplitude for the incongruent condition of the ANT : the children who still respond strongly to tactile stimuli at the end of the sequence are also the children who have a higher response to cognitive conflict in the incongruent condition.
- This result suggests that deficit in RS and deficit in EA are linked.

We find an association between N140 in the tactile modality and N2 amplitude, but also with the executive composite score at the BRIEF-P, suggesting that sensory regulation mechanisms could be markers of executive attention development.

Such findings could help to promote new screening and remediation tools for children at-risk of atypical attentional development.

REFERENCES

- Anquetil, M., Roche-Labarbe, N. & Rossi, S. (under review). Tactile sensory processing as a precursor of executive attention : towards early detection impairments and of attention neurodevelopmental disorders. Wires Cognitive Science.
- Bubic, A., Von Cramon, D. Y., & Schubotz, R. I. (2010). Prediction, cognition and the brain. Frontiers in Human Neuroscience, 4.
- Dumont V., Giovannella M., Zuba D., Clouard R., Durduran T., Guillois B. and Roche-Labarbe N. (2022). Somatosensory prediction in the premature neonate brain. Developmental Cognitive Neuroscience (accepted article).
- Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1456), 815-836.
- McClelland, M. M., Cameron, C. E., Duncan, R., Bowles, R. P., Acock, A. C., Miao, A., & Pratt, M. E. (2014). Predictors of early growth in achievement: The head-toes-kneesacademic shoulders task. Frontiers in Psychology, 5.
- Rueda, M. R., Fan, J., McCandliss, B. D., Halparin, J. D., Gruber, D. B., Lercari, L. P., & Posner, M. I. (2004). Development of attentional in childhood. Neuropsychologia, 42(8), networks 1029-1040.
- Wright, I., Waterman, M., Prescott, H., & Murdoch-Eaton, D. (2003). A new Stroop-like measure of inhibitory function development: developmental trends. Journal of Child Typical *Psychology and Psychiatry*, 44(4), 561-575.



Repetition suppression of the tactile N140 is associated with BRIEF-P executive composite score : children who do not suppress tactile information across the sequence have higher BRIEF-P composite scores, reflecting impaired executive functioning.

Scan the flash code below to find this poster on PDF version



