



IQ testing in bacteria

Vic Norris, Camille Ripoll

► To cite this version:

Vic Norris, Camille Ripoll. IQ testing in bacteria. EMBO Workshop, Bacterial cell biophysics, growth, division, size and shape, Dec 2022, EIN GEDI, Israel. hal-03941000

HAL Id: hal-03941000

<https://normandie-univ.hal.science/hal-03941000>

Submitted on 16 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IQ testing in bacteria

Vic Norris and Camille Ripoll

Bacterial Communication and Anti-infection strategies, University of Rouen, France



Sensing the intensity of working of constituents, I , is important for many biological systems because once a constituent becomes limiting, the system can no longer grow exponentially. The solution is to duplicate this constituent – and perhaps divide – before it becomes limiting (1). Sensing the quantity, Q , of constituents in a system is also important to ensure the viability of the daughter subsystems produced by a division. At the level of the cell, DNA becomes limiting if it is not replicated. We report the results of the simulation of a turbidostat of bacteria written to address the question of how the decision to initiate DNA replication might be made by testing both I and Q .

If it is assumed that control over the replication of DNA and over cell division developed at some early stage in the evolution of cells, it might then be assumed that the nature of this control was independent of the action of sophisticated enzymes. I have argued that a powerful selection exists for cells (1) to replicate their DNA and their catalytic functions before they become limiting for growth and (2) to divide before they get too big (2). These actions require cells sensing the intensity, I , with which their constituents are working and sensing the quantity, Q , of their mass. To explore this hypothesis, we have simulated the growth of bacteria in a turbidostat. Each

bacterium has seven constituents: the two DNA strands, m miscellaneous material (lipids, metabolites, ions etc.), e enzymes (catalysing the production of enzymes), and ryb osomes (composite RNA polymerases and ribosomes) that can be either i inactive or f fast (expressing the strand encoding ribosomes) or s slow (expressing the strand encoding the enzymes). Each DNA strand is expressed by ribosomes that advance through a fixed number of intervals; each interval can be saturated with ribosomes (i.e., there is a maximum number of ribosomes per interval). Each newborn bacterium has a $NewBornMass = Ni + Nf + Ns + Ne + Nm$ (where N = number and the mass of each constituent is one unit). Growth rate and composition is calculated via iteration of equations based on kinetic constants kif (inactive to fast), kis (inactive to slow), and kem (enzyme catalysing miscellaneous from miscellaneous).

Intensity sensing is used to initiate DNA replication. Each bacterium divides at constant time after initiation of DNA replication. Strand segregation is semi-conservative (3, 4). Every time division occurs, a randomly chosen bacterium is discarded. The program allows the value of many variables to be calculated.

Virtual bacterium x1000 copies



1. Norris V, Amar P. 2012. Chromosome Replication in *Escherichia coli*: Life on the Scales. *Life* 2:286-312.
2. Norris V. 2015. Why do bacteria divide? *Front Microbiol* 6:322.
3. Norris V. 2019. Does the Semiconservative Nature of DNA Replication Facilitate Coherent Phenotypic Diversity? *J Bacteriol* 201.
4. Norris V, Ripoll C. 2021. Generation of Bacterial Diversity by Segregation of DNA Strands. *Frontiers in Microbiology* 12.