



HAL
open science

The invasive species *Rangia cuneata*: A new food source for herring gull (*Larus argentatus*)?

Jean-philippe Pezy, Ambre Pezy, Aurore Raoux

► To cite this version:

Jean-philippe Pezy, Ambre Pezy, Aurore Raoux. The invasive species *Rangia cuneata*: A new food source for herring gull (*Larus argentatus*)?. *Ecosphere*, 2022, 13 (4), 10.1002/ecs2.4058 . hal-03959821

HAL Id: hal-03959821

<https://normandie-univ.hal.science/hal-03959821>

Submitted on 28 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

ARTICLE

Coastal and Marine Ecology

The invasive species *Rangia cuneata*: A new food source for herring gull (*Larus argentatus*)?

 Jean-Philippe Pezy¹  | Ambre Pezy² | Aurore Raoux¹

¹Normandie Université, UNICAEN, UNIROUEN, Laboratoire Morphodynamique Continentale et Côtière, Caen, France

²Independent Researcher, Caen, France

Correspondence

Jean-Philippe Pezy
Email: jean-philippe.pezy@unicaen.fr

Funding information

Agence de l'Eau Seine-Normandie

Handling Editor: Hunter S. Lenihan

Abstract

Continued human population growth and the associated development of coasts lead to dramatic effects on marine coastal ecosystems altering food resources and habitats for several species and resulting in novel species interactions. The Bay of the Seine, located in the eastern part of the English Channel, is one of the most anthropized sea in the world. Here, we report an observation of the European herring gulls (*Larus argentatus*) feeding on the nonindigenous species (NIS) *Rangia cuneata* in the Bay of the Seine. Observations of otter prints (*Lutra lutra*) and excrements were also found near the bivalves, suggesting that this species also feed on the NIS bivalve. Our study suggests that the spread of NIS in coastal marine ecosystem is expected to drive changes in the foraging behavior of top predators.

KEYWORDS

brackish, ecosystem, English Channel, foraging innovation, non-indigenous species, prey–predator relations

INTRODUCTION

Intentional or unintentional species introductions by human activities to regions outside their native ranges have been ongoing for thousands of years (Carlton & Geller, 1993). However, these ecosystem invasions by nonindigenous species (NIS) are considered nowadays as one of the greatest threats to biodiversity worldwide (Mack et al., 2000; Vitousek et al., 1996). NIS introductions into coastal zones have been accelerated through the world in recent decades mainly due to increased human activities such as shipping, aquaculture, maritime traffic, and tourism (Boudouresque & Verlaque, 2002). This increase in NIS introduction occurring in the context of global climate change leads to modifications in the ecosystem structure and function (Vitousek et al., 1996) with potential repercussions on their ability

to provide goods and services to humans (Pimentel et al., 2000; Williamson & Fitter, 1996). In fact, NIS present a wide range of threats to native ecosystems and can be responsible for the decline in native species by competition and predation, and thus can lead to a loss of biodiversity (Sax & Gaines, 2008; Wilcove et al., 1998). Moreover, NIS can also cause direct economic impacts on human activity structure and aquaculture. Invasive species can also cause disappearance of the original communities. A famous example is the invasion of the ctenophore *Mnemiopsis leidyi* in the Black Sea, which led to a collapse of pelagic fisheries (GESAMP, 1997). Finally, it is worth to note that in the marine environment, these introductions are most of the time irreversible.

Apart from these negative effects, NIS may harmlessly be integrated into ecosystems and even

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Ecosphere* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

contribute to enrich the biodiversity (Gouletquer, 2016). Recently, studies have highlighted the potential positive role of NIS to regional species richness and conservation goals (Sax & Gaines, 2008). In fact, NIS can provide shelter (Wonham et al., 2005) and be responsible for an increase in habitat leading to an increase in abundance and species richness (Crooks, 2002). They can also provide a new source of food for native species (Bulleri et al. 2006). For instance, Cadée (2000, 2001, 2008) reported that the native European herring gulls (*Larus argentatus*) fed both on the NIS Pacific oysters (*Crassostrea gigas*) and on the NIS *Ensis directus* in the Wadden Sea. Finally, Holman et al. (2019) highlighted a novel herring gull's foraging behavior on the widespread sea squirt *Ciona intestinalis* from under floating pontoons in a recreational marina in Ireland.

Actually, a recent inventory of NIS along the Normandy coast concludes with a total of 152 NIS (Pezy, Baffreau, et al., 2021). Between Caen and Ouistreham was constructed between 1844 and 1857 the Canal de Caen à la Mer, a navigational channel of 14 km. Two NIS was reported in 1886 (*Amphibalanus amphitrite*) and in 1898 (*Mytilopsis leucophaeata*) (Pezy, Baffreau, et al., 2021). Recently, on 9 August 2017 for the first time *Rangia cuneata* was observed in the Canal de Caen à la Mer by fresh valves collected on the banks (Kerckhof et al., 2018). Since this observation, several dead shells were observed near the sea bottom with some barnacles fixed (personal observations). This brackish-water, suspension-feeding bivalve *R. cuneata* (G. B. Sowerby I, 1832) is native to the Gulf of Mexico (Carlton, 1992) and extended its range northward to the Chesapeake Bay (Hopkins & Andrews, 1970) and to the Hudson River (Carlton, 1992). In Europe, this species was observed for the first time in the Antwerp harbor, in Belgium in 2005 (Verween et al., 2006) and has since spread to estuaries in southern regions of the North Sea (Wiese et al., 2016). In its native range (Gulf of Mexico), *R. cuneata* is typically found in estuaries in salinities ranging between 5‰ and 15‰ (Swingle & Bland, 1974). *Rangia cuneata* is a bivalve with a remarkable salinity tolerance, being able to adapt to salinities varying from 0‰ to 33‰ (Cooper, 1981; Hopkins et al., 1974). The conditions of the Canal de Caen à la Mer are optimum to establish *R. cuneata* population (Faillietaz et al., 2020).

In this study, we report an observation of the European herring gulls feeding on the NIS *R. cuneata* in the Bay of the Seine (eastern part of the English Channel). Observations of otter prints and excrements were also found near the *Rangia* shells, suggesting that this species also feed on the NIS bivalve.

MATERIALS AND METHODS

Observations on herring gulls consuming *R. cuneata* at the Ouistreham harbor were made on August 2019. In fact, observations of herring gulls breaking the bivalve by shell dropped on a hard area were recorded at the Ouistreham harbor and along the bike path bordering the Canal de Caen à la Mer. *Rangia cuneata* shells that accumulated at the Ouistreham harbor along the bike path were collected between 12 and 23 August 2020 (Figures 1, 2a–c, 3, and 4). Only the right valves (the most abundant) were counted and measured. A total of 156 *R. cuneata* specimens were collected among them, 89 were used for length measurements (the 67 other valves were broken). The second site corresponds to the Herouville basin near the Caen City (Figures 1 and 2d). These observations were made along the banks and at the level of an old port structure on 25 August 2020. On this site, a lot of rubble (e.g., concrete block, wooden planks) are present. On a wooden board, located at the level of a path frequented by animals (traces of hair), paw prints were found. Empty shells of *R. cuneata* (29 individuals) have been found near the concrete banks of the Canal de Caen à la Mer. The two valves were still connected to each other by the ligament, but the commissure was chipped. Observations of otter excrements were also found during the same period near the Herouville basin (Figure 2e).

RESULTS

Of the 156 bivalves collected at Ouistreham from the pavement, 42.9% were broken and 57.1% were intact (two valves). The average length of 89 specimens of *R. cuneata* dropped and consumed by herring gulls was 35.3 mm (SD 4.6 mm); the range was 24–45 mm. During the different samplings, the herring gulls were observed to collect *R. cuneata* specimen by diving. Once the specimen collected, the herring gull flies to a high altitude and drop it onto a hard surface near the capture location (Figure 4).

The 29 shells of *R. cuneata* collected close to the Herouville basin were all intact. The average length of 29 specimens of *R. cuneata* was 37.0 mm (SD = 3.9 mm); the range was 31–47 mm. Observations of otter prints (*Lutra lutra*) and excrements were also found near the *Rangia* shells at the Herouville basin (Figure 2e). Further surveys carried out on Canal de Caen à la Mer did not result in the observation of other shells of *R. cuneata* without the presence of *L. lutra* excrements near them (Table 1).




FIGURE 1 Sampling sites of *Rangia cuneata* shell observation in Normandy, at Ouistreham harbor and Herouville basin in 2020



FIGURE 2 Individuals of *Rangia cuneata* from Ouistreham harbor (a, b, c) and individuals of *R. cuneata* from Herouville basin (d), and other excrements (e)




FIGURE 3 *Rangia cuneata* (G. B. Sowerby I, 1832) from Herouville basin, 0.80 m depth. (a) Left valve 77.4 mm long. (b) Right valve. (c) Same specimen, viewed from the dorsal region of the two valves. (d) View of the inside of the right valve. (e) View of the inside of the left valve. (f) View of the frontal region of the two valves



FIGURE 4 Individuals of *Rangia cuneata* broken by herring gulls from Ouistreham harbor

TABLE 1 Number and min, max, and mean size (\pm SD; in centimeters) of *Rangia cuneata* specimens recorded at Ouistreham and Herouville basin between 12 and 25 August 2020

<i>Rangia cuneata</i> specimens	Ouistreham	Herouville basin
Number	156	29
Min size	2.4	3.1
Max size	4.5	4.7
Mean size	3.5 \pm 0.5	3.7 \pm 0.4

DISCUSSION

The herring gull is considered as an apex predator and is known to be an opportunistic species with a wide trophic range. This species is also known to easily adapt to a new source of food (Burger & Gochfeld, 1996) such as anthropogenic food sources (rubbish dumps and bins) (Horton et al., 1983). In this paper, the NIS bivalve *R. cuneata* have been reported to be preyed upon by herring gulls in the Bay of the Seine, one of the most anthropized sea in the world (Halpern et al., 2008). Indeed, herring gull observations capturing the bivalve *R. cuneata* and taken it into the air and dropped from a certain height to break were recorded at the Ouistreham harbor along the bike path. This herring gull consumption of NIS can be explained in part by the fact that coastal habitats have been profoundly altered by human activities where most gull species commonly reside (Firth et al., 2016) and have led to dramatic effects on marine coastal ecosystems altering food resources and habitats for several species and resulting in novel species interactions. Thus, this new behavior in the Bay of the Seine seems to have developed as a consequence of these changes with, especially, the construction of the Canal de Caen la Mer. This herring gulls drop catch behavior on clams and more particularly on the bivalve *R. cuneata* has already been observed at a coastal estuary in Virginia, USA (Cristol et al., 2017; Gamble & Cristol, 2002). In addition, this smashing method is known for a long time (Oldham, 1930) and occurs worldwide (Cadée, 2001). Moreover, it is worth to note that observation of gulls feeding on NIS species has been already recorded in the literature (Cadée, 2000, 2001, 2008; Holman et al., 2019). For instance, Cadée (2000, 2001, 2008) reported herring gulls carrying the NIS Pacific oysters (*C. gigas*) or the NIS *E. directus* to high altitude and drop them onto rocks to crack them open and to feed them. Similarly, in 2018, Holman et al. (2019) observed a new behavior of herring gull feeding on *C. intestinalis* in a recreational marina in Ireland. In this study, the authors observed a gull diving down below the pontoons and emerging with a sea squirt of the non-native species *C. intestinalis*. All these studies highlighted that in these last decades, herring gulls have learned to prey on several NIS and thus that the spread of NIS in coastal

marine ecosystem is expected to drive changes in the foraging behavior of top predators. In addition that Faillettaz et al. (2020) found that the densities of *R. cuneata* (110.45 ± 86.08 individuals m^{-2}) in the Ouistreham marina largely dominating the benthos community and was similar to thus found in northern Europe. The authors also stated that a population of *R. cuneata* is now consistently established in Normandy (Ouistreham marina). Thus, these densities of *R. cuneata* could represent a large proportion of the herring gull diet (Faillettaz et al., 2020).

In Normandy, the otter lives in the catchment areas of the Orne, la Vire, l'Aure, and La Seulles, and so, it can be met in the Canal de Caen la Mer. This species is also known to be a potential native predator of NIS (Miranda et al., 2008). Observations of otter prints and excrements were found near the *Rangia* shells; this observation may suggest that the otter could feed on the NIS bivalve. As this species is a nocturnal species, its diet composition can be studied mainly by indirect methods such as the analyses of the fecal samples (Kruuk, 2006). In fact, several authors highlighted that in fecal samples, it is most common to find remains, which can be used for the identification of prey possibly to species level (Mason & Macdonald, 1986; Reynolds & Aebischer, 1991). However, as pointed out by Carss and Parkinson (1996), this method cannot be appropriate for detecting some food taxa, such as mollusks, or when only the prey soft parts are eaten. Thus, nocturnal observations with photographic evidence are needed in order to analyze the potential proportion of *Rangia* consumed by this species. In addition, it will be interesting to repeat this kind of study after some years in order to analyze whether the herring gulls improve their foraging method as suggested by Barash et al. (1975).

ACKNOWLEDGMENTS

This research was funded by the Agence de l'Eau Seine-Normandie in the CHABIMANOR project.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data (Pezy, Pezy, & Raoux, 2021) are available from Dryad: <https://doi.org/10.6084/m9.figshare.17152145>.

ORCID

Jean-Philippe Pezy  <https://orcid.org/0000-0002-8293-5090>

REFERENCES

- Barash, D. P., P. Donovan, and R. Myrick. 1975. "Clam Dropping Behavior of the Glaucous-Winged Gull (*Larus glaucescens*)."
Wilson Bull 87: 60–4.

- Boudouresque, C. F., and M. Verlaque. 2002. "Assessing Scale and Impact of Ship-Transported Alien Macrophytes in the Mediterranean Sea." In *Alien Marine Organisms Introduced by Ships in the Mediterranean and Black Seas*. CIESM Workshop Monographs 20, edited by F. Briand, 53–61. Monaco: CIESM.
- Bulleri, F., L. Airoidi, G. M. Branca, and M. Abbiati. 2006. "Positive Effects of the Introduced Green Alga, *Codium fragile* ssp. *tomentosoides*, on Recruitment and Survival of Mussels." *Marine Biology* 148: 1213–20.
- Burger, J., and M. Gochfeld. 1996. "Family Laridae." In *Handbook of the Birds of the World*, Vol 3, edited by J. Del Hoyo, A. Elliot, and J. Sargatal, 572–623. Barcelona: Lynx Edicions.
- Cadée, G. C. 2000. "Herring Gulls Feeding on a Recent Invader in the Wadden Sea, *Ensis directus*." In *The Evolutionary Biology of the Bivalvia*. Special Publication 177, edited by E. M. Haroer, J. D. Taylor, and J. A. Crame, 459–64. London: Geological Society.
- Cadée, G. C. 2001. "Herring Gulls Learn to Feed on a Recent Invader in the Dutch Wadden Sea, the Pacific Oyster *Crassostrea gigas*." *Basteria* 65: 33–42.
- Cadée, G. C. 2008. "Herring Gulls Feeding Again on Pacific Oysters *Crassostrea gigas* in the Dutch Wadden Sea near Texel." *Basteria* 72: 33–6.
- Carlton, J. T., and J. B. Geller. 1993. "Ecological Roulette: The Global Transport of Non-Indigenous Marine Organisms." *Science* 261: 78–82.
- Carlton, J. T. 1992. "Introduced Marine and Estuarine Mollusks of North America: An End-of-the-20th-Century Perspective." *Journal of Shellfish Research* 11(2): 489–505.
- Carss, D. N. and S. G. Parkinson. 1996. "Errors Associated with Otter *Lutra lutra* Faecal Analysis. I. Assessing General Diet from Spraints." *Journal of Zoology* 238: 301–17.
- Cooper, R. B. 1981. "Salinity Tolerance of *Rangia Cuneata* (Pelecypoda:Mactridae) in Relation to its Estuarine Environment: A review." *Walkerana* 1: 19–31.
- Cristol, D. A., J. G. Akst, M. K. Curatola, E. G. Dunlavy, K. A. Fisk, and K. E. Moody. 2017. "Age-Related Differences in Foraging Ability among Clam-Dropping Herring Gulls (*Larus argentatus*)." *The Wilson Journal of Ornithology* 129: 301–10.
- Crooks, J. 2002. "Characterizing Ecosystem-Level Consequences of Biological Invasions: The Role of Ecosystem Engineers." *Oikos* 97: 153–66.
- Faillietaz, R., C. Roger, M. Mathieu, J. P. Robin, and K. Costil. 2020. "Establishment and Population Features of the Non-Native Atlantic *Rangia*, *Rangia cuneata* (Mollusca: Bivalvia), in Northwestern France." *Aquatic Invasions* 15: 367–81.
- Firth, L. B., A. M. Knights, D. Bridger, A. Evans, N. Mieskowska, P. J. Moore, N. E. O'Connor, E. V. Sheehan, R. C. Thompson, and S. J. Hawkins. 2016. "Ocean Sprawl: Challenges and Opportunities for Biodiversity Management in a Changing World." In *Oceanography and Marine Biology*, edited by R. N. Hughes, D. J. Hughes, I. P. Smith, and A. C. Dale, 201–78. Boca Raton, FL: CRC Press.
- Gamble, J. R., and D. A. Cristol. 2002. "Drop-Catch Behaviour Is Play in Herring Gulls, *Larus argentus*." *Animal Behaviour* 63: 339–45.
- GESAMP (IMO/FAO/UNESCO-IOC/WMO/WHO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection) 1997. "Opportunistic settlers and the problem of the ctenophore *Mnemiopsis leidyi* invasion in the Black Sea." Report Number 58.
- Gouletquer, P. 2016. *Guide des Organismes Exotiques Marins*. Berlin: Editions Belin.
- Halpern, B. S., S. Walbridge, K. A. Selkoe, C. V. Kappel, F. Micheli, C. D'Agrosa, and R. Watson. 2008. "A Global Map of Human Impact on Marine Ecosystems." *Science* 319: 948–52.
- Holman, L. E., M. Rius, and T. M. Blackburn. 2019. "Observations of a Novel Predatory Gull Behavior on an Invasive Ascidian: A New Consequence of Coastal Urban Sprawl?" *Ecosphere* 10: e02636.
- Hopkins, S. H., and J. D. Andrews. 1970. "*Rangia cuneata* on the East Coast: Thousand Mile Range Extension, or Resurgence." *Science* 167: 868–9.
- Hopkins, S. H., J. W. Anderson, and K. Horvath. 1974. "Biology of the Clam *Rangia Cuneata*: What We Know and What it Means." *Proceedings of the National Shellfisheries Association* 64.
- Horton, N., T. Brough, and J. B. A. Rochard. 1983. "The Importance of Refuse Tips to Gulls Wintering in an Inland Area of Southeast England." *Journal of Applied Ecology* 20: 751–65.
- Kerckhof, F., M. Devleeschouwer, and N. Hamers. 2018. "De Amerikaanse Brakwaterstrandschelp *Rangia cuneata* (G. B. Sowerby, 1832) Aangetroffen in Frankrijk." *De Strandvlo* 37: 141–5.
- Kruuk, H. 2006. *Otters. Ecology, Behaviour and Conservation*. Oxford: Oxford University Press.
- Mack, R. N., D. Simberloff, W. M. Lonsdale, H. Evans, M. Clout, and F. A. Bazzaz. 2000. "Biotic Invasions: Causes, Epidemiology, Global Consequences, and Control." *Ecological Applications* 10: 689–710.
- Mason, C. F., and S. M. Macdonald. 1986. *Otters: Ecology and Conservation*. Cambridge: Cambridge University Press.
- Miranda, R., G. H. Copp, J. Williams, K. Beyer, and R. E. Gozlan. 2008. "Do Eurasian Otters *Lutra lutra* (L.) in the Somerset Levels Prey Preferentially on Non-Native Fish Species?" *Fundamental and Applied Limnology* 172: 339–47.
- Oldham, C. 1930. "Locomotive Habit of *Dreissena polymorpha*." *J. Conchol.* 19: 25–26.
- Pezy, J. P., A. Baffreau, A. M. Rusig, I. Mussio, A. Raoux, and J. C. Dauvin. 2021. "Non-Indigenous Species in Marine and Brackish Waters along the Normandy Coast in the English Channel." *BioInvasions Records* 10(4): 755–74.
- Pezy, J.-P., A. Pezy, and A. Raoux. 2021. "*Rangia cuneata* size date.docx." Figshare. Dataset. <https://doi.org/10.6084/m9.figshare.17152145.v1>.
- Pimentel, D., L. Lach, R. Zuniga, and D. Morrison. 2000. "Environmental and Economic Costs of Non-Indigenous Species in the United States." *Bioscience* 50: 53–65.
- Reynolds, J. C., and N. J. Aebischer. 1991. "Comparison and Quantification of Carnivore Diet by Faecal Analysis: A Critique, with Recommendations, Based on a Study of the Fox *Vulpes vulpes*." *Mammal Review* 21: 97–122.
- Sax, D. F., and S. D. Gaines. 2008. "Species Invasions and Extinction: The Future of Native Biodiversity on Islands." *Proceedings of the National Academy of Sciences* 105: 11490–7.
- Swingle, H. A., and D. G. Bland. 1974. "Distribution of the Estuarine Clam *Rangia cuneata* in Coastal Waters of Alabama." *Alabama Marine Resources Bulletin* 10: 9–16.
- Verween, A., F. Kerckhof, M. Vincx, and S. Degraer. 2006. "First European Record of the Invasive Brackish Water Clam *Rangia cuneata* (G.B. Sowerby I, 1831) (Mollusca: Bivalvia)." *Aquatic Invasions* 1: 198–203.

- Vitousek, P. M., C. M. D'Antonio, L. L. Loope, and R. Westbrooks. 1996. "Biological Invasions as Global Environmental Change." *American Journal of Science* 84: 468–78.
- Wiese, L., O. Niehus, B. Faass, and V. Wiese. 2016. "Ein weiteres Vorkommen von *Rangia cuneata* in Deutschland (Bivalvia: Mactridae)." *Schriften zur Malakozoologie*. 29: 53–60.
- Wilcove, D. S., D. Rothstein, J. Dubow, A. Phillips, and E. Losos. 1998. "Quantifying Threats to Imperiled Species in the United States." *Bioscience* 48: 607–15.
- Williamson, M., and A. Fitter. 1996. "The Varying Success of Invaders." *Ecology* 77: 1661–6.
- Wonham, M. J., M. O'Connor, and C. D. G. Harley. 2005. "Positive Effects of a Dominant Invader on Introduced and

Native Mudflat Species." *Marine Ecology Progress Series* 289: 109–16.

How to cite this article: Pezy, Jean-Philippe, Ambre Pezy, and Aurore Raoux. 2022. "The Invasive Species *Rangia Cuneata*: A New Food Source for Herring Gull (*Larus Argentatus*)?." *Ecosphere* 13(4): e4058. <https://doi.org/10.1002/ecs2.4058>