

A national-scale field study to evaluate potential N 2 O emissions: Community composition predicts soil potential denitrification in Irish grasslands

C Deveautour, Murilo Veloso, A M Duff, D Wall, R Carolan, K G Richards, O Flaherty, Fiona Brennan, Paula Rojas-Pinzon, James Rambaud

▶ To cite this version:

C Deveautour, Murilo Veloso, A M Duff, D Wall, R Carolan, et al.. A national-scale field study to evaluate potential N 2 O emissions: Community composition predicts soil potential denitrification in Irish grasslands. ISME18, Aug 2022, Lausanne, Switzerland. hal-04168225

HAL Id: hal-04168225 https://normandie-univ.hal.science/hal-04168225

Submitted on 17 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.












Deveautour, C.^{1,2,3}, Rojas-Pinzon, P.A.^{1,2,6}, Veloso, M.³, Rambaud, J.¹, Duff, A. M.¹, Wall, D.¹, Carolan, R.⁴, Philippot, L.⁵, Richards, K.G.¹, O'Flaherty, V.², Brennan, F¹.


¹ Environmental Research Centre, Teagasc, Johnstown Castle, Co. Wexford, Ireland ² School of Natural Sciences and Ryan Institute, National University of Ireland, University Road, Galway, Ireland H91 TK33 ³ Institut Polytechnique UniLaSalle, Unité AGHYLE, Campus Rouen, 76130, Mont-Saint-Aignan, Normandie, France. ⁴Agri-Food and Biosciences Institute (AFBI), Newforge Lane, Belfast BT9 5PX, Northern Ireland ⁵ University Bourgogne Franche-Comte, INRAE, AgroSup Dijon, Department of Agroecology, Dijon, France ⁶University of Vienna, Centre for Microbiology and Environmental Systems Science, Division of Microbial Ecology. Vienna, Austria.



A national-scale field study to evaluate potential N_2 0 emissions:



136 samples collected in 32 sites in Ireland, together with soil physicochemical properties, climate, functional genes abundance and soil communities composition. Soil potential denitrification was assessed under optimal conditions using incubations.








Soils' potential denitrification varied greatly between sites and even at the same sites, with soil P, N and pH being the most predictive. In most of the soils, emissions were in the form of N_2O rather than N_2 .

Among biotic factors, prokaryotic community composition was more predictive than the denitrifiers community abundance:

We evaluated the prokaryotic (16S) and fungal (ITS) community composition using Illumina sequencing and determined the abundance of genes involved in denitrification (nirK, nirS, nosZ clade I and II) with qPCR.







The prokaryotic community composition better predicted the potential denitrification rate. The relative abundance of **Actinobacteriota** and **Crenarchaeota** was positively correlated to potential denitrification (R = 0.35 & 0.41, respectively).

Long story short...

- Soil phosphorus was as important predicting PD as was pH and total nitrogen.
- The prokaryotic community was more predictive of soil PD than the fungal community.
- Soil community structure, rather than denitrifiers' community abundance, was an important predictor.
- Actinobacteriota and Crenarchaeota were correlated to complete denitrification.

Our open access paper



