



**HAL**  
open science

## Methods for studying the bacterial urinary microbiome of non menopausal women

Camille Villenave, Sandrine Dahyot, Marie Leoz, Maxime Grand, Chervin Hassel, Martine Pestel-Caron

► **To cite this version:**

Camille Villenave, Sandrine Dahyot, Marie Leoz, Maxime Grand, Chervin Hassel, et al.. Methods for studying the bacterial urinary microbiome of non menopausal women. ECCMID, Apr 2023, Copenhagen, Denmark. hal-04184256

**HAL Id: hal-04184256**

**<https://normandie-univ.hal.science/hal-04184256>**

Submitted on 21 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P-1814

# Methods for studying the bacterial urinary microbiome of non-menopausal women

C. Villenave<sup>1</sup>, S. Dahyot<sup>2</sup>, M. Leoz<sup>1</sup>, M. Grand<sup>1</sup>, C. Hassel<sup>1</sup>, M. Pestel-Caron<sup>2</sup>

<sup>1</sup>Univ Rouen Normandie, UNICAEN, INSERM, DYNAMICURE UMR 1311 F-76000 Rouen, France; <sup>2</sup>Univ Rouen Normandie, UNICAEN, DYNAMICURE UMR 1311, CHU Rouen, Department of bacteriology, F-76000 Rouen, France  
[camille.villenave@univ-rouen.fr](mailto:camille.villenave@univ-rouen.fr); tel+33 2.35.14.82.99


## Background

Next generation sequencing technologies (NGS) as well as the development of extended cultures (EC) have allowed the recent discovery of an **Urinary Microbiota (UM)**<sup>1</sup>. Nevertheless, there are still many unknowns regarding the UM associated with healthy subjects or **urinary tract infections (UTIs)**.

## Objectives


- Locally implement **EC** and **NGS** (Illumina and Nanopore) to describe **female bacterial UM**
- Identify **urotype** (dominant bacterial taxon) for each urine specimen
- Compare bacterial UM of patients with **UTIs** with that of **healthy volunteers (HV)**

## Methods




## Results


### Analysis of UM by culture



### Analysis of UM by NGS



### UM of UTI vs HV



## Conclusions

- EC** improves detection of uropathogens compared to SC procedure<sup>1</sup>. Both **EC** and **NGS** results showed a **difference in the composition of the UM** between **HV** and **UTI** patients. This study highlights a high occurrence of **Lactobacillus** in the female UM, as already reported<sup>2</sup>.
- Nevertheless, these results must to be confirmed with a larger sample collection, and the analyses of the NGS data need to be further investigated, particularly by using other databases than those suggested by the manufacturers.

1. Price, Travis K., Tanaka Dune, Evann E. Hilt, Krystal J. Thomas-White, Stephanie Kliethermes, Cynthia Brincat, Linda Brubaker, Alan J. Wolfe, Elizabeth R. Mueller, and Paul C. Schreckenberger. "The Clinical Urine Culture: Enhanced Techniques Improve Detection of Clinically Relevant Microorganisms." *Journal of Clinical Microbiology* 54, no. 5 (May 2016): 1216–22.  
 2. Ugarcina Perovic, Svetlana, Magdalena Ksiezarek, Joana Rocha, Elisabete Alves Cappelli, Márcia Sousa, Teresa Gonçalves Ribeiro, Filipa Grosso, and Luísa Peixe. "Urinary Microbiome of Reproductive-Age Asymptomatic European Women." *Theis. Microbiology Spectrum* 10, no. 6 (December 21, 2022): e01308-22.