



Short-Course Versus Prolonged-Course Antimicrobial Therapy in Adults With Catheter-Related Septic Thrombosis: a Propensity-Weighted Retrospective Study

Marine Stoldick, Margot Vannier, Mikael Verdalles-Cazes, Manuel Etienne,
Kévin Alexandre

► To cite this version:

Marine Stoldick, Margot Vannier, Mikael Verdalles-Cazes, Manuel Etienne, Kévin Alexandre. Short-Course Versus Prolonged-Course Antimicrobial Therapy in Adults With Catheter-Related Septic Thrombosis: a Propensity-Weighted Retrospective Study. ECCMID, Apr 2023, Copenhagen, Denmark. hal-04184293

HAL Id: hal-04184293

<https://normandie-univ.hal.science/hal-04184293>

Submitted on 21 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short-Course Versus Prolonged-Course Antimicrobial Therapy in Adults With Catheter-Related Septic Thrombosis: a Propensity-Weighted Retrospective Study

Marine STOLDICK¹; Margot VANNIER²; Mikael VERDALLES-CAZES³; Manuel ETIENNE⁴; Kévin ALEXANDRE⁴

¹CHU Rouen, Department of Infectious diseases, F-76000 Rouen, France ²CHU Rouen, Department of Biostatistics, F-76000 Rouen, France ³ CHU Rouen, Department of Radiology, F-76000 Rouen, France ⁴Univ Rouen Normandie, Inserm UMR 1311 DYNAMICURE, CHU Rouen, Department of Infectious diseases, F-76000 Rouen, France

About septic thrombosis ...

- 30% of hospitalized patients are catheterized
- 20% mortality per catheter-related bacteremia
- 17 000 €

Why reduce antimicrobial therapy duration for catheter-related septic thrombosis?

LESS

- Adverse events
- Clostridium difficile*
- Antimicrobial resistance
- Hospitalization duration

≤ 3 weeks VS > 3 weeks

of antimicrobial therapy



MORE

- Relapse ?
- Mortality ?

Methods

DESIGN


- Retrospective
- Monocentric
- Between 2015 and 2020
- Propensity score

INCLUSION

- Adults
- Catheter-related deep venous septic thrombosis
- Upper limb

EXCLUSION


- Missing data
- Secondary localization
- Prolonged treatment for a different infection



Results


63 years old
60% male

58% solid tumor
4% oncohematology




Catheters:

- 32% long-term
- 35% short-term
- 33% others



30% chemotherapy
18% parenteral nutrition

36% CoNS
21% *S. aureus*
10% GNB
7% NFGNB
7% *Candida* spp.



Antimicrobial therapy duration	≤3 weeks (n = 75)		>3 weeks (n = 107)		wHR (95% CI)
	n	%	n	%	
30-day all-cause mortality or relapse	13	17.3%	21	19.6%	0.66 (0.33-1.36)
90-day mortality	19	25.3%	24	22.4%	0.73 (0.41-1.30)
90-day relapse	2	2.7%	4	3.7%	0.91 (0.17-4.95)

Discussion

- A 3-week antimicrobial therapy seem as safe as a longer-course antimicrobial therapy to treat catheter-related septic thromboses
- Caution should be exercised with *Candida* septic thromboses and oncohematologic patients