



HAL
open science

Thiosulfate sulfurtransferase deficiency promotes oxidative distress and aberrant NRF2 function in the brain

Yang Luo, Laurent Chatre, Shaden Melhem, Zayana M Al-Dahmani, Natalie Z.M. Homer, Anneke Miedema, Leo E Deelman, Matthew R Groves, Martin Feelisch, Nicholas M Morton, et al.

► **To cite this version:**

Yang Luo, Laurent Chatre, Shaden Melhem, Zayana M Al-Dahmani, Natalie Z.M. Homer, et al.. Thiosulfate sulfurtransferase deficiency promotes oxidative distress and aberrant NRF2 function in the brain. *Redox Biology*, 2023, 68, pp.102965. 10.1016/j.redox.2023.102965 . hal-04294976

HAL Id: hal-04294976

<https://normandie-univ.hal.science/hal-04294976>

Submitted on 20 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Journal Pre-proof

Thiosulfate sulfurtransferase deficiency promotes oxidative distress and aberrant NRF2 function in the brain

Yang Luo, Laurent Chatre, Shaden Melhem, Zayana M. Al-Dahmani, Natalie Z.M. Homer, Anneke Miedema, Leo E. Deelman, Matthew R. Groves, Martin Feelisch, Nicholas M. Morton, Amalia Dolga, Harry van Goor

PII: S2213-2317(23)00366-X

DOI: <https://doi.org/10.1016/j.redox.2023.102965>

Reference: REDOX 102965

To appear in: *Redox Biology*

Received Date: 14 October 2023


Revised Date: 6 November 2023

Accepted Date: 14 November 2023

Please cite this article as: Y. Luo, L. Chatre, S. Melhem, Z.M. Al-Dahmani, N.Z.M. Homer, A. Miedema, L.E. Deelman, M.R. Groves, M. Feelisch, N.M. Morton, A. Dolga, H. van Goor, Thiosulfate sulfurtransferase deficiency promotes oxidative distress and aberrant NRF2 function in the brain, *Redox Biology* (2023), doi: <https://doi.org/10.1016/j.redox.2023.102965>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier B.V.



Thiosulfate Sulfurtransferase Deficiency Promotes Oxidative Distress and Aberrant NRF2 Function in the Brain

Yang Luo^{1,2}, Laurent Chatre³, Shaden Melhem⁴, Zayana M. Al-Dahmani⁵, Natalie Z.M. Homer⁶, Anneke Miedema², Leo E. Deelman⁷, Matthew R. Groves⁵, Martin Feelisch⁸, Nicholas M. Morton^{4,9}, Amalia Dolga¹, Harry van Goor^{2*}

1. University of Groningen, Dept. of Molecular Pharmacology, Groningen Research Institute of Pharmacy, Faculty of Science and Engineering, Groningen, the Netherlands

2. University Medical Center Groningen, Dept. of Pathology and Medical Biology, Groningen, the Netherlands

3. Université de Caen Normandie, CNRS, Normandie Univ, ISTCT UMR6030, GIP Cyceron, F-14000 Caen, France

4. Centre for Cardiovascular Science, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, United Kingdom

5. University of Groningen, Dept. of Pharmacy, Drug Design, Groningen, the Netherlands


6. Mass Spectrometry Core, Edinburgh Clinical Research Facility, University of Edinburgh/BHF Centre for Cardiovascular Sciences, Queen's Medical Research Institute, Edinburgh, University of Edinburgh

7. University of Groningen, University Medical Center Groningen, Dept. of Clinical Pharmacy and Pharmacology, Groningen, the Netherlands

8. Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, United Kingdom

9. Centre for Systems Health and Integrated Metabolic Research, School of Science and Technology, Nottingham Trent University, Nottingham, United Kingdom

Graphical abstract:



Created with Biorender.com

Abstract

Thiosulfate sulfurtransferase (TST, EC 2.8.1.1) was discovered as an enzyme that detoxifies cyanide by conversion to thiocyanate (rhodanide) using thiosulfate as substrate; this rhodanese activity was subsequently identified to be almost exclusively located in mitochondria. More recently, the emphasis regarding its function has shifted to hydrogen sulfide metabolism, antioxidant defense, and mitochondrial function in the context of protective biological processes against oxidative distress. While TST has been described to play an important role in liver and colon, its function in the brain remains obscure. In the present study, we therefore sought to address its potential involvement in maintaining cerebral redox balance in a murine model of global TST deficiency (*Tst*^{-/-} mice), primarily focusing on characterizing the biochemical phenotype of TST loss in relation to neuronal activity and sensitivity to oxidative stress under basal conditions. Here, we show that TST deficiency is associated with a perturbation of the *reactive species interactome* in the brain cortex secondary to altered ROS and RSS (specifically, polysulfide) generation as well as mitochondrial OXPHOS remodeling. These changes were accompanied by aberrant Nrf2-Keap1 expression and thiol-dependent antioxidant function. Upon challenging mice with the redox-active herbicide paraquat (25mg/kg i.p. for 24hrs), *Tst*^{-/-} mice displayed a lower antioxidant capacity compared to wildtype controls (C57BL/6J mice). These results provide a first glimpse into the molecular and metabolic changes of TST deficiency in the brain and suggest that pathophysiological conditions associated with aberrant TST expression and/or activity renders neurons more susceptible to oxidative stress-related malfunction.

Keywords: Thiosulfate sulfurtransferase (TST), Reactive species, Antioxidant, Oxidative distress, Nrf2 signaling...

Abbreviations

TST	Thiosulfate sulfurtransferase
CBS	Cystathionine beta synthase
CSE	Cystathionine gamma lyase
MPST	3-mercaptopyruvate sulfurtransferase
H ₂ S	Hydrogen sulfide
SQOR	Sulfide quinone oxidoreductase
PDO	Persulfide dioxygenase
DHLA	Dihydrolipoic acid
GSH	Glutathione
BMI	Body mass index
HFD	High-fat diet
SOD	Superoxide dismutase
ROS	Reactive oxygen species
2-PTS	2-propenyl thiosulfate
DSS	Dextran sulfate sodium
RNS	Reactive nitrogen species
RSS	Reactive sulfur species
RSI	Reactive species interactome
GSSG	Glutathione disulfide
GPX	Glutathione peroxidase
GR	Glutathione reductase
FCCP	Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone
TMPD	N,N,N',N'-tetramethyl-p-phenylenediamine
PRDX	Peroxiredoxin
NOS	Nitric oxide synthase
CAT	Catalase
NRF2	Nuclear factor erythroid 2-related factor 2
KEAP1	Kelch-like ECH-associated protein 1
HMOX1	Heme Oxygenase 1
TXN	Thioredoxin
NQO1	NAD(P)H dehydrogenase [quinone] 1
GCLC	Glutamate cysteine ligase, catalytic subunit
GCLM	Glutamate cysteine ligase, modifier subunit
GSTM1	Glutathione S-transferase Mu

Introduction

The enzyme thiosulfate sulfurtransferase (TST, EC 2.8.1.1) was first identified in 1933 as a cyanide (CN⁻) detoxification enzyme that utilizes thiosulfate (S₂O₃²⁻) as sulfur donor to form thiocyanate (SCN⁻; also known as rhodanide).^{1,2} The use of thiosulfate as an antidote to cyanide poisoning relies on this enzymatic reaction. Another known rhodanese enzyme, 3-mercaptopyruvate sulfurtransferase (MPST, EC 2.8.1.2), shares a high sequence homology with TST, and both enzymes are thought to be evolutionarily closely related.^{3,4} TST is abundantly expressed in various tissues and plays a vital role in three important biological processes: the metabolism of hydrogen sulfide (H₂S; hereinafter referred to as sulfide), modulation of mitochondrial respiratory chain activity, and involvement in the thiol-dependent antioxidant system.⁵⁻⁸

In the context of sulfide metabolism, TST oxidizes the endogenous gasotransmitter H₂S in a similar fashion as sulfide quinone oxidoreductase (SQOR) and persulfide dioxygenase (ETHE1/PDO).⁹ TST also has the capacity to form H₂S from thiosulfate by utilizing dihydrolipoic acid (DHLA), the reduced form of α -lipoic acid.^{10,11} Balancing both functions, H₂S oxidation and production, it seems that the primary function of TST is in the detoxification of H₂S and less so in the production of H₂S.¹² TST is almost exclusively expressed in mitochondria, the organelle that synthesizes ATP by oxidative phosphorylation (OXPHOS) through stepwise reduction of oxygen in the Electron Transport Chain (ETC) and the generated proton gradient using five enzyme complexes (Complex I-V).^{13,14} TST activity appears to modulate reduced nicotinamide adenine dinucleotide (NADH) dehydrogenase (Complex I) and other mitochondrial complexes including succinate dehydrogenase (Complex II) via interaction with their iron-sulfur clusters.^{6,7} In addition to this interaction between mitochondrial activity and TST, phosphorylation of TST may modulate its activity and thereby affect the stability of sulfur in the iron-sulfur centers of Complexes III and IV, further impacting electron transport and ATP production.¹⁵ In mammalian cells, TST is linked to two thiol-dependent antioxidant systems, the thioredoxin system and the GSH system.¹⁵ TST exerts anti-oxidative properties via donating sulfur for these two systems and an increase of TST activity can (in)directly activate their reactive oxygen species (ROS) scavenging function.^{12,16}

Oxidative stress has classically been defined as an imbalance in the generation of pro-oxidant and antioxidant species in favor of the former.¹⁷ A series of experimental observations promoted a redefinition of oxidative stress as a condition that is associated with alterations in redox signaling and control, and an updated interpretation of the original concept distinguishes physiological oxidative stress (or 'oxidative eustress') from excessive and deleterious oxidative stress (coined 'oxidative distress').¹⁸ Besides ROS, some other types of short-lived molecules such as reactive nitrogen (RNS) and reactive sulfur species (RSS) contribute to cellular signaling processes and merit consideration specifically in the context of sensing and adaptation to changes in environmental and/or metabolic conditions. These considerations are part of the stress signaling paradigm and have recently been conceptualized in the form of the Reactive Species Interactome (RSI) framework.^{19,20}

The transcription factor Nrf2 (nuclear factor (erythroid-derived 2)-like 2) is a master regulator of cellular homeostasis that controls the expression of genes related to redox homeostasis.²¹ The

primary mechanism of NRF2 regulation is at the protein level. The majority of research has focused on the involvement of the electrophile and redox sensor Kelch-like ECH-associated protein 1 (KEAP1) in regulating NRF2 protein levels in response to metabolic demands.²² The association between KEAP1 and NRF2 is disrupted by electrophilic alteration or oxidation of cysteine thiols in KEAP1, enabling cells/organisms to respond to environmental stress. Following dissociation from KEAP1, NRF2 escapes destruction and targets antioxidant response element (ARE) genes, which leads to an enhancement of antioxidant capacity.²³ Moreover, various studies linked the TST-related antioxidant system to nuclear factor erythroid 2-related factor 2 (Nrf2) signaling due to its transcriptional activation of GSH-related enzymes, one of the first lines of defense against oxidative stress.^{24–27} Here, H₂S mediates direct persulfidation of KEAP1 (Kelch-like ECH-associated protein 1) and thereby contributes to sulfide-mediated NRF2 regulation.²⁸

Several studies have demonstrated that TST provides an essential protective function against disease.¹⁵ Studies investigating the mechanistic effects of TST, which were identified in polygenic lean mouse adipocytes, have indicated its potential as a predictor for diabetes.²⁶ In an Icelandic cohort of almost 700 people, an inverse correlation was observed between *TST* mRNA levels in subcutaneous adipose tissue and body mass index (BMI).²⁹ Upregulation of adipose TST was detected in lean mice challenged with a high-fat diet (HFD), further supporting its likely protective effects.²⁹ Additionally, mice with transgenic *Tst* overexpression in mature adipocytes, showed resistance against HFD-induced obesity and exhibited higher protein expression of mitochondrial superoxide dismutase 2 (SOD2) and higher mRNA levels of peroxiredoxin 3 (*Prdx3*) compared to control mice, while cytosolic superoxide dismutase 1 (SOD1) remained unchanged.²⁹ These findings provide evidence of an interaction between TST and mitochondrial ROS. Consistent with a function of TST for ROS removal, knockdown of *Tst* in the preadipocyte cell line 3T3-L1 generated higher levels of mitochondrial ROS in the presence of oxidative stress stimuli.²⁹ Furthermore, ROS-related adiponectin release from 3T3-L1 adipocytes was decreased in cells treated with the TST inhibitor, 2-PTS (2-propenyl thiosulfate), while adiponectin was increased following addition of the TST substrate thiosulfate.²⁹ TST enables H₂S detoxification in colon tissue, and has reduced expression in colon mucosa of patients with ulcerative colitis and colon cancer.^{30–32} During the development of dextran sulfate sodium (DSS) induced colitis in mice, the expression of *Tst* mRNA, protein and activity significantly decrease.³³

Taken together, these findings indicate that TST has an important function in maintaining the redox balance in colon and adipocytes, but there are no studies elucidating the importance of TST in redox signaling within the brain. A clinical report of rhodanese (i.e. TST) deficiency in a rare neurological and mitochondrial human disorder known as Leber's Hereditary Optic Atrophy (LHON)³⁴ prompted us to investigate the underlying molecular and metabolic changes of TST deficiency in the brain. From physiology to pathology, the brain consumes about 20% of the whole-body supply of oxygen and energy²⁶. The cerebral cortex, with its intense neuronal activity and high metabolic activity, has a particularly high susceptibility to perturbation by oxidative stress.^{35–37} In Alzheimer's Disease, increased oxidative stress is associated with the progression of the neurodegenerative pathology along with amyloid and hyperphosphorylated tau.^{38,39} This is due to high mitochondrial generation of ROS, including superoxide anion (O₂[•]

) and hydrogen peroxide (H₂O₂).^{19,40} Hence, we hypothesized that TST plays a vital role in enabling an appropriate redox balance in the brain, and that global gene silencing of TST in mice would provide novel insights into the central role of TST in neurological antioxidant signaling and tissue protection.

Journal Pre-proof

Methods

Experimental animals

All the experiments were performed according to international guidelines set out by the ethical committees of the University of Edinburgh and within the framework of the Animals (Scientific Procedures) Act (1986) of the United Kingdom Home Office. Animals were maintained in standard housing conditions with a 12-hour light and 12-hour dark cycle (7 a.m. to 7 p.m.) and *ad libitum* access to rodent chow. All studies used female mice housed in cages of 3-6 individual littermates. The female mice used for the study originated from C57BL/6N *Tst*^{-/-} mice²⁹ backcrossed onto a C57BL/6J genetic background for more than 10 generations. Pieces of brain cortex were snap frozen and stored at -80 °C until use.

Induction of oxidative stress *in vivo* using paraquat

To test the efficacy of paraquat dichloride (PQ, 36541-SIGMA) as an inducer of oxidative stress in the brain cortex tissue *in vivo*, PQ (in physiological saline 0.9%) was administered to WT and *Tst*^{-/-} mice by intraperitoneal injection at a concentration of (25 mg/kg body weight). 24 hours later, mice were culled by decapitation and organs were harvested for analysis.

Total ATP, GSH/GSSG and MPST activity measurements

Total ATP content was measured with the CellTiter-Glo 2.0 Luminescent assay (G9241, Promega). GSH/GSSG content and ratio were measured with the Glutathione GSH/GSSG Assay Kit (MAK440, Sigma-Aldrich) following the manufacturer's instructions. MPST enzyme activity was detected with MPST ELISA kit (MBS9912889, Mybiosource) according to manufacturer's instructions. All analyses were performed using freshly extracted total protein from brain cortex samples, and using microplate multimode reader Spark (Tecan). For MPST enzyme activity, absorbance reading were obtained by a Synergy H4 Hybrid Reader (Biotek).

Protein extraction and immunoblotting

Protein extraction from brain cortex samples was performed using 50 mM Tris·HCl pH 7.5, 150 mM NaCl, 1% (v/v) Triton X-100, 0.1% (w/v) SDS. Extracted proteins were centrifuged to remove sample debris. The protein content was determined with the Bicinchoninic acid (BCA) Protein Assay (23227, ThermoFisher Scientific/Pierce). 20 µg of protein was loaded for SDS/PAGE. After blotting, *Trans*-blot Turbo nitrocellulose membranes (Bio-rad) were stained with Ponceau S staining solution for protein loading reference pictures, then cut for different membrane immunostaining, and blocked with blocking buffer (5% (w/v) milk in 1× Tris-buffered saline (TBS), 0.1% (v/v) Tween) for 1 h. Blocked membranes were incubated with primary antibodies in blocking buffer overnight at 4°C, and finally with HRP-conjugated secondary antibodies for 1 h at room temperature. Detection was performed using Clarity Western ECL substrate (Bio-Rad) and Odyssey Fc Imager (LI-COR). Representative immunoblots are shown. Primary antibodies include TST (HPA003643, Atlas Antibodies), MPST (Ab224043, Abcam), GPX4 (SAB4300725, Sigma-Aldrich), OXPHOS complexes (total OXPHOS Rodent WB Antibody Cocktail, AB110413, ABCAM), SOD1 (HPA001401, Sigma-Aldrich), SOD2 (HPA001814, Sigma-Aldrich), KEAP1 (MABS5164, Millipore), NRF2 (AB62352, ABCAM).

Mitochondrial isolation and high-resolution respirometry

Approximate 50 mg of brain cortex from C57BL/6J and *Tst*^{-/-} mice were harvested and mitochondria were isolated utilizing a semi-automated pump-controlled tissue rupture system (0.7 mL/min) (PCC) and syringes (SGE, Trajan© Scientific, Australia) attached to a cell homogenizer (Isobiotech, EMBL Heidelberg, Germany) installed on a pump (ProSense, Oosterhout, NL).⁴¹ The isolated mitochondria were suspended in isolation buffer (250mM sucrose, 20mM HEPES, 3mM EDTA, pH 7.5). The mitochondrial concentration was quantified by Pierce® BCA Protein Assay Kit (Thermo Scientific). All the experiments were conducted on ice.

The mitochondrial respiration parameters were analyzed using an O2K oxygraphy (Oroboros systems, Innsbruck, Austria). 250 µg crude mitochondria from wildtype and *Tst*^{-/-} mouse cortexes were monitored under continuous stirring at 750 rpm in 1 mL MiR05 (0.5 mM EGTA, 3 mM MgCl₂, 60 mM lactobionic acid, 20 mM taurine, 10 mM KH₂PO₄, 20mM HEPES, 110mM D-Sucrose, BSA, 1 g/l essentially fatty acid free, at pH7.4). Oxygen polarography was performed at 37 °C and the oxygen flux per tissue mass (pmol O₂/s/mg) were recorded in real-time using the DatLab5 software. The mitochondrial respiration was assessed with the use of different substrates as follows: 5 mM pyruvate, 2 mM malate and 1.5 µM FCCP to uncouple all the ETC complexes and measure complex I respiration; subsequently, 0.5 µM rotenone was added to inhibit complex I, followed by 10 mM succinate to measure complex II respiration. In addition, 2.5 µM antimycin A was added to prevent both complex I and II mediated respiration due to inhibition of complex III, followed by 2 mM ascorbate combined with 0.5 mM TMPD to measure complex IV-mediated respiration. The measurement sequence was completed by injection of 10 mM sodium dithionite.

Detection of the ROS O₂⁻ and H₂O₂, RNS NO and ONOO⁻, and the RSS H₂S and H₂S_n

Fluorescent molecules were used to measure ROS, specifically O₂⁻ (DHE, D7008, Sigma-Aldrich) and H₂O₂ (DCF-DA, D6883, Sigma-Aldrich), the RNS NO (DAF-2DA AB145283, ABCAM) and ONOO⁻ (DHR123, D1054, Sigma-Aldrich), and the RSS H₂S (7-Azido-4-Methylcoumarin, 802409, Sigma-Aldrich) and H₂S_n (Sulfane Sulfur Probe 4/SSP4, SB10-10, Tebubio) in freshly extracted total proteins from brain cortex samples. This was done by spectrofluorometry using a microplate multimode reader Spark (Tecan).

Total SOD activity, catalase activity, total nitric oxide synthase activity, and total peroxidase activity

Total SOD activity was measured with the Superoxide Dismutase Activity Assay Kit (AB65354, ABCAM), which assesses the inhibition activity of xanthine oxidase by SOD. Catalase activity was assessed with the Catalase Activity Assay Kit (AB83464, ABCAM). Total Nitric Oxide Synthase activity was measured with the Nitric Oxide Synthase (NOS) Activity Assay Kit (MAK407, Sigma-Aldrich). Total peroxidase activity was assessed with the Peroxidase Activity Assay Kit (MAK092, Sigma-Aldrich). Analyses were performed following manufacturer's instructions, with freshly extracted total proteins from brain cortex samples, and using microplate multimode reader Spark (Tecan).

Glutamate content

Glutamate concentration was measured with the Glutamate Assay Kit (AB138883, ABCAM). Analysis was performed following manufacturer's instructions, with freshly extracted total proteins from brain cortex samples, and using microplate multimode reader Spark (Tecan).

RNA isolation, cDNA synthesis and quantitative real-time PCR

RNA was extracted from tissue slices according to the instruction of miRNeasy® Mini Kit (Qiagen, Germany). Total RNA concentration was quantified by Nanodrop® (ND-1000 Spectrophotometer). Equal amount of cDNA synthesis and amplification were performed using Superscript II kit and RANDOM primers (Thermo Fisher Scientific, USA), according to the manufacture's protocol. The cDNA synthesis mix are incubated as follows: 10 mins at 25 °C, 50mins at 42 °C, 15 mins under 70 °C via a Veriti 96 Well Thermal Cycler (Thermo Fisher Scientific). For qRT-PCR, 500 ng of cDNA samples were mixed with SYBR green, forward primers, and reverse primers (Primers used for quantification are shown in Supplementary Table.1). The qPCR was run on LightCycler® 480 II (Roche) and the protocol was as follows: 2 min at 50 °C, 10 min at 95 °C, 40 cycles of 15s 95 °C and 60s 60 °C, and the dissociation curve is run under 15s 95 °C, 15s 60 °C, 15s 95 °C. RT-qPCR data was analyzed on LightCycler 480 software. Expression levels were normalized to GAPDH and displayed as 2-delta Ct.

Thiosulfate measurements in the cortex using LC-MS/MS

Frozen cortex (~25 mg, exact weight recorded) was homogenised in 160 mM EPPS, 16 mM DTPA and monobromobimane, pH 8.0 buffer (300 µL) and acetonitrile (300 µL) with 46 mM monobromobimane (MBBr) (25 µL) for 30 min, room temperature in the dark. The reaction was stopped with ethyl acetate (1 mL), the mixture centrifuged (400 x g for 15 min at 4 °C), and the upper organic layer (~800 µL) passed through a 96-well Filter+ Plate (Biotage, Sweden), along with an MBB-derivatised calibration curve of thiosulfate (0, 0.01, 0.025, 0.05, 0.1, 0.25, 0.5, 1, 2.5, 5, 10, 20, 90 and 100 µM), followed by PLD+ extraction (Biotage, Sweden), reduction to dryness and resuspension in water (100 µL). Samples were analyzed on an I-Class UPLC system (Waters, UK) on a 16 min chromatographic run using an HSS T3 (2.1 x 150 mm; 1.8 mm) column kept at 50°C. MBB-thiosulfate was eluted using water (0.1% formic acid) / methanol (0.1% formic acid) system at 0.4 mL/min from 0-100% methanol over 6 mins, re-equilibrating to 0% methanol at 15 mins. Multiple reaction monitoring parameters for the thiosulfate MBB derivative was: m/z 303.0 → 205.1 on a QTrap 6500+ (AB Sciex) with the following compound settings: declustering potential -65 V, collision exit potential -15 V, collision energy -18 V. Source conditions were -4.5 kV ion spray voltage, 600 °C temperature, gases (GS1=40, GS2=60, Curtain gas= 40 units). Analyst® v1.7 was used for instrument control and data acquisition. Data analysis was carried out using MultiQuant v3.0 (AB Sciex). Linear regression analysis of the peak areas of MBB-thiosulfate was used to calculate the quantity of thiosulfate in each sample, corrected for tissue mass.

Statistics

The data are presented as mean \pm standard deviation (SD). Statistical differences between two conditions were calculated using an unpaired T-test. Statistical differences among more than two conditions were calculated using an one-way ANOVA test. Statistical analysis was performed in GraphPad Prism (version 9.1.0), and repeat numbers are specified in the figure legends.

Journal Pre-proof

Results

***Tst*^{-/-} cerebral cortex exhibits a steady H₂S level and a dysregulated GSH pathway**

To investigate how TST influences the GSH pathway in the brain, we performed studies in the cortical brain region of conventional knock-out mice in which *Tst* deletion (termed *Tst*^{-/-}) has been successfully validated (Fig. 1A). Although the difference in thiosulfate concentrations in the cerebral cortex between wildtype and knockout mice did not reach statistical significance (likely due to small sample size), mean thiosulfate concentration was approx. 3-fold higher in *Tst*^{-/-} mice compared to WT (Fig. 1B). Next, sulfide and polysulfide concentrations were assessed in the cortex. While tissue steady-state levels of both, H₂S and H₂S_n were found to be maintained at similar levels in *Tst*^{-/-} and C57BL/6J mice, mean concentrations were lower for H₂S and higher for polysulfides (Fig. 1B). To investigate whether MPST can compensate for the lack of TST in the brain of the *Tst*^{-/-} mice, we analysed *Mpst* mRNA and protein levels. MPST protein and *Mpst* mRNA was decreased in *Tst*^{-/-} mice (Fig.S1A-D). Interestingly, the MPST activity was not changed, although it tended to be higher in the brain of *Tst*^{-/-} mice compared to C57BL/6J mice (Fig. S1E). To verify whether the GSH pathway was altered in the brain of *Tst*^{-/-} mice, we measured its GSH and GSSG content. GSH was 36% lower in *Tst*^{-/-} mice, while its oxidized form, GSSG, increased 5 times, causing the GSH/GSSG ratio to decline 7.2-fold in *Tst*^{-/-} mice compared to wildtype controls (Fig. 1C). This indicates a redox imbalance secondary to TST enzyme activity loss. Moreover, we found a significant decrease of GPX4 protein (Fig. 1D; Fig. S1F) and the transcriptional level of GR significantly decreased in *Tst*^{-/-} brain cortex (Fig. 4J). Altogether, our results established that expression and activity of TST exert a major effect on both H₂S metabolism and the GSH antioxidant pathway (Fig. 1E).

TST deletion increases ATP in the cerebral cortex and elevates ROS via modulation of OXPHOS activity

Since TST exists almost exclusively in mitochondria and is capable of modulating mitochondrial respiratory complexes, we examined the impact of *Tst* deletion on mitochondrial activity in the cortical brain areas. For that purpose, we performed total ATP measurements generated by OXPHOS *via* luminescence. Total ATP was increased by 22% in the *Tst*^{-/-} mice cortex compared with C57BL/6J mice (Fig. 2A). Next, we measured the protein levels of OXPHOS complexes, and found no significant differences between the two mouse genotypes (Fig. 2B; Fig. S2A). In contrast, basal respiration and complex IV activity represented by oxygen consumption rate were markedly increased in the cortex of *Tst*^{-/-} mice, while no significant differences in Complex I capacity were observed among either mouse genotypes (Fig. 2C). The increase of complex IV activity in the absence of TST indicates an impact of this enzyme on mitochondrial capacity. The function of TST as modifier of GSH to form glutathione persulfide (which is a more efficient ROS scavenger than GSH itself) has been well characterized.^{12,16} We next determined the extent of O₂-derived reactive species formation from OXPHOS. The O₂⁻ level was 10% higher in the cortex of *Tst*^{-/-} mice, while tissue levels of H₂O₂ were 57% higher in the cerebral region of *Tst*^{-/-} mice compared to C57BL/6J control mice

(Fig. 2D).

***Tst* deficiency reprograms the antioxidant defense against endogenous ROS.** To determine whether antioxidant expression/activity changed in the cortex of *Tst*^{-/-} mice, we checked both protein and total enzyme activity levels of superoxide dismutase 1 (SOD1) and superoxide dismutase 2 (SOD2). Both, SOD1 and SOD2 are vital endogenous antioxidant enzymes. The expression of SOD1 and SOD2 proteins was unchanged between the two groups, whereas total SOD enzyme activity showed a 34% decrease in the *Tst*^{-/-} mice (Fig.3A; Fig. S3A). When H₂O₂ is excessively produced, catalase participates in H₂O₂ breakdown. The catalase activity in *Tst*^{-/-} was 7.6% higher relative to that observed in C57BL/6J mice (Fig.3B). In addition, the total peroxidase from *Tst*^{-/-} mice exhibited unaltered activity compared with C57BL/6J mice (Fig.3C). In the context of the RSI framework, we also measured nitric oxide synthase (NOS) activity and RNS (NO and ONOO⁻) production. NOS activity and NO content were similar in the brain cortex of either genotype (Fig.S3B and D), while ONOO⁻ amount in cortexes of *Tst*^{-/-} mice was 10% lower than in C57BL/6J mice (Fig.S3C). In summary, we found decreased SOD enzyme activity, increased catalase activity and unchanged peroxidase activity, when ROS is elevated due to the global loss of TST in murine brain tissue as schematically depicted in Fig. 3D.




Fig. 1. $Tst^{-/-}$ mice brain cortexes display a dysregulated GSH pathway. A-D: Data are expressed as mean \pm SD, student T-test. **A.** *Tst* mRNA and Tst protein levels of $Tst^{-/-}$ (red) and C57BL/6J (black) mice (n=4). **B.** Thiosulfate content in C57BL/6J mice (Black, n=3) and $Tst^{-/-}$ mice brain cortexes (Red, n=4) measured by HPLC-MS/MS. H₂S and H₂S_n concentrations were measured using fluorescent probes, and show that H₂S decreased 5% and H₂S_n increased 5% in the $Tst^{-/-}$ (red) compared with C57BL/6J mice (black) (n=4). **C.** GSH and GSSG contents in mice cortexes in the presence (black) or absence (red) of TST (n=4). **D.** Quantification of immunoblots of cerebral cortical area for GPX4 protein level in $Tst^{-/-}$ (red) compared with C57BL/6J mice (black) (n=3). The original blotting picture is shown in **Fig S1F**. **E.** Brief schematic showing H₂S enzymatic pathway. Sulfide is an endogenously produced gaseous signaling molecule via CBS, CSE and MPST. Polysulfides (H₂S_n) became novel reactive sulfur species, which is derived from H₂S with MPST. TST converts thiosulfate produced via non-enzymatic reactions to sulphite. Then oxidation of sulphite to sulphate is dependent on the glutathione system. GPX4 and GR is the cofactor of conversion between GSH and GSSG.




Fig. 2. $Tst^{-/-}$ mice brain cortexes display metabolism reprogramming. A-D: Data expressed as mean \pm SD, student T-test. **A.** Total ATP level of C57BL/6J (black) and $Tst^{-/-}$ (red) mice cortical area (n=4). **B.** OXPHOS protein levels of two genotypes detected by OXPHOS cocktail antibody (n=3). Quantification is normalized to ponceau stain. Immunoblot is displayed in **Fig. S2**. **C.** Corrected oxygen consumption rate of 250ug crude isolated mitochondria from mice cerebral cortexes with various substrates. Respiratory capacity differences of basal respiration, Complex I, Complex II and Complex IV from C57BL/6J (black) and $Tst^{-/-}$ (red) were analyzed. (n=3 biological samples, independent experiments were repeated at least 3 times). **D.** Relative ROS contents as O_2^- and H_2O_2 of C57BL/6J (black) and $Tst^{-/-}$ (red) (n=4).




Fig. 3. *Tst*^{-/-} mice cerebral cortical regions display reprogrammed antioxidant defense. A-D: Data expressed as mean ± SD, student T-test. **A.** SOD protein expression (n=3) and enzyme activity (n=4) levels of C57BL/6J (black) and *Tst*^{-/-} (red) mice cerebral cortical area. Original immunoblots are presented in **Fig.S3A. 2**. **B.** Catalase enzyme activity measurements in C57BL/6J (black) and *Tst*^{-/-} (red) mice (n=4). **C.** Peroxidase activity in C57BL/6J (black) and *Tst*^{-/-} (red) mice (n=4). **D.** Created with Biorender.com. Brief scheme of response to excessive ROS with or without TST presence. SOD1 and SOD2 are vital endogenous antioxidant elements for H₂O₂ catalyzation from O₂⁻.




Fig. 4. *Tst*^{-/-} mice brain cortexes display dysregulated Nrf2-keap1 signaling expression and increased glutamate content. **A-B, D-G:** Data expressed as mean \pm SD, student T-test. **A.** Expression of *Keap1* mRNA in C57BL/6J (black) (n=4) and *Tst*^{-/-} (red) (n=4) mice levels of cortical area. **B.** Expression of *Nrf2* mRNA in C57BL/6J (black) (n=4) and *Tst*^{-/-} (red) (n=4) mice levels of cortical area. **C.** WB analysis of the two proteins. **D.** Quantification of immunoblots of cerebral cortical area for NRF2 protein level in mice cortexes in the presence from C57BL/6J (black) (n=3) and *Tst*^{-/-} mice (n=3). **E.** Quantification of immunoblots of cerebral cortical area for KEAP1 protein level in mice cortexes in the presence from C57BL/6J (black) (n=3) and *Tst*^{-/-} mice (n=3). **F.** Glutamate content measurements in both genotypes (n=4). **G.** *Tst*^{-/-} mice brain cortexes display

dysregulated ARE genes. P value of each gene is indicated between genotypes. All experiments were conducted with repeat number as 4 per genotype.

***Tst* absence results in decrease of Nrf2-Keap1 pathway expression level and increased glutamate content**

Activation of the Nrf2-Keap1 system results in an antioxidant defense response which is, at least in part, regulated by H₂S and KEAP1 protein level.^{21,22,28} To investigate whether *Tst* deficiency causes a dysregulation of Nrf2 signaling, qRT-PCR was conducted for genes regulated by antioxidant response elements (AREs) under control of NRF2. *Nrf2* and *Keap1* mRNA were significantly decreased in *Tst*^{-/-} cortical brain samples compared with C57BL/6J mice (Fig.4A and B). NRF2 protein showed significant reduction and KEAP1 protein had an increased expression in *Tst*^{-/-} mice brain cortexes when compared to those of control mice (Fig.4C-E). At the same time, different ARE genes including *Hmox1* and *Txn2* mRNA showed relative reduction in the case of *Tst* deficiency, while *Gstm1*, *Txn1*, *Noq1* and *Cat* revealed no significant difference between the two types of mice (Fig.4G).

GCLC, GCLM and GR are important ARE genes for proteins protecting against oxidative stress. All these genes were found to be significantly reduced in the *Tst*^{-/-} brain (Fig.4G). Since both GCLC and GCLM are critical for *de novo* GSH biosynthesis we also measured glutamate content and found a 1.8-fold increase in *Tst*^{-/-} compared to wildtype mice, corroborating the effect of *Tst* deletion on this important antioxidant pathway (Fig.4F).

***Tst*^{-/-} mouse brain displayed deteriorated antioxidant capacity when facing paraquat-induced oxidative stress**

In order to show the effects of TST loss when facing oxidative stress, we analyzed several oxidative stress markers including ROS, RSS, GSH/GSSG contents and GPX4 protein abundance. Under basal TST deficiency condition, ROS showed a significant increase in the *Tst*^{-/-} mouse brains. Upon challenging both genotypes of mice with 25mg/kg paraquat (PQ) for 24hrs, both genotypes of mouse brain cerebral cortices showed significantly increased ROS compared to their basal levels. Cerebral cortex of *Tst*^{-/-} mice showed 1.5-fold increase of H₂O₂ compared to those of C57BL/6J mice (Fig.5A). As for reactive sulfur species, brain cortexes of *Tst*^{-/-} mice showed similar amounts of H₂S in the cortical brain area compared to C57BL/6J mice. Upon PQ challenging, there was no difference in both H₂S and H₂S_n intensity between the two genotypes (Fig.5B).

The parameters relevant to the GSH content were also assessed. Following paraquat stimulation, both mouse genotypes had significantly higher GSH content compared to their basic levels in brain tissue. However, GSH amount generated in the brain cortex of *Tst*^{-/-} mice was less than that of C57BL/6J mice. In contrast to the accumulation of GSSG in basal cerebral cortical area of *Tst*^{-/-} mice, PQ-treated *Tst*^{-/-} mice also had relatively lower GSSG compared to C57BL/6J mice. Therefore, in the cerebral cortex of PQ-treated mice the ratio of GSH/GSSG showed a significant increase when compared with C57BL/6J mice (Fig.5C). Furthermore, we tested GPX4 protein expression and found that GPX4 had higher levels in PQ injected animals in response to acute redox damage in both mouse genotypes. In the mouse brain of *Tst*^{-/-} mice,

GPX4 protein level was slightly lower than in C57BL/6J mice (Fig.5D). In summary, this additional challenge experiment showed an overall deteriorated antioxidant capacity in murine cerebral cortices in conditions of global TST deficiency.




Fig. 5 PQ challenge exacerbated *Tst*^{-/-} mice brain cortexes disrupted antioxidant capacity. A-D: Data are expressed as mean \pm SD, one-way ANOVA test. **A.** Relative ROS contents as O₂⁻ and H₂O₂ of mice brain cortexes in the presence (hatched pattern) or absence (no pattern) of 25mg/kg 24hrs from C57BL/6J (black) (n=4-5) and *Tst*^{-/-} mice (n=4). **B.** Relative H₂S and H₂S_n concentrations of mice brain cortexes in the presence (hatched pattern) or absence (no pattern) of 25mg/kg 24hrs from C57BL/6J (black) (n=4-5) and *Tst*^{-/-} mice (n=4). **C.** GSH and GSSG contents in mice cortexes in the presence (hatched pattern) or absence (no pattern) of 25mg/kg 24hrs from C57BL/6J (black) (n=4-5) and *Tst*^{-/-} mice (n=4). **D.** Quantification of immunoblots of cerebral cortical area for GPX4 protein level in mice cortexes in the presence (hatched pattern) or absence (no pattern) of 25mg/kg 24hrs from C57BL/6J (black) (n=3) and *Tst*^{-/-} mice (n=3). The immunoblots are shown in **Fig S4**.

Discussion

TST plays an important role in the oxidation of the endogenous gasotransmitter hydrogen sulfide, complementing the action of sulfide quinone oxidoreductase (SQOR) and persulfide dioxygenase (ETHE1/PDO).^{9,42} Whether or not TST also contributes to antioxidant protection in the brain has not been investigated before. In *Tst*^{-/-} mice, circulating thiosulfate and sulfide level were markedly elevated, while cerebral cortex displayed similar steady-state levels of sulfide and thiosulfate, as observed in the liver.⁴³ This indicates that the response to *Tst* knock-down is organ specific in this chronic but viable sulfide elevation model. Starting from organ specificity, the free thiol antioxidant system of brain and liver revealed opposite responses in terms of GSH content. In *Tst*^{-/-} mice, GSH was decreased in the cerebral cortex, while there was no difference in GSH content between the two genotypes in the liver.⁴³ Since NADPH/H⁺ dependent glutathione reductase (GR) is important for the recycling of GSSG to GSH⁴⁴, its down-regulation in the brain of *Tst*^{-/-} mice explains the alterations in GSSG levels observed. In support of an activation role of TST on ROS scavengers such as GSH,¹⁵ *Tst*^{-/-} mice showed a lower GSH/GSSG ratio and higher ROS levels, indicating a redox signaling imbalance along with lower GPX4 and GR expression. Meanwhile, the *Mpst* transcription level of brain and liver in *Tst*^{-/-} mice were reduced. In contrast, the liver functions as a major detoxification organ, and despite lower mRNA for *Mpst* protein levels were upregulated, perhaps for compensatorily enhanced sulfide removal,⁴³ while MPST protein significantly decreased in the brain cortex and in mitochondria in *Tst*^{-/-} mice.

TST interacts with iron-sulfur cluster proteins including Complex I (NADH dehydrogenase), Complex II (succinate dehydrogenase) and Complex III (Cytochrome bc1) of the mitochondrial respiratory chain in oxidative pathway.⁴⁵ TST directly interacts with Complex I via sulfide transfer.⁶ In addition, TST transfers sulfane sulfur to Complex II and modifies its iron-sulfur structure via alteration of its phosphorylation status. Those functions led us to explore the metabolic phenotypes of the brain tissues of C57BL/6J and *Tst*^{-/-} mice. The unexpected finding of mitochondrial activity remodeling was supported by several observations, including unchanged OXPHOS protein expression, increased basal respiration capacity and Complex IV activity in the presence of ascorbate and TMPD. However, the cerebral cortex of *Tst*^{-/-} mice produced more ATP, and higher amounts of ROS compared to C57BL/6J control mice. ATP, the universal energy currency, is mainly synthesized in mitochondria by OXPHOS. The representative role of Complex IV as regulatory center of OXPHOS limits the rate of respiratory chain reaction.⁴⁶ Hence, our finding of enhanced Complex IV activity is in line with the observation of enhanced ATP levels in *Tst*^{-/-} mice.

ROS are generated during the incomplete reduction of $O_2^{47,48}$. In mitochondria, the major sources of $O_2^{\bullet -}$ are the OXPHOS Complexes I and III.^{20,49} $O_2^{\bullet -}$ will be further processed quickly to H_2O_2 via SOD1 and SOD2.⁵⁰ In the absence of TST, the significant inhibition of total SOD activity can be explained by excessive H_2O_2 in the brain cortex, which could possibly slow down H_2O_2 production as oxidative distress response. The metabolism of H_2O_2 to H_2O is catalyzed by catalase, glutathione peroxidase, and peroxiredoxin 1.⁵¹ Compared to $O_2^{\bullet -}$, the steadier feature of H_2O_2 allows for mitochondrial modulation of the cellular antioxidant system.⁵² The observed alterations in ROS metabolizing enzymes indicate the importance of TST in overall antioxidant capacity to protect the brain.

Our study of the Nrf2 pathway in *Tst*^{-/-} mice highlights a potent interaction between Nrf2 signaling and the GSH antioxidant system⁵³⁻⁵⁶. Importantly, Nrf2 is a major transcriptional factor regulating multiple processes related to GSH synthesis and regeneration.^{57,58} Beyond GSH content change, the transcriptional and protein levels of Nrf2 appear to be lower and its intracellular inhibitor Keap1 had higher protein expression in the brains of *Tst*^{-/-} mice. A possible reason for the fact that TST deletion causes disrupted Nrf2-Keap1 signaling, might be found in the abnormal cysteine metabolism within the redox sensor protein KEAP1. TST's main function is sulfide oxidation as cysteine metabolism downstream, and KEAP1 protein contains 25 cysteines within mouse homologue, thus accumulated KEAP1 were detected in the brain of *Tst*^{-/-} mice, consequently lower amount of Nrf2 protein was released from KEAP1 ubiquitin E3 ligase adapter, which further influenced the ARE gene transcription process.^{22,59,60} In light with reduced transcription of *Keap1*, we found an increased KEAP1 protein level, and we suggest a post-transcriptional cellular redox-sensing mechanism as operative.

Similar reduction in Nrf2 activation in *Tst*^{-/-} mice using transcription factor binding site (TFBS) enrichment analysis were found earlier.⁴³ Additionally, glutamate-cysteine ligase catalytic (GCLC) and modifier (GCLM) subunits together catalyze the rate-limiting step in GSH biosynthesis from glutamate under conditions of oxidative distress.²⁶ *Nrf2*^{-/-} mice exhibited a decrease in GSH content and GCL levels,⁶¹ which supported our results on the brains of *Tst*^{-/-} mice. The decrease in Nrf2 mRNA and protein translate into a reduced function of the GSH antioxidant system. Moreover, the decrease in Nrf2-mediated free thiol antioxidants influences various cellular processes, ranging from synthesis, oxidation and reduction of GSH. Among the heterogeneity of brain cell types, the expression of *Tst* mRNA is known to be highest in astrocytes (<https://www.proteinatlas.org/ENSG00000128311-TST/single+cell+type/brain>). Astrocytes comprise the majority of glial cells in the mammalian central nervous system (CNS) and serve as major provider of GSH for neurons.^{62,63} Astrocytes contribute to neuronal activity, and their interaction play distinct, cooperative roles in neuronal redox homeostasis. Astrocytes could initiate and promote an increase in gene expression of Nrf2-mediated antioxidant pathway in response to mild oxidative stress, which results in the synthesis of GSH using glutamate.⁶⁴ When astrocytes function abnormally, glutamate, which is an excitatory neurotransmitter, will accumulate to an excitotoxic level and contribute to harmful effects on neurons, eventually causing impaired brain function.⁶⁵ Thus, any deficiency in astrocyte capacity to produce GSH (or reduce GSSG) will render neurons particularly vulnerable to conditions or insults known to be associated with enhanced oxidative stress.⁶⁶⁻⁷⁰ Both oxidative stress and elevated glutamate contribute and are associated with brain pathologies.⁶⁵ The decreased Nrf2 expression, reduced

GSH and increased glutamate content, together with an altered ROS landscape in the CNS, secondary to a loss of TST are therefore key observations of the present study.

Relevant to Nrf2 signaling, another important thiol-dependent antioxidant system is the thioredoxin (TXN) system.⁷¹ TXN2 is a mitochondria-specific protein, whereas TXN1 is found in the cytosol, both of them operate using NADPH as cofactor.⁷² Down-regulated *Txn2* mRNA level observed in the cerebral cortex of *Tst*^{-/-} mice, could be explained by loss of TST's function to interact with mitochondrial NADH dehydrogenase (Complex I) and other iron-cluster proteins.⁶ Our findings provide a better understanding of the role of TST in linking Nrf2-keap1 signaling and thiol antioxidant pathways.

When challenging mice with paraquat-induced oxidative stress, the ROS production in *Tst*^{-/-} mouse brain was dramatically increased due to the hampered antioxidant capacity, indicating the importance of the interaction of TST with the antioxidant pathway. The nature of GSH is to provide a protective effect on free radical scavenging and H₂O₂ detoxification⁷³ therefore, after stimulation, increased GSH content was detected in both genotypes, although *Tst*^{-/-} mouse brain had lower GSH levels than control mice. Unlike the basal condition, PQ-treated mice biosynthesized less GSSG when the brain tissue of *Tst*^{-/-} mice was compared to the WT animals. PQ injection also induced an increase in GPX4 protein expression in response to excessive oxidative distress. GPX4 protein in *Tst*^{-/-} mice was slightly lower than in C57BL/6J mice. In summary, this additional challenge experiment showed an overall deteriorated antioxidant capacity in murine cerebral cortex under the model of global TST deficiency.

In the context of brain function, both mitochondria and NRF2 functions are particularly important because the brain is highly susceptible to oxidative distress due to its high metabolic rate and relatively low levels of antioxidants compared to other tissues⁷⁴. During oxidative stress, structural and functional damage occurs to neurons and disrupts the normal actions of neurotransmitters, leading to altered signaling from neurons. This can lead to impaired cognitive function, affect behavior, and contribute to the development and progression of neurological and neurodegeneration diseases⁷⁵. Taken together, our data unmask several tissue-specific mechanisms linked to TST function that are related to mitochondrial and Nrf2-Keap1 signaled antioxidant functions, and the role of TST in maintaining the health and function of brain tissues.

Conclusion

Our study on mouse cerebral cortex revealed that TST deficiency promotes a dysregulation of the reactive species interactome through both generation of ROS and RSS, coupled with mitochondrial OXPHOS remodeling. In addition, TST deficiency elicits the dysregulation of thiol-dependent antioxidant systems with altered GSH levels and aberrant Nrf2 pathway expression. Moreover, the absence of TST is characterized by a deteriorated antioxidant buffering ability in the murine brain cortex when challenged with a redox cyler as demonstrated by paraquat *in vivo*.

Funding

Y.L. was supported by China Scholarship Council (Grant No.: 202008520033). Z. M. Al-Dahmani was supported by the Islamic Development Bank. A.M.D. was supported by a Rosalind Franklin Fellowship co-funded by the European Union and the University of Groningen. S.M. was supported by a British Heart Foundation 4Y PhD Scholarship (FS/4yPhD/F/20/34126). L.C was supported by the CNRS.

Contribution Statement

YL, AM and HvG designed and conceptualized the study, wrote and revised the manuscript. YL conducted parts of research, LC designed and performed parts of study, edited the manuscript. YL and LC analyzed the data. ZMM conducted parts of experiments. SM, NZMH and NMM performed experiments on the animal model, provided samples, and edited the manuscript. MG and MF revised the manuscript. AM and HvG supervised this study.

Declaration of competing interest

The authors hereby declare that there is no competing financial interests or personal relationships that could have influenced the work in this paper.

Data availability

Data will be made available on request.

Acknowledgements

All authors greatly appreciate the technical assistance from Wierd Kooistra and Albert Gerding on experimental training. LC-MS/MS data were obtained at the University of Edinburgh, Edinburgh Clinical Research Facility, Mass Spectrometry Core, RRID:SCR_021833 using the AB SCIEX QTRAP 6500 plus system, RRID:SCR_021831. We acknowledge Joanna Simpson for expertise and Scott Denham for operation of the LC-MS/MS instrumentation. YL and HvG are members of COST (European Cooperation in Science and Technology) Action CA20121 (Bench to bedside transition for pharmacological regulation of NRF2 in noncommunicable diseases, BenBedPhar).

Appendix A. Supplementary data

Supplementary data to this article can be found online.

Reference:

1. Lang K. Die Rhodanbildung Im Tierkorper [Thiocyanogen in the Bodies of Animals]. *Biochem Z.* 1933;259:243-256.
2. Hall AH, Saiers J, Baud F. Which cyanide antidote? *Crit Rev Toxicol.* 2009;39(7):541-552. doi:10.1080/10408440802304944
3. Nagahara N, Okazaki T, Nishinoz T. Cytosolic Mercaptopyruvate Sulfurtransferase Is Evolutionarily Related to Mitochondrial Rhodanese: Striking similarity in active site amino acid sequence and the increase in the mercaptopyruvate sulfurtransferase activity of rhodanese by site-directed mutagenesis. *J Biol Chem.* 1995;270(27):16230-16235.
4. Nagahara N, Nishino T. Mercaptopyruvate sulfurtransferase role of amino acid residues in the active site of rat liver mercaptopyruvate sulfurtransferase: cDNA cloning, overexpression, and site-directed mutagenesis. *J Biol Chem.* 1996;271:27395-27401.
5. Wang R. Physiological implications of hydrogen sulfide: A whiff exploration that blossomed. *Physiol Rev.* 2012;92(2):791-896. doi:10.1152/physrev.00017.2011
6. Pagani S, Galante YM. Interaction of rhodanese with mitochondrial NADH dehydrogenase. *Biochim Biophys Acta (BBA)- Protein Struct Mol Enzymol.* 1983;742(2):278-284. doi:10.1016/0167-4838(83)90312-6
7. F Bonomi, S Pagani, P Cerletti, C Cannella. Rhodanese-mediated sulfur transfer to succinate dehydrogenase. *Eur J Biochem.* 1977;72(1):17-24. doi:10.1111/j.1432-1033.1977.tb11219.x
8. Nandi DL, Horowitz PM, Westley J. Rhodanese as a thioredoxin oxidase. *Int J Biochem Cell Biol.* 2000;32(4):465-473. doi:10.1016/S1357-2725(99)00035-7
9. Hildebrandt TM, Grieshaber MK. Three enzymatic activities catalyze the oxidation of sulfide to thiosulfate in mammalian and invertebrate mitochondria. *FEBS J.* 2008;275(13):3352-3361. doi:10.1111/j.1742-4658.2008.06482.x
10. Villarejol M, Westley J. Rhodanese-catalyzed reduction of thiosulfate by reduced lipoic acid. *J Biol Chem.* 1963;238:1185-1186. doi:10.1016/s0021-9258(18)81283-2
11. Iciek M, Kowalczyk-Pachel D, Bilska-Wilkosz A, Kwiecién I, Górny M, Wødek L. S-sulfhydration as a cellular redox regulation. *Biosci Rep.* 2016;36(2). doi:10.1042/BSR20150147
12. Libiad M, Sriraman A, Banerjee R. Polymorphic variants of human rhodanese exhibit differences in thermal stability and sulfur transfer kinetics. *J Biol Chem.* 2015;290(39):23579-23588. doi:10.1074/jbc.M115.675694
13. Xu S, Zhang X, Liu C, et al. Role of Mitochondria in Neurodegenerative Diseases: From an Epigenetic Perspective. *Front Cell Dev Biol.* 2021;9. doi:10.3389/fcell.2021.688789
14. Wang Y, Xu E, Musich PR, Lin F. Mitochondrial dysfunction in neurodegenerative diseases and the potential countermeasure. *CNS Neurosci Ther.* 2019;25(7):816-824. doi:10.1111/cns.13116

15. Kruithof PD, Lunev S, Aguilar Lozano SP, et al. Unraveling the role of thiosulfate sulfurtransferase in metabolic diseases. *Biochim Biophys Acta (BBA)-Mol Basis Dis.* 2020;1866(6). doi:10.1016/j.bbadis.2020.165716
16. Libiad M, Motl N, Akey DL, et al. Thiosulfate sulfurtransferase-like domain-containing 1 protein interacts with thioredoxin. *J Biol Chem.* 2018;293(8):2675-2686. doi:10.1074/jbc.RA117.000826
17. Sies H. Findings in redox biology: From H₂O₂ to oxidative stress. *Journal of Biological Chemistry.* 2020;295(39). doi:10.1074/jbc.X120.015651
18. Jones DP. Redefining Oxidative Stress. *Antioxid Redox Signal.* 2006;8(9-10):1865-1879. doi:10.1089/ars.2006.8.1865
19. Cortese-Krott MM, Koning A, Kuhnle GGC, et al. The reactive species interactome: evolutionary emergence, biological significance, and opportunities for redox metabolomics and personalized Medicine. *Antioxid Redox Signal.* 2017;27(10):684-712. doi:10.1089/ars.2017.7083
20. Malard E, Valable S, Bernaudin M, Pérès E, Chatre L. The reactive species interactome in the brain. *Antioxid Redox Signal.* 2021;35(14):1176-1206. doi:10.1089/ars.2020.8238
21. Suzuki T, Yamamoto M. Molecular basis of the Keap1–Nrf2 system. *Free Radic Biol Med.* 2015;88:93-100. doi:10.1016/j.freeradbiomed.2015.06.006
22. Saito R, Suzuki T, Hiramoto K, et al. Characterizations of Three Major Cysteine Sensors of Keap1 in Stress Response. *Mol Cell Biol.* 2016;36(2):271-284. doi:10.1128/MCB.00868-15
23. Raghunath A, Sundarraj K, Nagarajan R, et al. Antioxidant response elements: Discovery, classes, regulation and potential applications. *Redox Biol.* 2018;17:297-314. doi:10.1016/j.redox.2018.05.002
24. Xie L, Gu Y, Wen M, et al. Hydrogen sulfide induces Keap1 S-sulfhydration and suppresses diabetes-accelerated atherosclerosis via Nrf2 activation. *Diabetes.* 2016;65(10):3171-3184. doi:10.2337/db16-0020
25. Cuadrado A, Manda G, Hassan A, et al. Transcription factor NRF2 as a therapeutic target for chronic diseases: A systems medicine approach. *Pharmacol Rev.* 2018;70(2):348-383. doi:10.1124/pr.117.014753
26. Abdul-Aziz A, Macewan DJ, Bowles KM, Rushworth SA. Oxidative stress responses and NRF2 in human leukaemia. *Oxid Med Cell Longev.* 2015;2015. doi:10.1155/2015/454659
27. Yang G, Zhao K, Ju Y, et al. Hydrogen sulfide protects against cellular senescence via s-sulfhydration of keap1 and activation of Nrf2. *Antioxid Redox Signal.* 2013;18(15):1906-1919. doi:10.1089/ars.2012.4645
28. Zhao S, Song T, Gu Y, et al. Hydrogen Sulfide Alleviates Liver Injury Through the S-Sulfhydrated-Kelch-Like ECH-Associated Protein 1/Nuclear Erythroid 2–Related

- Factor 2/Low-Density Lipoprotein Receptor–Related Protein 1 Pathway. *Hepatology*. 2021;73(1):282-302. doi:10.1002/hep.31247
29. Morton NM, Beltram J, Carter RN, et al. Genetic identification of thiosulfate sulfurtransferase as an adipocyte-expressed antidiabetic target in mice selected for leanness. *Nat Med*. 2016;22(7):771-779. doi:10.1038/nm.4115
 30. Levine J, Ellis CJ, Furne JK, Springfield J, Levitt MD. Fecal hydrogen sulfide production in ulcerative colitis. *Am J Gastroenterol*. 1998;93(1):83-87. doi:10.1111/j.1572-0241.1998.083_c.x
 31. Picton R, Eggo MC, Merrill GA, Langman MJS, Singh S. Mucosal protection against sulphide: Importance of the enzyme rhodanese. *Gut*. 2002;50(2):201-205. doi:10.1136/gut.50.2.201
 32. Kaczor-Kamińska M, Kaminski K, Wróbel M. The expression and activity of rhodanese, 3-mercaptopyruvate sulfurtransferase, cystathionine γ -lyase in the most frequently chosen cellular research models. *Biomolecules*. 2021;11(12). doi:10.3390/biom11121859
 33. Taniguchi E, Matsunami M, Kimura T, et al. Rhodanese, but not cystathionine- γ -lyase, is associated with dextran sulfate sodium-evoked colitis in mice: A sign of impaired colonic sulfide detoxification? *Toxicology*. 2009;264(1-2):96-103. doi:10.1016/j.tox.2009.07.018
 34. Poole CJ, Kind PR. Deficiency of thiosulphate sulphurtransferase (rhodanese) in Leber's hereditary optic neuropathy. *BMJ*. 1986;292(6530):1229-1230. doi:10.1136/bmj.292.6530.1229
 35. Cogley JN, Fiorello ML, Bailey DM. 13 reasons why the brain is susceptible to oxidative stress. *Redox Biol*. 2018;15:490-503. doi:10.1016/j.redox.2018.01.008
 36. Vandresen-Filho S, Martins WC, Bertoldo DB, Mancini G, De Bem AF, Tasca CI. Cerebral cortex, hippocampus, striatum and cerebellum show differential susceptibility to quinolinic acid-induced oxidative stress. *Neurol Sci*. 2015;36(8):1449-1456. doi:10.1007/s10072-015-2180-7
 37. Trombetta-Lima M, Sabogal-Guáqueta AM, Dolga AM. Mitochondrial dysfunction in neurodegenerative diseases: A focus on iPSC-derived neuronal models. *Cell Calcium*. 2021;94. doi:10.1016/j.ceca.2021.102362
 38. Sbdio JI, Snyder SH, Paul BD. Redox mechanisms in neurodegeneration: From disease outcomes to therapeutic opportunities. *Antioxid Redox Signal*. 2019;30(11):1450-1499. doi:10.1089/ars.2017.7321
 39. Marmolejo-Garza A, Medeiros-Furquim T, Rao R, Eggen BJJ, Boddeke E, Dolga AM. Transcriptomic and epigenomic landscapes of Alzheimer's disease evidence mitochondrial-related pathways. *Biochim Biophys Acta Mol Cell Res*. 2022;1869(10). doi:10.1016/j.bbamcr.2022.119326

40. Rink C, Khanna S. Significance of brain tissue oxygenation and the arachidonic acid cascade in stroke. *Antioxid Redox Signal* . 2011;14(10):1889-1903. doi:10.1089/ars.2010.3474
41. Honrath B, Matschke L, Meyer T, et al. SK2 channels regulate mitochondrial respiration and mitochondrial Ca²⁺ uptake. *Cell Death Differ*. 2017;24(5):761-773. doi:10.1038/cdd.2017.2
42. Libiad M, Yadav PK, Vitvitsky V, Martinov M, Banerjee R. Organization of the human mitochondrial hydrogen sulfide oxidation pathway. *J Biol Chem* . 2014;289(45):30901-30910. doi:10.1074/jbc.M114.602664
43. Carter RN, Gibbins MTG, Barrios-Llerena ME, et al. The hepatic compensatory response to elevated systemic sulfide promotes diabetes. *Cell Rep* . 2021;37(6):109958. doi:10.1016/j.celrep.2021.109958
44. Capelletti MM, Manceau H, Puy H, Peoc'h K. Ferroptosis in liver diseases: An overview. *Int J Mol Sci* . 2020;21(14):4908. doi:10.3390/ijms21144908
45. Rydz L, Wróbel M, Jurkowska H. Sulfur administration in fe-s cluster homeostasis. *Antioxidants*. 2021;10(11). doi:10.3390/antiox10111738
46. Kadenbach B. Complex IV – The regulatory center of mitochondrial oxidative phosphorylation. *Mitochondrion*. 2021;58:296-302. doi:10.1016/j.mito.2020.10.004
47. Bourgonje AR, Feelisch M, Faber KN, Pasch A, Dijkstra G, van Goor H. Oxidative stress and redox-modulating therapeutics in Inflammatory bowel disease. *Trends Mol Med* . 2020;26(11):1034-1046. doi:10.1016/j.molmed.2020.06.006
48. Sies H, Jones DP. Reactive oxygen species (ROS) as pleiotropic physiological signalling agents. *Nat Rev Mol Cell Biol* . 2020;21(7):363-383. doi:10.1038/s41580-020-0230-3
49. Snezhkina A V., Kudryavtseva A V., Kardymon OL, et al. ROS generation and antioxidant defense systems in normal and malignant cells. *Oxid Med Cell Longev*. 2020;2019. doi:10.1155/2019/6175804
50. Zhao RZ, Jiang S, Zhang L, Yu Z Bin. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *Int J Mol Med* . 2019;44(1):3-15. doi:10.3892/ijmm.2019.4188
51. Pannala VR, Dash RK. Mechanistic characterization of the thioredoxin system in the removal of hydrogen peroxide. *Free Radic Biol Med* . 2015;78:42-55. doi:10.1016/j.freeradbiomed.2014.10.508
52. Nolfi-Donagan D, Braganza A, Shiva S. Mitochondrial electron transport chain: Oxidative phosphorylation, oxidant production, and methods of measurement. *Redox Biol* . 2020;37. doi:10.1016/j.redox.2020.101674
53. Corsello T, Komaravelli N, Casola A. Role of hydrogen sulfide in nrf2-and sirtuin-dependent maintenance of cellular redox balance. *Antioxidants*. 2018;7(10). doi:10.3390/antiox7100129

54. Ryoo I geun, Kwak MK. Regulatory crosstalk between the oxidative stress-related transcription factor Nfe2l2/Nrf2 and mitochondria. *Toxicol Appl Pharmacol*. 2018;359:24-33. doi:10.1016/J.TAAP.2018.09.014
55. Song X, Long D. Nrf2 and ferroptosis: A new research direction for neurodegenerative diseases. *Front Neurosci*. 2020;14. doi:10.3389/fnins.2020.00267
56. Dodson M, Castro-Portuguez R, Zhang DD. NRF2 plays a critical role in mitigating lipid peroxidation and ferroptosis. *Redox Biol*. 2019;23. doi:10.1016/j.redox.2019.101107
57. Yang Y, Dieter MZ, Chen Y, Shertzer HG, Nebert DW, Dalton TP. Initial characterization of the glutamate-cysteine ligase modifier subunit Gclm(-/-) knockout mouse: Novel model system for a severely compromised oxidative stress response. *J Biol Chem*. 2002;277(51):49446-49452. doi:10.1074/jbc.M209372200
58. Chen Y, Shertzer HG, Schneider SN, Nebert DW, Dalton TP. Glutamate cysteine ligase catalysis: Dependence on ATP and modifier subunit for regulation of tissue glutathione levels. *J Biol Chem*. 2005;280(40):33766-33774. doi:10.1074/jbc.M504604200
59. Dinkova-Kostova AT, Kostov R V., Canning P. Keap1, the cysteine-based mammalian intracellular sensor for electrophiles and oxidants. *Arch Biochem Biophys*. 2017;617:84-93. doi:10.1016/j.abb.2016.08.005
60. Dayalan Naidu S, Muramatsu A, Saito R, et al. C151 in KEAP1 is the main cysteine sensor for the cyanoenone class of NRF2 activators, irrespective of molecular size or shape. *Sci Rep*. 2018;8(1). doi:10.1038/s41598-018-26269-9
61. Hoang YD, Nakamura BN, Chan JY, Luderer U. Decreased ovarian glutathione and glutamate cysteine ligase subunit protein levels in Nrf2 knockout mice. *Biol Reprod*. 2008;78(Suppl_1):116-116. doi:10.1093/biolreprod/78.s1.116a
62. Sjöstedt E, Zhong W, Fagerberg L, et al. An atlas of the protein-coding genes in the human, pig, and mouse brain. *Science (1979)*. 2020;367(6482). doi:10.1126/science.aay4106
63. Human Protein Atlas proteintatlas.org.
64. Baxter PS, Hardingham GE. Adaptive regulation of the brain's antioxidant defences by neurons and astrocytes. *Free Radic Biol Med*. 2016;100:147-152. doi:10.1016/j.freeradbiomed.2016.06.027
65. Maher P, van Leyen K, Dey PN, Honrath B, Dolga A, Methner A. The role of Ca²⁺ in cell death caused by oxidative glutamate toxicity and ferroptosis. *Cell Calcium*. 2018;70:47-55. doi:10.1016/j.ceca.2017.05.007
66. Schousboe A, Scafidi S, Bak LK, Waagepetersen HS, McKenna MC. Glutamate Metabolism in the Brain Focusing on Astrocytes. *Adv Neurobiol*. Published online 2014:13-30. doi:10.1007/978-3-319-08894-5_2
67. McBean GJ. Cysteine, glutathione, and thiol redox balance in astrocytes. *Antioxidants (Basel)*. 2017;6(3). doi:10.3390/antiox6030062

68. Raps SP, Lai JCK, Hertz L, Cooper AJL. Glutathione is present in high concentrations in cultured astrocytes but not in cultured neurons. *Brain Res.* 1989;493(2):398-401. doi:10.1016/0006-8993(89)91178-5
69. Makar TK, Nedergaard M, Preuss A, Gelbard AS, Perumal AS, Cooper AJL. Vitamin E, Ascorbate, Glutathione, Glutathione Disulfide, and Enzymes of Glutathione Metabolism in Cultures of Chick Astrocytes and Neurons: Evidence that Astrocytes Play an Important Role in Antioxidative Processes in the Brain. *J Neurochem.* 2008;62(1):45-53. doi:10.1046/j.1471-4159.1994.62010045.x
70. Yudkoff M, Pleasure D, Cregar L, et al. Glutathione turnover in cultured astrocytes: studies with [¹⁵N]glutamate. *J Neurochem.* 1990;55(1):137-145. doi:10.1111/j.1471-4159.1990.tb08831.x
71. Masutani H, Ueda S, Yodoi J. The thioredoxin system in retroviral infection and apoptosis. *Cell Death Differ.* 2005;12(S1):991-998. doi:10.1038/sj.cdd.4401625
72. Tang H, Kim M, Lee M, et al. Overexpression of thioredoxin-2 attenuates age-related muscle loss by suppressing mitochondrial oxidative stress and apoptosis. *JCSM Rapid Commun.* 2022;5(1):130-145. doi:10.1002/rco2.57
73. Owen JB, Butterfield DA. Measurement of Oxidized/Reduced Glutathione Ratio. In: *Methods Mol Biol.* ; 2010:269-277. doi:10.1007/978-1-60761-756-3_18
74. Bellissimo MI, Amado D, Abdalla DSP, Ferreira EC, Cavalleiro EA, da Graça Naffah-Mazzacoratti M. Superoxide dismutase, glutathione peroxidase activities and the hydroperoxide concentration are modified in the hippocampus of epileptic rats. *Epilepsy Res.* 2001;46(2):121-128. doi:10.1016/S0920-1211(01)00269-8
75. Singh A, Kukreti R, Saso L, Kukreti S. Oxidative Stress: A Key Modulator in Neurodegenerative Diseases. *Molecules.* 2019;24(8):1583. doi:10.3390/molecules24081583

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof